Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

SKU: 9780134296074-TEST-BANK

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

1) Addition and subtraction: If $\vec{A} - \vec{B} = 0$, then the vectors $\vec{A}$ and $\vec{B}$ have equal magnitudes and are directed in the opposite directions from each other.

A) True

B) False

Answer: B

2) Addition and subtraction: Under what condition is $|\vec{A} - \vec{B}| = A + B$?

A) Vectors $\vec{A}$ and $\vec{B}$ are in perpendicular directions.

B) Vectors $\vec{A}$ and $\vec{B}$ are in opposite directions.

C) The magnitude of vector $\vec{B}$ is zero.

D) Vectors $\vec{A}$ and $\vec{B}$ are in the same direction.

E) The statement is never true.

Answer: B

3) Addition and subtraction: If $A > B$, under what condition is $|\vec{A} - \vec{B}| = A - B$?

A) Vectors $\vec{A}$ and $\vec{B}$ re in perpendicular directions.

B) Vectors $\vec{A}$ and $\vec{B}$ are in the same direction.

C) The statement is never true.

D) Vectors $\vec{A}$ and $\vec{B}$ are in opposite directions.

E) The statement is always true.


Answer: B

TBEXAM.COM

TBEXAM.COM

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

4) Addition and subtraction: For the vectors shown in the figure, express vector $\vec{S}$ in terms of vectors $\vec{M}$ and $\vec{N}$.



Answer: $\vec{S} = \vec{M} - \vec{N}$

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

5) Components: Which of the following is an accurate statement?
A) The magnitude of a vector is independent of the coordinate system used.
B) Rotating a vector about an axis passing through the tip of the vector does not change the vector.
C) The magnitude of a vector can be zero even though one of its components is not zero.
D) It is possible to add a scalar quantity to a vector.
E) Even though two vectors have unequal magnitudes, it is possible that their vector sum is zero.
Answer: A

6) Components: The magnitude of a vector can never be less than the magnitude of one of its components.
A) True
B) False
Answer: A

7) Components: If the magnitude of vector $\vec{A}$ is less than the magnitude of vector $\vec{B}$, then the x component of $\vec{A}$ is less than the x component of $\vec{B}$.
A) True
B) False
Answer: B

8) Components: If the eastward component of vector $\vec{A}$ is equal to the westward component of vector $\vec{B}$ and their northward components are equal. Which one of the following statements about these two vectors is correct?
TBEXAM.COM
A) Vector $\vec{A}$ is perpendicular to vector $\vec{B}$.
B) Vectors $\vec{A}$ and $\vec{B}$ point in opposite directions.
C) The magnitude of vector $\vec{A}$ is twice the magnitude of vector $\vec{B}$.
D) Vector $\vec{A}$ is parallel to vector $\vec{B}$.
E) The magnitude of vector $\vec{A}$ is equal to the magnitude of vector $\vec{B}$.
Answer: E

9) Unit vectors: If all the components of a vector are equal to 1, then that vector is a unit vector.
A) True
B) False
Answer: B

10) Scalar (dot) product: If the dot product of two nonzero vectors is zero, the vectors must be perpendicular to each other.
A) True
B) False
Answer: A

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

11) Scalar (dot) product: If two nonzero vectors point in the same direction, their dot product must be zero.
A) True
B) False
Answer: B

12) Scalar (dot) product: The value of the dot product of two vectors depends on the particular coordinate system being used.
A) True
B) False
Answer: B

13) Vector (cross) product: If two vectors are perpendicular to each other, their cross product must be zero.
A) True
B) False
Answer: B

14) Vector (cross) product: If two vectors point in opposite directions, their cross product must be zero.
A) True
B) False
Answer: A


15) Vector (cross) product: If $\vec{A}$ and $\vec{B}$ are nonzero vectors for which $\vec{A} \cdot \vec{B} = 0$, it must follow that
A) $\vec{A}$ is parallel to $\vec{B}$.
TBEXAM.COM
B) $|\vec{A} \times \vec{B}| = 1$.
C) $|\vec{A} \times \vec{B}| = AB$.
D) $\vec{A} \times \vec{B} = 0$.
Answer: C

16) Addition and subtraction: You walk 55 m to the north, then turn 60° to your right and walk another 45 m. How far are you from where you originally started?
A) 94 m
B) 87 m
C) 50 m
D) 46 m
Answer: B

TBEXAM.COM

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

17) Addition and subtraction: Vectors $\vec{A}$ and $\vec{B}$ are shown in the figure. Vector $\vec{C}$ is given by $\vec{C} = \vec{B} - \vec{A}$. The magnitude of vector $\vec{A}$ is 16.0 units, and the magnitude of vector $\vec{B}$ is 7.00 units. What is the magnitude of vector $\vec{C}$?



A) 9.53
B) 15.5
C) 9.00
D) 16.2
E) 17.5


Answer: D

TBEXAM.COM

TBEXAM.COM

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

18) Addition and subtraction: Vectors $\vec{A}$ and $\vec{B}$ are shown in the figure. Vector $\vec{C}$ is given by $\vec{C} = \vec{B} - \vec{A}$. The magnitude of vector $\vec{A}$ is 16.0 units, and the magnitude of vector $\vec{B}$ is 7.00 units. What is the angle of vector $\vec{C}$, measured counterclockwise from the +*x*-axis?



A) 16.9°
B) 22.4°
C) 292°
D) 73.1°
E) 287°

TBEXAM.COM

Answer: E

19) Addition and subtraction: A rabbit trying to escape a fox runs north for 8.0 m, darts northwest for 1.0 m, then drops 1.0 m down a hole into its burrow. What is the magnitude of the net displacement of the rabbit?

A) 8.8 m
B) 10 m
C) 8.1 m
D) 66 m

Answer: A

TBEXAM.COM

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

20) Addition and subtraction: You walk 53 m to the north, then turn 60° to your right and walk another 45 m. Determine the direction of your displacement vector. Express your answer as an angle relative to east.

A) 57° N of E
B) 69° N of E
C) 63° N of E
D) 50° N of E

Answer: C

21) Components: Vector $\vec{A}$ has a magnitude 5.00 and points in a direction 40.0° clockwise from the negative y axis. What are the x and y components of vector $\vec{A}$.

A) $A_X = 4.29$ and $A_y = 2.16$
B) $A_X = 3.83$ and $A_y = 3.21$
C) $A_X = -3.21$ and $A_y = -3.83$
D) $A_X = -3.21$ and $A_y = 3.83$
E) $A_X = 3.83$ and $A_y = -3.21$

Answer: C

22) Components: The components of vector $\vec{A}$ are $A_x = +3.90$ and $A_y = -4.00$. What is the angle measured counterclockwise from the +x-axis to vector $\vec{A}$?

A) 134°
B) 46.0°
C) 314°
D) 136°
E) 224°

TBEXAM.COM

Answer: C

TBEXAM.COM

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

23) Components: Vector $\vec{A}$ has a magnitude of 5.5 cm and points along the x-axis. Vector $\vec{B}$ has a magnitude of 7.5 cm and points at +30° above the negative x-axis.

(a) Determine the x and y components of Vector $\vec{A}$.
(b) Determine the x and y components of Vector $\vec{B}$.
(c) Determine x and y components of the sum of these two vectors.
(d) Determine the magnitude and direction of the sum of these two vectors.

Answer: (a) $A_x = 5.5$ cm, $A_y = 0$
(b) $B_x = -6.5$ cm, $B_y = 3.8$ cm
(c) $R_x = -1.0$ cm, $R_y = 3.8$ cm
(d) 3.9 cm at 75° above -x-axis

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

24) Components: Vector $\vec{A}$ has a magnitude of 75.0 cm and points at 30° above the positive x-axis. Vector $\vec{C}$ has a magnitude of 25.0 cm and points along the negative x-axis. Vector $\vec{C}$ has a magnitude of 40.0 cm and points at 45° below the negative x-axis.

(a) Determine the x and y components of Vector $\vec{A}$.

(b) Determine the x and y components of Vector $\vec{B}$.

(c) Determine the x and y components of Vector $\vec{C}$.

(d) Determine x and y components of the sum of these three vectors.

(e) Determine the magnitude and direction of the sum of these three vectors.

Answer: (a) A_x = 65 cm, A_y = 38 cm

(b) B_x = -25 cm, B_y = 0

(c) C_x = -28 cm, C_y = -28 cm

(d) R_x = 12 cm, R_y = 9.2 cm

(e) 15 cm at 38° above +x-axis

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

25) Components: A helicopter is flying horizontally with a speed of 444 m/s over a hill that slopes upward with a 2% grade (that is, the "rise" is 2% of the "run"). What is the component of the helicopter's velocity perpendicular to the sloping surface of the hill?

A) 444 m/s

B) 220 m/s

C) 435 m/s

D) 8.9 m/s

Answer: D

TBEXAM.COM

26) Components: An apple falls from an apple tree growing on a 20° slope. The apple hits the ground with an impact velocity of 16.2 m/s straight downward. What is the component of the apple's impact velocity parallel to the surface of the slope?

A) 15 m/s

B) 5.5 m/s

C) 8.7 m/s

D) 12 m/s

Answer: B

TBEXAM.COM

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

27) Components: The components of vector $\vec{A}$ are A_x = +2.2 and A_y = -6.9 , and the components of vector $\vec{B}$ are given are B_x = -6.1 and B_y = -2.2. What is the magnitude of the vector $\vec{B}$ - $\vec{A}$?

A) 91
B) 6.1
C) 9.9
D) 9.5
E) 0.76

Answer: D

28) Components: The components of vector $\vec{B}$ are B_x = -3.5 and B_y = -9.7, and the components of vector $\vec{C}$ are C_x = -6 and C_y = +8.1. What is the angle (less than 180 degrees) between vectors $\vec{B}$ and $\vec{C}$?

A) 56°
B) 106°
C) 163°
D) 124°
E) 17°

Answer: D

29) Components: An airplane undergoes the following displacements: First, it flies 66 km in a direction 30° east of north. Next, it flies 49 km due south. Finally, it flies 100 km 30° north of west. Using vector components, determine how far the airplane ends up from its starting point.

TBEXAM.COM

A) 79 km
B) 76 km
C) 81 km
D) 78 km
E) 82 km


Answer: A

TBEXAM.COM

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 8th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

30) Components: In the figure, the magnitude of vector $\vec{A}$ is 18.0 units, and the magnitude of vector $\vec{B}$ is 12.0 units. What vector $\vec{C}$ must be added to the vectors $\vec{A}$ and $\vec{B}$ so that the resultant of these three vectors points in the -x direction and has a magnitude of 7.50 units? Use vector components to find your answer, and express vector $\vec{C}$ by giving its magnitude and the angle it makes with the +x-axis taking counterclockwise to be positive.



.COM


Answer: 15.5, 209°

TBEXAM.COM

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

31) Components: Three forces are exerted on an object placed on a tilted floor. Forces are vectors. The three forces are directed as shown in the figure. If the forces have magnitudes F_1 = 1.0 N, F_2 = 8.0 N and F_3 = 7.0 N, where N is the standard unit of force, what is the component of the *net force* $\vec{F}_{\text{net}} = \vec{F}_1 + \vec{F}_2 + \vec{F}_3$ parallel to the floor?



A) 6.0 N
B) 7.8 N
C) 2.5 N
D) 5.1 N


Answer: C

TBEXAM.COM

TBEXAM.COM

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

32) Components: As shown in the figure, three force vectors act on an object. The magnitudes of the forces as shown in the figure are $F_1 = 80.0$ N, $F_2 = 60.0$ N, and $F_3 = 40.0$ N, where N is the standard SI unit of force. The resultant force acting on the object is given by



A) 20.0 N at an angle 34.3° with respect to +x-axis.
B) 40.0 N at an angle 60.0° with respect to +x-axis.
C) 60.0 N at an angle 90.0° with respect to +x-axis.
D) 35.5 N at an angle 34.3° with respect to +x-axis.
E) 180 N at an angle 60.0° with respect to +x-axis.

Answer: D

TBEXAM.COM

33) Components: A teacher sends her students on a treasure hunt. She gives the following instructions:
1. Walk 300 m north
2. Walk 400 m northwest
3. Walk 700 m east-southeast and the treasure is buried there.
As all the other students walk off following the instructions, Jane physics student quickly adds the displacements and walks in a straight line to find the treasure. How far and in what direction does Jane need to walk?

A) 284 m in a direction 28.2° west of north
B) 481 m in a direction 40.9° north of east
C) 187 m in a direction 67.3° north of east
D) 399 m in a direction 52.5° north of east
E) The treasure position cannot be reached in one straight walk.

Answer: B

TBEXAM.COM

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

34) Unit vectors: Vector $\vec{A} = -3.00\,\hat{i} + 3.00\,\hat{j}$ and vector $\vec{B} = 3.00\,\hat{i} + 4.00\,j'$. What is vector $\vec{C} = \vec{A} + \vec{B}$?

A) $0.00\,\hat{i} + 3.00\,\hat{j}$

B) $0.00\,\hat{i} + 7.00\,\hat{j}$

C) $-3.00\,\hat{i}$ -3.00 $\hat{j}$

D) $-3.00\,\hat{i} + 7.00\,\hat{j}$

E) $7.00\,\hat{i} + 7.00\,\hat{j}$

Answer: B

35) Unit vectors: Vector $\vec{A} = 1.00\,\hat{i} + -2.00\,\hat{j}$ and vector $\vec{B} = 3.00\,\hat{i} + 4.00\,\hat{j}$ What are the magnitude and direction of vector $\vec{C} = \vec{A} + \vec{B}$?

A) 6.00 in a direction 63.4° counterclockwise from the positive *x* axis

B) 7.21 in a direction 33.7° counterclockwise from the positive *x* axis

C) 4.47 in a direction 26.6° counterclockwise from the positive *x* axis

D) 7.21 in a direction 56.3° counterclockwise from the positive *x* axis

E) 4.47 in a direction 6.34° counterclockwise from the positive *x* axis

Answer: C

36) Unit vectors: What is the magnitude of $\vec{A} + \vec{B} + \vec{C}$, where $\vec{A} = 1.00\,\hat{i} + 4.00\,\hat{j} - 1.00\,\hat{k}$, $\vec{B} = 3.00\,\hat{i} - 1.00\,\hat{j} - 4.00\,\hat{k}$ and $\vec{C} = -1.00\,\hat{i} + 1.00\,\hat{j}$?

A) 10.76

TBEXAM.COM

B) 7.07

C) 8.12

D) 2.00

E) 6.78

Answer: B

37) Unit vectors: If $\vec{A} = +4\,\hat{i} - 2\,\hat{j} - 3\,\hat{k}$ and $\vec{C} = -4\,\hat{i} - 2\,\hat{j} - 3\,\hat{k}$, which of the following numbers is closest to the magnitude of $\vec{A} - \vec{C}$?

A) 9

B) 8

C) 10

D) 11

E) 7

Answer: B

T B E X A M . C O M


TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

38) Unit vectors: Vector $\vec{A} = -1.00\,\hat{i} + -2.00\,\hat{j}$ and vector $\vec{B} = 3.00\,\hat{i} + 4.00\,\hat{j}$ What are the magnitude and direction of vector $\vec{C} = 3.00\,\vec{A} + 2.00\,\vec{B}$?

A) 3.61 in a direction 56.3° counterclockwise from the positive *x*-axis
B) 6.72 in a direction 34.4° counterclockwise from the positive *x*-axis
C) 3.61 in a direction -56.3° counterclockwise from the positive *x*-axis
D) 5.00 in a direction 56.3° counterclockwise from the positive *x* axis
E) 3.61 in a direction 33.7° counterclockwise from the positive *x*-axis

Answer: E

39) Unit vectors: Vectors $\vec{A}$ and $\vec{B}$ are shown in the figure. What is $|-5.00\,\vec{A} + 4.00\,\vec{B}|$



A) 1028
B) 34.0
C) $-2.00\,\hat{i} - 32.0\,\hat{j}$
D) 31.8
E) $-32.0\,\hat{i} - 2.00\,\hat{j}$

Answer: D

TBEXAM.COM

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

40) Scalar (dot) product: Determine the scalar product of $\vec{A} = 6.0\hat{i} + 4.0\hat{j} - 2.0\hat{k}$ and $\vec{B} = 5.0\hat{i} - 6.0\hat{j} - 3.0\hat{k}$.

A) 12

B) $30\hat{i} + 24\hat{j} + 6\hat{k}$

C) 60

D) $30\hat{i} - 24\hat{j} + 6\hat{k}$

E) undefined

Answer: A

41) Scalar (dot) product: Determine the angle between the directions of vector $\vec{A} = 3.00\hat{i} + 1.00\hat{j}$ and vector $\vec{B} = -3.00\hat{i} + 3.00\hat{j}$.

A) 30.0°

B) 117°

C) 26.6°

D) 88.1°

E) 45.2°

Answer: B

42) Scalar (dot) product: The scalar product of vector $\vec{A} = 3.00\hat{i} + 2.00\hat{j}$ and vector $\vec{B}$ is 10.0. Which of the following vectors could be vector $\vec{B}$?

A) $2.00\hat{i} + 4.00\hat{j}$

B) $4.00\hat{i} + 6.00\hat{j}$

C) $2.00\hat{i} + 2.00\hat{j}$

D) $12.0\hat{i}$

E) $5.00\hat{i} + 4.00\hat{j}$

Answer: C

TBEXAM.COM

43) Scalar (dot) product: The angle between vector $\vec{A} = 2.00\hat{i} + 3.00\hat{j}$ and vector $\vec{B}$ is 45.0°. The scalar product of vectors $\vec{A}$ and $\vec{B}$ is 3.00. If the *x* component of vector $\vec{B}$ is positive, what is vector $\vec{B}$.

A) $0.871\hat{i} + 0.419\hat{j}$

B) $4.76\hat{i} + 0.952\hat{j}$

C) $2.96\hat{i} + -0.973\hat{j}$

D) $3.42\hat{i} + 0.684\hat{j}$

E) $1.15\hat{i} + 0.231\hat{j}$

Answer: E

T B E X A M . C O M

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

44) Scalar (dot) product: What is the angle between the vector $\vec{A} = +3\hat{i} - 2\hat{j} - 3\hat{k}$ and the +y-axis?

A) 155°
B) 90°
C) 65°
D) 115°
E) 25°

Answer: D

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

45) Scalar (dot) product: If $\vec{A} = 3\,\hat{i} - \hat{j} + 4\hat{k}$ and $\vec{B} = x\,\hat{i} + \hat{j} - 5\hat{k}$, find x so $\vec{B}$ will be perpendicular to $\vec{A}$.

Answer: 7

46) Scalar (dot) product: Two boys searching for buried treasure are standing underneath the same tree. One boy walks 18 m east and then 18 m north. The other boy walks 16 m west and then 11 m north. Find the scalar product of their net displacements from the tree.

Answer: -90 m^2

TBEXAM.COM

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

47) Scalar (dot) product: A rectangular box is positioned with its vertices at the following points:

A = (0,0,0) C = (2,4,0) E = (0,0,3) G = (2,4,3)
B = (2,0,0) D = (0,4,0) F = (2,0,3) H = (0,4,3)

TBEXAM.COM

If the coordinates all have three significant figures, the angle between the line segments AG and AH is closest to:


A) 22.5°.
B) 21.8°.
C) 45.0°.
D) 36.9°.
E) 26.6°.

Answer: B

TBEXAM.COM

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

48) Scalar (dot) product: For the vectors shown in the figure, assume numbers are accurate to two significant figures. The scalar product $\vec{A} \times \vec{C}$ is closest to



A) zero.
B) -45.
C) -16.
D) 45.
E) 16.

Answer: C

49) Vector (cross) product: What is the vector product of $\vec{A} = 2.00\,\hat{i} + 3.00\,\hat{j} + 1.00\hat{k}$ and $\vec{B} = 1.00\,\hat{i} - 3.00\,\hat{j} - 2.00\hat{k}$?

A) $-5.00\,\hat{i} + 2.00\,\hat{j} - 6.00\hat{k}$
B) $-3.00\,\hat{i} + 5.00\,\hat{j} - 9.00\hat{k}$
C) $-4.00\,\hat{i} + 3.00\,\hat{j} - 1.00\hat{k}$
D) $2.00\,\hat{i} - 9.00\,\hat{j} - 2.00\hat{k}$
E) $-9.00\,\hat{i} - 3.00\,\hat{j} - 3.00\hat{k}$

Answer: B

TBEXAM.COM

50) Vector (cross) product: What is the magnitude of the cross product of a vector of magnitude 2.00 m pointing east and a vector of magnitude 4.00 m pointing 30.0° west of north?

A) 8.00
B) 6.93
C) -4.00
D) 4.00
E) -6.93

Answer: B

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

51) Vector (cross) product: If the magnitude of the cross product of two vectors is one-half the dot product of the same vectors, what is the angle between the two vectors?

Answer: 26.6°

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

52) Vector (cross) product: If $\vec{C} = -4\hat{i} - 2\hat{j} - 3\hat{k}$, what is $\vec{C} \times \hat{j}$?

A) $+3\hat{i} - 4\hat{k}$
B) $+3\hat{i} + 4\hat{k}$
C) $-3\hat{i} - 2\hat{j} + 4\hat{k}$
D) $-3\hat{i} + 4\hat{k}$
E) $+3\hat{i} + 2\hat{j} - 4\hat{k}$

Answer: A


53) Vector (cross) product: If $\vec{B} = -2\hat{i} - 6\hat{j} + 2\hat{k}$ and $\vec{C} = -2\hat{i} - 2\hat{j} - 3\hat{k}$, which of the following numbers is closest to the magnitude of $\vec{C} \times \vec{B}$?

A) 21
B) 13
C) 25
D) 17
E) 9

Answer: C

54) Vector (cross) product: For the vectors shown in the figure, find the magnitude and direction of $\vec{B} \times \vec{A}$, assuming that the quantities shown are accurate to two significant figures.

TBEXAM.COM




A) 31, directed into the plane
B) 31, directed on the plane
C) 31, directed out of the plane
D) 26, directed into the plane
E) 26, directed out of the plane

Answer: D

TBEXAM.COM

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

55) Vector (cross) product: For the vectors shown in the figure, find the magnitude and direction of the vector product $\vec{A} \times \vec{C}$ that the quantities shown are accurate to two significant figures.



A) 45, directed on the plane
B) 16, directed out of the plane
C) 45, directed out of the plane
D) 16, directed into the plane
E) 45, directed into the plane

Answer: C

TBEXAM.COM

TBEXAM.COM

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

Answer Key
Testname: UNTITLED2

1) B
ID: upg2 2.1.1-1
Diff: 0
Objective:

2) B
ID: upg2 2.1.1-2
Diff: 0
Objective:

3) B
ID: upg2 2.1.1-3
Diff: 0
Objective:

4) $\vec{S} = \vec{M} - \vec{N}$
ID: upg2 2.1.1-4
Diff: 0
Objective:

5) A
ID: upg2 2.1.2-1
Diff: 0
Objective:

6) A
ID: upg2 2.1.2-2
Diff: 0
Objective:

7) B
ID: upg2 2.1.2-3
Diff: 0
Objective:

8) E
ID: upg2 2.1.2-4
Diff: 0
Objective:

9) B
ID: upg2 2.1.3-1
Diff: 0
Objective:

10) A
ID: upg2 2.1.4-1
Diff: 0
Objective:

11) B
ID: upg2 2.1.4-2
Diff: 0
Objective:

12) B
ID: upg2 2.1.4-3
Diff: 0
Objective:

TBEXAM.COM

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

TBEXAM.COM

13) B
ID: upg2 2.1.5-1
Diff: 0
Objective:

14) A
ID: upg2 2.1.5-2
Diff: 0
Objective:

15) C
ID: upg2 2.1.5-3
Diff: 0
Objective:

16) B
ID: upg2 2.2.1-1
Diff: 0
Objective:

17) D
ID: upg2 2.2.1-2
Diff: 0
Objective:

18) E
ID: upg2 2.2.1-3
Diff: 0
Objective:

TBEXAM.COM

19) A
ID: upg2 2.2.1-4
Diff: 0
Objective:

20) C
ID: upg2 2.2.1-5
Diff: 0
Objective:

21) C
ID: upg2 2.2.2-1
Diff: 0
Objective:

22) C
ID: upg2 2.2.2-2
Diff: 0
Objective:

23) (a) A_x = 5.5 cm, A_y = 0
(b) B_x = -6.5 cm, B_y = 3.8 cm
(c) R_x = -1.0 cm, R_y = 3.8 cm
(d) 3.9 cm at 75° above -x-axis
ID: upg2 2.2.2-3
Diff: 0
Objective:

TBEXAM.COM

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

TBEXAM.COM

24) (a) A_x = 65 cm, A_y = 38 cm
(b) B_x = -25 cm, B_y = 0
(c) C_x = -28 cm, C_y = -28 cm
(d) R_x = 12 cm, R_y = 9.2 cm
(e) 15 cm at 38° above +x-axis
ID: upg2 2.2.2-4
Diff: 0
Objective:

25) D
ID: upg2 2.2.2-5
Diff: 0
Objective:

26) B
ID: upg2 2.2.2-6
Diff: 0
Objective:

27) D
ID: upg2 2.2.2-7
Diff: 0
Objective:

28) D
ID: upg2 2.2.2-8
Diff: 0
Objective:

TBEXAM.COM

29) A
ID: upg2 2.2.2-9
Diff: 0
Objective:

30) 15.5, 209°
ID: upg2 2.2.2-10
Diff: 0
Objective:

31) C
ID: upg2 2.2.2-11
Diff: 0
Objective:

32) D
ID: upg2 2.2.2-12
Diff: 0
Objective:

33) B
ID: upg2 2.2.2-13
Diff: 0
Objective:

34) B
ID: upg2 2.2.3-1
Diff: 0
Objective:

TBEXAM.COM

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank

Answer Key
Testname: UNTITLED2

35) C
ID: upg2 2.2.3-2
Diff: 0
Objective:

36) B
ID: upg2 2.2.3-3
Diff: 0
Objective:

37) B
ID: upg2 2.2.3-4
Diff: 0
Objective:

38) E
ID: upg2 2.2.3-5
Diff: 0
Objective:

39) D
ID: upg2 2.2.3-6
Diff: 0
Objective:

40) A
ID: upg2 2.2.4-1
Diff: 0
Objective:

41) B
ID: upg2 2.2.4-2
Diff: 0
Objective:

42) C
ID: upg2 2.2.4-3
Diff: 0
Objective:

43) E
ID: upg2 2.2.4-4
Diff: 0
Objective:

44) D
ID: upg2 2.2.4-5
Diff: 0
Objective:

45) 7
ID: upg2 2.2.4-6
Diff: 0
Objective:

46) -90 m^2
ID: upg2 2.2.4-7
Diff: 0
Objective:

TBEXAM.COM

T B E X A M . C O M

TBEXAM.COM Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank
THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES

TBEXAM.COM

47) B
ID: upg2 2.2.4-8
Diff: 0
Objective:

48) C
ID: upg2 2.2.4-9
Diff: 0
Objective:

49) B
ID: upg2 2.2.5-1
Diff: 0
Objective:

50) B
ID: upg2 2.2.5-2
Diff: 0
Objective:

51) 26.6°
ID: upg2 2.2.5-3
Diff: 0
Objective:

52) A
ID: upg2 2.2.5-4
Diff: 0
Objective:

53) C
ID: upg2 2.2.5-5
Diff: 0
Objective:

54) D
ID: upg2 2.2.5-6
Diff: 0
Objective:

55) C
ID: upg2 2.2.5-7
Diff: 0
Objective:

TBEXAM.COM

T B E X A M . C O M

TBEXAM.COM THE WORLD'S LARGEST TEST BANK AND SOLUTIONS DATABASE WITH UNBEATABLE RATES
Physics for Scientists and Engineers with Modern Physics 5th edition Giancoli Test Bank