Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

SKU: 9780134437736-SOLUTIONS

INSTRUCTOR'S SOLUTIONS MANUAL

MATTHEW G. HUDELSON

BASIC TECHNICAL MATHEMATICS

AND

BASIC TECHNICAL MATHEMATICS WITH CALCULUS

ELEVENTH EDITION

Allyn J. Washington

Dutchess Community College

Richard S. Evans

Corning Community College

Pearson

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual



This work is protected by United States copyright laws and is provided solely for the use of instructors in teaching their courses and assessing student learning. Dissemination or sale of any part of this work (including on the World Wide Web) will destroy the integrity of the work and is not permitted. The work and materials from it should never be made available to students except by instructors using the accompanying text in their classes. All recipients of this work are expected to abide by these restrictions and to honor the intended pedagogical purposes and the needs of other instructors who rely on these materials.

The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expressed or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson from electronic files supplied by the author.

Copyright © 2018, 2014, 2009 Pearson Education, Inc.
Publishing as Pearson, 330 Hudson Street, NY NY 10013

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.



ISBN-13: 978-0-13-443589-3
ISBN-10: 0-13-443589-3

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

Instructor's Solutions Manual for
Basic Technical Mathematics and
Basic Technical Mathematics with Calculus, 11th Edition

Chapter 1 Basic Algebraic Operations 1

Chapter 2 Geometry 104

Chapter 3 Functions and Graphs 171

Chapter 4 The Trigonometric Functions 259

Chapter 5 Systems of Linear Equations; Determinants 346

Chapter 6 Factoring and Fractions 490

Chapter 7 Quadratic Equations 581

Chapter 8 Trigonometric Functions of Any Angle 666

Chapter 9 Vectors and Oblique Triangles 723

Chapter 10 Graphs of the Trigonometric Functions 828

Chapter 11 Exponents and Radicals 919

Chapter 12 Complex Numbers 1001

Chapter 13 Exponential and Logarithmic Functions 1090

Chapter 14 Additional Types of Equations and Systems of Equations 1183

Chapter 15 Equations of Higher Degree 1290

Chapter 16 Matrices; Systems of Linear Equations 1356

Chapter 17 Inequalities 1477

Chapter 18 Variation 1598

Chapter 19 Sequences and the Binomial Theorem 1634

Chapter 20 Additional Topics in Trigonometry 1696

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

Chapter 21 Plane Analytic Geometry 1812

Chapter 22 Introduction to Statistics 2052

Chapter 23 The Derivative 2126

Chapter 24 Applications of the Derivative 2310

Chapter 25 Integration 2487

Chapter 26 Applications of Integration 2572

Chapter 27 Differentiation of Transcendental Functions 2701

Chapter 28 Methods of Integration 2839

Chapter 29 Partial Derivatives and Double Integrals 2991

Chapter 30 Expansion of Functions in Series 3058

Chapter 31 Differential Equations 3181

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

Chapter 1


Basic Algebraic Operations

1.1 Numbers

1. The numbers –7 and 12 are integers. They are also rational numbers since they can be written as $\frac{-7}{1}$ and $\frac{12}{1}$.

2. The absolute value of –6 is 6, and the absolute value of –7 is 7. We write these as $|-6| = 6$ and $|-7| = 7$.

3. $-6 < -4$; –6 is to the left of –4.



4. The reciprocal of $\frac{3}{2}$ is $\frac{1}{3/2} = 1 \times \frac{2}{3} = \frac{2}{3}$.

5. 3 is an integer, rational $\left(\frac{3}{1}\right)$, and real.

$\sqrt{-4}$ is imaginary.

6. $\frac{\sqrt{7}}{3}$ is irrational (because $\sqrt{7}$ is an irrational number) and real.

-6 is an integer, rational $\left(\frac{-6}{1}\right)$, and real.

7. $-\frac{\pi}{6}$ is irrational (because π is an irrational number) and real.

$\frac{1}{8}$ is rational and real.

8. $-\sqrt{-6}$ is imaginary.

$-2.33 = \frac{-233}{100}$ is rational and real.

9. $|3| = 3$

$|-3| = 3$

$\left|-\frac{\pi}{2}\right| = \frac{\pi}{2}$

10. $|-0.857| = 0.857$


$|\sqrt{2}| = \sqrt{2}$

$\left|-\frac{19}{4}\right| = \frac{19}{4}$


Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual


11. $6 < 8$; 6 is to the left of 8.




12. $7 > 5$; 7 is to the right of 5.




13. $\pi < 3.1416$; π (3.1415926…) is to the left of 3.1416.




14. $-4 < 0$; -4 is to the left of 0.



15. $-4 < -|-3|$; -4 is to the left of $-|-3|$, $\left(-|-3| = -(3) = -3\right)$.




16. $-\sqrt{2} > -1.42$; $(-\sqrt{2} = -(1.414\ldots) = -1.414\ldots), -\sqrt{2}$ is to the right of -1.42.



17. $-\frac{2}{3} > -\frac{3}{4}; -\frac{2}{3} = -0.666\ldots$ is to the right of $-\frac{3}{4} = -0.75$.



18. $-0.6 < 0.2$; -0.6 is to the left of 0.2.



Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

19. The reciprocal of 3 is $\frac{1}{3}$.

The reciprocal of $-\frac{4}{\sqrt{3}}$ is $-\frac{1}{4/\sqrt{3}} = -\frac{\sqrt{3}}{4}$.


The reciprocal of $\frac{y}{b}$ is $\frac{1}{y/b} = \frac{b}{y}$.

20. The reciprocal of $-\frac{1}{3}$ is $-\frac{1}{1/3} = -\frac{3}{1} = -3$.


The reciprocal of $0.25 = \frac{1}{4}$ is $\frac{1}{1/4} = \frac{4}{1} = 4$.

The reciprocal of $2x$ is $\frac{1}{2x}$.

21. Find 2.5, $-\frac{12}{5} = -2.4; -\frac{3}{4} = -0.75; \sqrt{3} = 1.732...$



22. Find $-\frac{7}{3} = -2.333...; -\frac{\sqrt{2}}{2} = -\frac{1.414\ldots}{2} = -0.707;\ 2\pi = 2 \times 3.14\ldots = 6.28;\ \frac{123}{19} = 6.47$.



23. An absolute value is not always positive, $|0| = 0$ which is not positive.

24. Since $-2.17 = -\frac{217}{100}$, it is rational.

25. The reciprocal of the reciprocal of any positive or negative number is the number itself.

The reciprocal of n is $\frac{1}{n}$; the reciprocal of $\frac{1}{n}$ is $\frac{1}{1/n} = 1 \cdot \frac{n}{1} = n$.

26. Any repeating decimal is rational, so $2.\overline{72}$ is rational. It turns out that $2.\overline{72} = \frac{30}{11}$.

27. It is true that any nonterminating, nonrepeating decimal is an irrational number.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

28. No, $|b-a|=|b|-|a|$, as shown below.

If $a>0$, then $|a|=a$.


If $b>a$ and $a>0$, then $|b|=b$.

If $b>a$ then $b-a>0$, then $|b-a|=b-a$.

Therefore, $|b-a|=b-a=|b|-|a|$.


The two sides of the expression are equivalent, one side is not less than the other.

29. List these numbers from smallest to largest: $-1,\ 9,\ \pi=3.14,\ \sqrt{5}=2.236,\ |-8|=8,\ -|-3|=-3,\ -3.1$.



So, from smallest to largest, they are $-3.1,\ -|-3|,\ -1,\ \sqrt{5},\ \pi,\ |-8|,\ 9$.

30. List these numbers from smallest to largest: $\frac{1}{5}=0.20,\ -\sqrt{10}=-3.16...,\ -|-6|=-6,\ -4,\ 0.25,\ |-\pi|=3.14...$.



So, from smallest to largest, they are $-|-6|,\ -4,\ -\sqrt{10},\ \frac{1}{5},\ 0.25,\ |-\pi|$.

31. If a and b are positive integers and $b>a$, then

(a) $b-a$ is a positive integer.

(b) $a-b$ is a negative integer.

(c) $\frac{b-a}{b+a}$, the numerator and denominator are both positive, but the numerator is less than the denominator, so the answer is a positive rational number than is less than 1.

32. If a and b are positive integers, then

(a) $a+b$ is a positive integer

(b) a/b is a positive rational number

(c) $a\times b$ is a positive integer

33. **(a)** Is the absolute value of a positive or a negative integer always an integer?

$|x|=x$, so the absolute value of a positive integer is an integer.

$|-x|=x$, so the absolute value of a negative integer is an integer.

(b) Is the reciprocal of a positive or negative integer always a rational number?

If x is a positive or negative integer, then the reciprocal of x is $\frac{1}{x}$. Since both 1 and x are integers, the reciprocal is a rational number.

Copyright © 2018 Pearson Education, Inc.


Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

34. **(a)** Is the absolute value of a positive or negative rational number rational?
$|x| = x$, so if x is a positive or negative rational number, the absolute value of it is also a rational number.


(b) Is the reciprocal of a positive or negative rational number a rational number?

A rational number is a number that can be expressed as a fraction where both the numerator and denominator are integers and the denominator is not zero. So a rational number $\frac{\text{integer } a}{\text{integer } b}$ has a reciprocal of $\frac{1}{\frac{\text{integer } a}{\text{integer } b}} = \frac{\text{integer } b}{\text{integer } a}$, which is also a rational number if integer a is not zero.


35. **(a)** If $x > 0$, then x is a positive number located to the right of zero on the number line.




(b) If $x < -4$, then x is a negative number located to the left of –4 on the number line.




36. **(a)** If $|x| < 1$, then $-1 < x < 1$.




(b) $|x| > 2$, then $x < -2$ or $x > 2$.



37. If $x > 1$, then $\frac{1}{x}$ is a positive number less than 1. Or $0 < \frac{1}{x} < 1$.



38. If $x < 0$, then $|x|$ is a positive number greater than zero.



39. $a + bj = a + b\sqrt{-1}$ is a real number when $\sqrt{-1}$ is eliminated, which is when $b = 0$. So $a + bj$ is a real number for all real values of a and $b = 0$.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

40. The variables are w and t.
The constants are c, 0.1, and 1.

41. $\frac{1}{C_T}=\frac{1}{C_1}+\frac{1}{C_2}$. Find C_T, where $C_1=0.0040\,\text{F}$ and $C_2=0.0010\,\text{F}$.

$$\frac{1}{C_T}=\frac{1}{0.0040}+\frac{1}{0.0010}$$

$$\frac{1}{C_T}=\frac{1(0.0040)+1(0.0010)}{0.0040\times 0.0010}$$

$$C_T=\frac{0.0040\times 0.0010}{0.0040+0.0010}=\frac{0.0000040}{0.0050}$$

$$C_T=0.00080\text{ F}$$

42. $|100V|=100V$


$|-200V|=200V$

$|-200V|>|100V|$

43. $N=\frac{a\text{ bits}}{\text{bytes}}\times\frac{1000\text{ bytes}}{1\text{ kilobyte}}\times n\,\text{kilobytes}$

$N=1000\,an\text{ bits}$

44.



$x=$ length of base in m


$y=$ the shortened length in centimetres.

$100x=$ length of base in cm

$y+L=100x,$ all dimensions in cm

$L=100x-y$

45. Yes, $-20\,^\circ\text{C}>-30\,^\circ\text{C}$ because $-30\,^\circ\text{C}$ is found to the left of $-20\,^\circ\text{C}$ on the number line.



46. For $I<4$ A, $R>12\ \Omega$.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

1.2 Fundamental Operations of Algebra

1. $16-2\times(-2)=16-(-4)=16+4=20$

2. $\dfrac{-18}{-6}+5-(-2)(3)=3+5-(-6)=8+6=14$

3. $\dfrac{-12}{8-2}+\dfrac{5-1}{2(-1)}=\dfrac{-12}{6}+\dfrac{4}{-2}=-2+(-2)=-4$

4. $\dfrac{7\times6}{0\times0}=\dfrac{42}{0}=$ is undefined , not indeterminate.

5. $5+(-8)=5-8=-3$

6. $-4+(-7)=-4-7=-11$

7. $-3+9=6$ or alternatively
$-3+9=+(9-3)=+(6)=6$

8. $18-21=-3$ or alternatively
$18-21=-(21-18)=-(3)=-3$

9. $-19-(-16)=-19+16=-3$

10. $-8-(-10)=-8+10=2$

11. $7(-4)=-(7\times4)=-28$

12. $-9(3)=-27$

13. $-7(-5)=+(7\times5)=35$

14. $\dfrac{-9}{3}=-3$

15. $\dfrac{-6(20-10)}{-3}=\dfrac{-6(10)}{-3}=\dfrac{-60}{-3}=20$

16. $\dfrac{-28}{-7(5-6)}=\dfrac{-28}{-7(-1)}=\dfrac{-28}{7}=-4$

17. $-2(4)(-5)=-8(-5)=40$

18. $-3(-4)(-6)=12(-6)=-72$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

19. $2(2-7)\div 10 = 2(-5)\div 10 = -10\div 10 = -1$

20. $\dfrac{-64}{-2|4-8|} = \dfrac{-64}{-2|-4|} = \dfrac{-64}{-2(4)} = \dfrac{-64}{-8} = 8$

21. $16\div 2(-4) = 8(-4) = -32$

22. $-20\div 5(-4) = -4(-4) = 16$

23. $-9-|2-10| = -9-|-8| = -9-8 = -17$

24. $(7-7)\div(5-7) = 0\div(-2) = 0$

25. $\dfrac{17-7}{7-7} = \dfrac{10}{0}$ is undefined

26. $\dfrac{(7-7)(2)}{(7-7)(-1)} = \dfrac{0(2)}{0(-1)} = \dfrac{0}{0}$ is indeterminate

27. $8-3(-4) = 8+12 = 20$

28. $-20+8\div 4 = -20+2 = -18$

29. $-2(-6)+\left|\dfrac{8}{-2}\right| = 12+|-4| = 12+4 = 16$

30. $\dfrac{|-2|}{-2} = \dfrac{2}{-2} = -1$

31. $10(-8)(-3)\div(10-50) = 10(-8)(-3)\div(-40)$
$= -80(-3)\div(-40)$
$= 240\div(-40)$
$= -6$

32. $\dfrac{7-|-5|}{-1(-2)} = \dfrac{7-5}{2} = \dfrac{2}{2} = 1$

33. $\dfrac{24}{3+(-5)} - 4(-9) = \dfrac{24}{-2} + (4\times 9) = -12+36 = 24$

34. $\dfrac{-18}{3} - \dfrac{4-|-6|}{-1} = \dfrac{-18}{3} - \dfrac{4-6}{-1} = -6 - \dfrac{-2}{-1} = -6-2 = -8$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

35. $$-7-\frac{|-14|}{2(2-3)}-3|6-8| = -7-\frac{14}{2(-1)}-3|-2|$$
$$= -7-\frac{14}{-2}-3(2)$$
$$= -7-(-7)-6$$
$$= -7+7-6$$
$$= -6$$

36. $$-7(-3)+\frac{-6}{-3}-|-9| = +(7\times 3)+2-9$$
$$= 21+2-9$$
$$= 14$$

37. $$\frac{3|-9-2(-3)|}{1-10} = \frac{3|-9+6|}{-9}$$
$$= \frac{3|-3|}{-9}$$
$$= \frac{9}{-9}$$
$$= -1$$

38. $$\frac{20(-12)-40(-15)}{98-|-98|} = \frac{-240+600}{98-98} = \frac{360}{0} = \text{is undefined}$$

39. $6(7)=(7)6$ demonstrates the commutative law of multiplication.

40. $6+8=8+6$ demonstrates the commutative law of addition.

41. $6(3+1)=6(3)+6(1)$ demonstrates the distributive law.

42. $4(5\times\pi)=(4\times 5)\pi$ demonstrates the associative law of multiplication.

43. $3+(5+9)=(3+5)+9$ demonstrates the associative law of addition.

44. $8(3-2)=8(3)-8(2)$ demonstrates the distributive law.

45. $(\sqrt{5}\times 3)\times 9=\sqrt{5}\times(3\times 9)$ demonstrates the associative law of multiplication.

46. $(3\times 6)\times 7=7\times(3\times 6)$ demonstrates the commutative law of multiplication.

47. $-a+(-b)=-a-b$, which is expression (d).

48. $b-(-a)=b+a=a+b$, which is expression (a).

49. $-b-(-a)=-b+a=a-b$, which is expression (b).

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

50. $-a-(-b)=-a+b=b-a$, which is expression (c).

51. Since $|5-(-2)|=|5+2|=|7|=7$ and $|-5-(-2)|=|-5+2|=|-3|=3$,
$|5-(-2)|>|-5-(-2)|$.

52. Since $|-3-|-7||=|-3-7|=|-10|=10$ and $||-3|-7|=|3-7|=|-4|=4$,
$|-3-|-7||>||-3|-7|$.

53. **(a)** The sign of a product of an even number of negative numbers is positive. Example: $-3(-6)=18$

(b) The sign of a product of an odd number of negative numbers is negative.
Example: $-5(-4)(-2)=-40$

54. Subtraction is not commutative because $x-y\neq y-x$. Example: $7-5=2$ does not equal $5-7=-2$

55. Yes, from the definition in Section 1.1, the absolute value of a positive number is the number itself, and the absolute value of a negative number is the corresponding positive number. So for values of x where $x>0$ (positive) or $x=0$ (neutral) then $|x|=x$.

Example: $|4|=4$.

The claim that absolute values of negative numbers $|x|=-x$ is also true.

Example: if x is -6, then $|-6|=-(-6)=6$.

56. The incorrect answer was achieved by subtracting before multiplying or dividing which violates the order of operations.
$24-6\div 2\times 3\neq 18\div 2\times 3=9\times 3=27$
The correct value is:
$24-6\div 2\times 3=24-3\times 3=24-9=15$

57. **(a)** $-xy=1$ is true for values of x and y that are negative reciprocals of each other or $y=-\dfrac{1}{x}$, providing that the number x in the denominator is not zero. So if $x=12$, then $y=-\dfrac{1}{12}$ and $-xy=-(12)\left(-\dfrac{1}{12}\right)=1$.

(b) $\dfrac{x-y}{x-y}=1$ is true for all values of x and y, providing that $x\neq y$ to prevent division by zero.

58. **(a)** $|x+y|=|x|+|y|$ is true for values where both x and y have the same sign or either are zero:
$|x+y|=|x|+|y|$, when $x\geq 0$ and $y\geq 0$ or when $x\leq 0$ and $y\leq 0$
Example:
$|6+3|=6+3=9$ and
$|6|+|3|=6+3=9$
Also,
$|-6+(-3)|=|-9|=9$
$|-6|+|-3|=6+3=9$

$|x+y|=|x|+|y|$ is not true however, when x and y have opposite signs
$|x+y|\neq|x|+|y|$, when $x>0$ and $y<0$; or $x<0$ and $y>0$.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

Example:
$|-21+6|=|-15|=15,$
$|-21|+|6|=21+6=27\neq 15$

$|4+(-5)|=|-1|=1,$
$|4|+|-5|=4+5=9\neq 1$

(b) In order for $|x-y|=|x|+|y|$ it is necessary that they have opposite signs or either to be zero. Symbolically, $|x-y|=|x|+|y|$ when $x\geq 0$ and $y\leq 0$; or when $x\leq 0$ and $y\geq 0$.

Example:
$|6-(-3)|=6+3=9$ and
$|6|+|-3|=6+3=9$

Example:
$|-11-7|=|-18|=18$
$|-11|+|-7|=11+7=18$

$|x-y|=|x|+|y|$ is not true, however, when x and y have the same signs.
$|x-y|\neq|x|+|y|$, when $x>0$ and $y>0$; or $x<0$ and $y<0$.
Example:
$|21-6|=|15|=15,$
$|21|+|6|=27\neq 15$

59. The total change in the price of the stock is $-0.68+0.42+0.06+(-0.11)+0.02=-0.29$.

60. The difference in altitude is $-86-(-1396)=1396-86=1310$ m

61. The change in the meter energy reading E would be:
$E_{change}=E_{used}-E_{generated}$
$E_{change}=2.1\text{ kW}\cdot\text{h}-1.5\text{ kW}(3.0\text{ h})$
$E_{change}=2.1\text{ kW}\cdot\text{h}-4.5\text{ kW}\cdot\text{h}$
$E_{change}=-2.4\text{ kW}\cdot\text{h}$

62. Assuming that this batting average is for the current season only which is just starting, the number of hits is zero and the total number of at-bats is also zero giving us a batting average $=\dfrac{\text{number of hits}}{\text{at}-\text{bats}}=\dfrac{0}{0}$ which is indeterminate, not 0.000.

63. The average temperature for the week is:
$T_{avg}=\dfrac{-7+(-3)+2+3+1+(-4)+(-6)}{7}\ ^\circ\text{C}$
$T_{avg}=\dfrac{-7-3+2+3+1-4-6}{7}\ ^\circ\text{C}$
$T_{avg}=\dfrac{-14}{7}\ ^\circ\text{C}=-2.0\ ^\circ\text{C}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

64. The vertical distance from the flare gun is

$d = (70)(5) + (-16)(25)$

$d = 350 + (-400)$

$d = 350 - 400$

$d = -50$ m

The flare is 50 m below the flare gun.

65. The sum of the voltages is

$V_{sum} = 6\text{V} + (-2\text{V}) + 8\text{V} + (-5\text{V}) + 3\text{V}$

$V_{sum} = 6\text{V} - 2\text{V} + 8\text{V} - 5\text{V} + 3\text{V}$

$V_{sum} = 10\text{V}$

66. **(a)** The change in the current for the first interval is the second reading – the first reading

$Change_1 = -2 \text{ lb/in}^2 - 7 \text{ lb/in}^2 = -9 \text{ lb/in}^2$.

(b) The change in the current for the middle intervals is the third reading – the second reading

$Change_2 = -9 \text{ lb/in}^2 - (-2 \text{ lb/in}^2) = -9 \text{ lb/in}^2 + 2 \text{ lb/in}^2 = -7 \text{ lb/in}^2$.

(c) The change in the current for the last interval is the last reading – the third reading

$Change_3 = -6 \text{ lb/in}^2 - (-9 \text{ lb/in}^2) = -6 \text{ lb/in}^2 + 9 \text{ lb/in}^2 = 3 \text{ lb/in}^2$.

67. The oil drilled by the first well is $100 \text{ m} + 200 \text{ m} = 300 \text{ m}$ which equals the depth drilled by the second well $200 \text{ m} + 100 \text{ m} = 300 \text{ m}$.

$100 \text{ m} + 200 \text{ m} = 200 \text{ m} + 100 \text{ m}$ demonstrates the commutative law of addition.

68. The first tank leaks $12\frac{\text{L}}{\text{h}}(7 \text{ h}) = 84 \text{ L}$. The second tank leaks $7\frac{\text{L}}{\text{h}}(12\text{h}) = 84\text{L}$.

$12 \times 7 = 7 \times 12$ demonstrates the commutative law of multiplication.

69. The total time spent browsing these websites is the total time spent browsing the first site on each day + the total time spent browsing the second site on each day

$t = 7 \text{ days} \times 25 \frac{\text{minutes}}{\text{day}} + 7 \text{ days} \times 15 \frac{\text{minutes}}{\text{day}}$

$t = 175 \text{ min} + 105 \text{ min}$

$t = 280 \text{ min}$

OR

$t = 7 \text{ days} \times (25 + 15) \frac{\text{minutes}}{\text{day}}$

$t = 7 \text{ days} \times 40 \frac{\text{minutes}}{\text{day}}$

$t = 280 \text{ min}$

which illustrates the distributive law.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

70. Distance = rate × time

$$d = \left(600\frac{\text{km}}{\text{h}} + 50\frac{\text{km}}{\text{h}}\right)\ 3\ \text{h}$$

$$d = 600\frac{\text{km}}{\text{h}}(3\text{h}) + 50\frac{\text{km}}{\text{h}}(3\text{h})$$

$$d = 1800\ \text{km} + 150\ \text{km} = 1950\ \text{km}$$

OR

$$d = \left(600\frac{\text{km}}{\text{h}} + 50\frac{\text{km}}{\text{h}}\right)\ 3\ \text{h}$$

$$d = \left(650\frac{\text{km}}{\text{h}}\right)\ 3\ \text{h}$$

$$d = 1950\ \text{km}$$

This illustrates the distributive law.

1.3 Calculators and Approximate Numbers

1. 0.390 has three significant digits since the zero is after the decimal. The zero is not necessary as a placeholder and should not be written unless it is significant.

2. 35.303 rounded off to four significant digits is 35.30.

3. In finding the product of the approximate numbers, $2.5 \times 30.5 = 76.25$, but since 2.5 has 2 significant digits, the answer is 76.

4. $38.3 - 21.9(-3.58) = 116.702$ using exact numbers; if we estimate the result, $40 - 20(-4) = 120$.

5. 8 cylinders is exact because they can be counted. 55 km/h is approximate since it is measured.

6. 0.002 mm thick is a measurement and is therefore an approximation. $7.50 is an exact price.

7. 24 hr and 1440 min (60 min/h × 24 h =1140 min) are both exact numbers.

8. 50 keys is exact because you can count them; 50 h of use is approximate since it is a measurement of time.

9. Both 1 cm and 9 g are measured quantities and so they are approximate.

10. The numbers 90 and 75 are exact counts of windows while 15 years is a measurement of time, hence it is approximate.

11. 107 has 3 significant digits; 3004 has 4 significant digits; 1040 has 3 significant digits (the final zero is a placeholder.)

12. 3600 has 2 significant digits; 730 has 2 significant digits; 2055 has 4 significant digits.

13. 6.80 has 3 significant digits since the zero indicates precision; 6.08 has 3 significant digits; 0.068 has 2 significant digits (the zeros are placeholders.)

14. 0.8730 has 4 significant digits; 0.0075 has 2 significant digits; 0.0305 has 3 significant digits.

15. 3000 has 1 significant digit; 3000.1 has 5 significant digits; 3000.10 has 6 significant digits.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

16. 1.00 has 3 significant digits since the zeros indicate precision; 0.01 has 1 significant digit since leading zeros are not significant; 0.0100 has 3 significant digits, counting the trailing zeros.

17. 5000 has 1 significant digit; 5000.0 has 5 significant digits; $500\overline{0}$ has 4 significant digits since the bar over the final zero indicates that it is significant.

18. 200 has 1 significant digit; $20\overline{0}$ has 3 significant digits; 200.00 has 5 significant digits.

19. **(a)** 0.010 has more decimal places (3) and is more precise.
(b) 30.8 has more significant digits (3) and is more accurate.

20. **(a)** Both 0.041 and 7.673 have the same precision as they have the same number of decimal places (3).
(b) 7.673 is more accurate because it has more significant digits (4) than 0.041, which has 2 significant digits.

21. **(a)** Both 0.1 and 78.0 have the same precision as they have the same number of decimal places.
(b) 78.0 is more accurate because it has more significant digits (3) than 0.1, which has 1 significant digit.

22. **(a)** 0.004 is more precise because it has more decimal places (3).
(b) 7040 is more accurate because it has more significant digits (3) than 0.004, which has only 1 significant digit.

23. **(a)** 0.004 is more precise because it has more decimal places (3).
(b) Both have the same accuracy as they both have 1 significant digit.

24. The precision and accuracy of $|-8.914|$ and 8.914 are the same.
(a) Both 50.060 and 8.914 have the same precision as they have the same number of decimal places (3).
(b) 50.060 is more accurate because it has more significant digits (5) than 8.914, which has 4 significant digits.

25. **(a)** 4.936 rounded to 3 significant digits is 4.94.
(b) 4.936 rounded to 2 significant digits is 4.9.

26. **(a)** 80.53 rounded to 3 significant digits is 80.5.
(b) 80.53 rounded to 2 significant digits is 81.

27. **(a)** -50.893 rounded to 3 significant digits is -50.9.
(b) -50.893 rounded to 2 significant digits is -51.

28. **(a)** 7.004 rounded to 3 significant digits is 7.00.
(b) 7.004 rounded to 2 significant digits is 7.0.

29. **(a)** 5968 rounded to 3 significant digits is 5970.
(b) 5968 rounded to 2 significant digits is $60\overline{0}0$.

30. **(a)** 30.96 rounded to 3 significant digits is 31.0.
(b) 30.96 rounded to 2 significant digits is 31.

31. **(a)** 0.9449 rounded to 3 significant digits is 0.945.
(b) 0.9449 rounded to 2 significant digits is 0.94.

32. **(a)** 0.9999 rounded to 3 significant digits is 1.00.
(b) 0.9999 rounded to 2 significant digits is 1.0.

33. **(a)** Estimate: $13+1-2=12$
(b) Calculator: $12.78+1.0495-1.633=12.1965$, which is 12.20 to 0.01 precision

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

34. **(a)** Estimate: $4\times17=68$
(b) Calculator: $3.64(17.06)=62.0984$, which is 62.1 to 3 significant digits

35. **(a)** Estimate $0.7\times4-9=-6$
(b) Calculator: $0.6572\times3.94-8.651=-6.061632$, which is -6.06 to 3 significant digits

36. **(a)** Estimate $40-26\div4=40-6.5=34$
(b) Calculator: $41.5-26.4\div3.7=34.3648649$, which is 34 to 2 significant digits

37. **(a)** Estimate $9+(1)(4)=9+4=13$
(b) Calculator: $8.75+(1.2)(3.84)=13.358$, which is 13 to 2 significant digits

38. **(a)** Estimate $30-\dfrac{20}{2}=30-10=20$

(b) Calculator: $28-\dfrac{20.955}{2.2}=18.475$, which is 18 to 2 significant digits

39. **(a)** Estimate $\dfrac{9(15)}{9+15}=\dfrac{135}{24}=6$, to 1 significant digit

(b) Calculator: $\dfrac{8.75(15.32)}{8.75+15.32}=5.569173$, which is 5.57 to 3 significant digits

40. **(a)** Estimate $\dfrac{9(4)}{2+5}=\dfrac{36}{7}=5$, to 1 significant digit

(b) Calculator: $\dfrac{8.97(4.003)}{2.0+4.78}=5.296$, which is 5.3 to 2 significant digits

41. **(a)** Estimate $4.5-\dfrac{2(300)}{400}=3.0$, to 2 significant digits

(b) Calculator: $4.52-\dfrac{2.056(309.6)}{395.2}=2.9093279$, which is 2.91 to 3 significant digits

42. **(a)** Estimate $8+\dfrac{15}{2+2}=12$, to 2 significant digits

(b) Calculator: $8.195+\dfrac{14.9}{1.7+2.1}=12.1160526$, which is 12 to 2 significant digits

43. $0.9788+14.9=15.8788$ since the least precise number in the question has 4 decimal places.

44. $17.311-22.98=-5.669$ since the least precise number in the question has 3 decimal places.

45. $-3.142(65)=-204.23$, which is -204.2 because the least accurate number has 4 significant digits.

46. $8.62\div1728=0.004988$, which is 0.00499 because the least accurate number has 3 significant digits.

47. With a frequency listed as 2.75 MHz, the least possible frequency is 2.745 MHz, and the greatest possible frequency is 2.755 MHz. Any measurements between those limits would round to 2.75 MHz.

48. For an engine displacement stated at 2400 cm^3, the least possible displacement is 2350 cm^3, and the greatest possible displacement is 2450 cm^3. Any measurements between those limits would round to 2400 cm^3.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

49. The speed of sound is $3.25 \text{ mi} \div 15 \text{ s} = 0.21666... \text{ mi/s} = 1144.0... \text{ ft/s}$. However, the least accurate measurement was time since it has only 2 significant digits. The correct answer is 1100 ft/s.

50. $4.4 \text{ s} - 2.72 \text{ s} = 1.68 \text{ s}$, but the answer must be given according to precision of the least precise measurement in the question, so the correct answer is 1.7 s.

51. **(a)** $2.2 + 3.8 \times 4.5 = 2.2 + (3.8 \times 4.5) = 19.3$

(b) $(2.2 + 3.8) \times 4.5 = 6.0 \times 4.5 = 27$

52. **(a)** $6.03 \div 2.25 + 1.77 = (6.03 \div 2.25) + 1.77 = 4.45$

(b) $6.03 \div (2.25 + 1.77) = 6.03 \div 4.02 = 1.5$

53. **(a)** $2 + 0 = 2$

(b) $2 - 0 = 2$

(c) $0 - 2 = -2$

(d) $2 \times 0 = 0$

(e) $2 \div 0 =$ error; from Section 1.2, an equation that has 0 in the denominator is undefined when the numerator is not also 0.

54. **(a)** $2 \div 0.0001 = 20\ 000$; $2 \div 0 =$ error

(b) $0.0001 \div 0.0001 = 1$; $0 \div 0 =$ error

(c) Any number divided by zero is undefined. Zero divided by zero is indeterminate.

55. $\pi = 3.14159265...$

(a) $\pi < 3.1416$

(b) $22 \div 7 = 3.1428$

$\pi < (22 \div 7)$

56. **(a)** $8 \div 33 = 0.2424... = 0.\overline{24}$

(b) $\pi = 3.14159265...$

57. **(a)** $1 \div 3 = 0.333...$ It is a rational number since it is a repeating decimal.

(b) $5 \div 11 = 0.454545...$ It is a rational number since it is a repeating decimal.

(c) $2 \div 5 = 0.400...$ It is a rational number since it is a repeating decimal (0 is the repeating part).

58. $124 \div 990 = 0.12525....$ the calculator may show the answer as 0.1252525253 because it has rounded up for the next 5 that doesn't fit on the screen.

59. $32.4 \text{ MJ} + 26.704 \text{ MJ} + 36.23 \text{ MJ} = 95.334 \text{ MJ}$. The answer must be to the same precision as the least precise measurement. The answer is 95.3 MJ.

60. We would compute $8(68.6) + 5(15.3) = 625.3$ and round to three significant digits for a total weight of 625 lb. The values 8 and 5 are exact.

61. We would compute $12(129) + 16(298.8) = 6328.8$ and round to three significant digits for a total weight of 6330 g. The values 12 and 16 are exact.

62. $V = (15.2\ \Omega + 5.64\ \Omega + 101.23\ \Omega) \times 3.55 \text{ A}$

$V = 122.07\ \Omega \times 3.55 \text{ A}$

$V = 433.3485 \text{ V}$

$V = 433 \text{ V}$ to 3 significant digits

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

63. $\dfrac{100(40.63+52.96)}{105.30+52.96} = 59.1386\ \% = 59.14\ \%$ to 4 signficiant digits

64. $T = \dfrac{50.45(9.80)}{1+100.9\div 23} = 91.779\ \text{N} = 92\ \text{N}$ to 2 significant digits

65. **(a)** Estimate $8\times 5-10=30,$ to 1 significant digit.
(b) Calculator: $7.84\times 4.932-11.317=27.34988$ which is 27.3 to 3 significant digits.

66. **(a)** Estimate $20-50\div 10=15$ to 2 significant digit.
(b) Calculator: $21.6-53.14\div 9.64=16.0875519$ which is 16.1 to 3 significant digits.

1.4 Exponents and Unit Conversions

1. $(-x^3)^2=\left[(-1)x^3\right]^2=(-1)^2(x^3)^2=(1)x^6=x^6$

2. $2x^0=2(1)=2$

3. $x^3x^4=x^{3+4}=x^7$

4. $y^2y^7=y^{2+7}=y^9$

5. $2b^4b^2=2b^{4+2}=2b^6$

6. $3k^5k=3k^{5+1}=3k^6$

7. $\dfrac{m^5}{m^3}=m^{5-3}=m^2$

8. $\dfrac{2x^6}{-x}=-2x^{6-1}=-2x^5$

9. $\dfrac{-n^5}{7n^9}=-\dfrac{n^{5-9}}{7}=-\dfrac{n^{-4}}{7}=-\dfrac{1}{7n^4}$

10. $\dfrac{3s}{s^4}=3s^{1-4}=3s^{-3}=\dfrac{3}{s^3}$

11. $\left(P^2\right)^4=P^{2(4)}=P^8$

12. $\left(x^8\right)^3=x^{8(3)}=x^{24}$

13. $\left(aT^2\right)^{30}=a^{30}T^{2(30)}=a^{30}T^{60}$

14. $\left(3r^2\right)^3=(3)^3r^{2(3)}=27r^6$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

15. $\left(\frac{2}{b}\right)^3 = \frac{(2)^3}{b^3} = \frac{8}{b^3}$

16. $\left(\frac{F}{t}\right)^{20} = \frac{F^{20}}{t^{20}}$

17. $\left(\frac{x^2}{-2}\right)^4 = \frac{x^{2(4)}}{(-2)^4} = \frac{x^8}{16}$

18. $\left(\frac{3}{n^3}\right)^3 = \frac{(3)^3}{n^{3(3)}} = \frac{27}{n^9}$

19. $(8a)^0 = 1$

20. $-v^0 = -1$

21. $-3x^0 = -3(1) = -3$

22. $-(-2)^0 = -1(1) = -1$

23. $6^{-1} = \frac{1}{6^1} = \frac{1}{6}$

24. $-w^{-5} = -\frac{1}{w^5}$

25. $\frac{1}{R^{-2}} = R^2$

26. $\frac{1}{-t^{-48}} = -t^{48}$

27. $(-t^2)^7 = \left[(-1)(t^2)\right]^7 = (-1)^7 t^{2(7)} = (-1)t^{14} = -t^{14}$

28. $(-y^3)^5 = \left[(-1)(y^3)\right]^5 = (-1)^5 y^{3(5)} = (-1)y^{15} = -y^{15}$

29. $-\frac{L^{-3}}{L^{-5}} = -L^{-3-(-5)} = -L^2$

30. $2i^{40}i^{-70} = 2i^{40+(-70)} = 2i^{-30} = \frac{2}{i^{30}}$

31. $\frac{2v^4}{(2v)^4} = \frac{2v^4}{(2)^4(v^4)} = \frac{2v^4}{16v^4} = \frac{1}{8}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

32. $\dfrac{x^2x^3}{(x^2)^3}=\dfrac{x^{2+3}}{x^{2(3)}}=\dfrac{x^5}{x^6}=\dfrac{1}{x}$

33. $\dfrac{(n^2)^4}{(n^4)^2}=\dfrac{n^{2(4)}}{n^{4(2)}}=\dfrac{n^8}{n^8}=1$

34. $\dfrac{(3t)^{-1}}{3t^{-1}}=\dfrac{(3)^{-1}t^{-1}}{3t^{-1}}=\dfrac{t}{3(3)t}=\dfrac{1}{9}$

35. $(\pi^0x^2a^{-1})^{-1}=\pi^{0(-1)}x^{2(-1)}a^{-1(-1)}=\pi^0x^{-2}a^1=\dfrac{a}{x^2}$

36. $(3m^{-2}n^4)^{-2}=(3)^{-2}m^{-2(-2)}n^{4(-2)}=3^{-2}m^4n^{-8}=\dfrac{m^4}{9n^8}$

37. $(-8g^{-1}s^3)^2=(-8)^2g^{-1(2)}s^{3(2)}=\dfrac{64s^6}{g^2}$

38. $ax^{-2}(-a^2x)^3=ax^{-2}(-1)^3(a^{2(3)})x^3=-\dfrac{a(a^6)x^3}{x^2}=-a^{1+6}x^{3-2}=-a^7x$

39. $\left(\dfrac{4x^{-1}}{a^{-1}}\right)^{-3}=\dfrac{(4)^{-3}x^{-1(-3)}}{a^{-1(-3)}}=\dfrac{x^3}{64a^3}$

40. $\left(\dfrac{2b^2}{y^5}\right)^{-2}=\dfrac{(2)^{-2}b^{2(-2)}}{y^{5(-2)}}=\dfrac{b^{-4}}{4y^{-10}}=\dfrac{y^{10}}{4b^4}$

41. $\dfrac{15n^2T^5}{3n^{-1}T^6}=\dfrac{5n^{2-(-1)}}{T}=\dfrac{5n^3}{T}$

42. $\dfrac{(nRT^{-2})^{32}}{R^{-2}T^{32}}=\dfrac{n^{32}R^{32-(-2)}T^{-2(32)}}{T^{32}}=\dfrac{n^{32}R^{34}T^{-64}}{T^{32}}=\dfrac{n^{32}R^{34}}{T^{32-(-64)}}=\dfrac{n^{32}R^{34}}{T^{96}}$

43. $7(-4)-(-5)^2=-28-25=-53$

44. $6-\left|-2\right|^5-(-2)(8)=6-32-(-16)=6-32+16=-10$

45. $-(-26.5)^2-(-9.85)^3=-(702.25)-(-955.671625)=253.421625$
which gets rounded to 253 because 702.25 and –955.671625 are both accurate to only 3 significant digits due to the original numbers having only 3 significant digits.

46. $-0.711^2-(-\left|-0.809\right|)^6=(-1)(0.711)^2-(-0.809)^6=(-1)(0.505521)-(0.2803439122)=-0.7858649122$
which gets rounded to 3 significant digits: –0.786.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

47. $\dfrac{3.07(-|-1.86|)}{(-1.86)^4+1.596}=\dfrac{-5.7102}{11.96883216+1.596}=\dfrac{-5.7102}{13.56483216}=-0.420956185$

which gets rounded to 3 significant digits: –0.421.

48. $\dfrac{15.66^2-(-4.017)^4}{1.044(-3.68)}=\dfrac{245.2356-260.379822692}{-3.84192}=\dfrac{-15.144222692}{-3.84192}=3.941837074$

which gets rounded to 3 significant digits: 3.94.

49. $2.38(-60.7)^2-\dfrac{254}{1.17^3}=2.38(3684.49)-\dfrac{254}{1.601613}$

$=8769.0862-158.5901213339$

$=8610.4960786661$

which gets rounded to 3 significant digits: 8610.

50. $4.2(4.6)+\dfrac{0.889}{1.89-1.09^2}$

$=19.32+\dfrac{0.889}{1.89-1.1881}$

$=19.32+\dfrac{0.889}{0.7019}$

$=19.32+0.889880728$

$=20.209880728$

which gets rounded to 2 significant digits: $2\overline{0}$.

51. $\left(\dfrac{1}{x^{-1}}\right)^{-1}=\dfrac{1^{-1}}{x^{-1(-1)}}=\dfrac{1}{x}$, which is the reciprocal of x.

52. $\left(\dfrac{0.2-5^{-1}}{10^{-2}}\right)^0=\left(\dfrac{0.2-\frac{1}{5}}{\frac{1}{100}}\right)^0=\left(\dfrac{0}{0.01}\right)^0=0^0\neq 1$, since $a^0=1$ requires that $a\neq 0$.

53. If $a^3=5$, then

$a^{12}=a^{3(4)}$

$a^{12}=\left(a^3\right)^4$

$a^{12}=(5)^4$

$a^{12}=625$

54. For any negative value of a, a will be negative, and a^2 will be positive, making all values of $\dfrac{1}{a^2}$ greater than $\dfrac{1}{a}$. Therefore, it is never the case for negative values of a, $a^{-2}<a^{-1}$.

55. $(x^a\cdot x^{-a})^5=(x^{a-a})^5=(x^0)^5=x^{0(5)}=x^0=1$, provided that $x\neq 0$.

56. $(-y^{a-b}\cdot y^{a+b})^2=((-1)y^{a-b+(a+b)})^2=(-1)^2(y^{2a})^2=y^{2a(2)}=y^{4a}$.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

57. $$\left(\frac{kT}{hc}\right)^3 (GkThc)^2 c = \frac{k^3T^3}{h^3c^3}\cdot(G^2k^2T^2h^2c^2)c$$
$$= \frac{k^3T^3}{h^3c^3}\cdot(G^2k^2T^2h^2c^3)$$
$$= \frac{(G^2k^{2+3}T^{2+3}c^{3-3})}{h^1}$$
$$= \frac{G^2k^5T^5}{h}$$

58. $$GmM(mr)^{-1}(r^{-2}) = \frac{GmM}{mr^{1+2}} = \frac{GM}{r^3}$$

59. $$\pi\left(\frac{r}{2}\right)^3\left(\frac{4}{3\pi r^2}\right) = \pi\left(\frac{r^3}{8}\right)\left(\frac{4}{3\pi r^2}\right) = \frac{4r}{24} = \frac{r}{6}$$

60. $$\frac{gM(2\pi fM)^{-2}}{2\pi fC} = \frac{gM}{2\pi fC(2\pi fM)^2}$$
$$= \frac{gM}{2\pi fC(4\pi^2 f^2 M^2)}$$
$$= \frac{gM}{8\pi^3 f^3 CM^2}$$
$$= \frac{g}{8\pi^3 f^3 CM}$$

61. $$2500\left(1+\frac{0.042}{4}\right)^{24} = \$2500(1.0105)^{24}$$
$$= \$2500(1.28490602753)$$
$$= \$3212.26700688$$
$$= \$3212.27$$

62. $$\frac{6.85(1000-20(6.85)^2+(6.85)^3)}{1850} = \frac{6.85(1000-20(46.9225)+321.419125)}{1850}$$
$$= \frac{6.85(1321.419125-938.45)}{1850}$$
$$= \frac{6.85(382.969125)}{1850}$$
$$= \frac{2623.33850625}{1850}$$
$$= 1.418020814$$
$$= 1.42 \text{ cm}$$

63. If $f(n) = \left(1+\frac{1}{n}\right)^n$ then $f(1) = 2^1 = 2.000$, $f(10) = 1.1^{10} = 2.594$,
$f(100) = 1.01^{100} = 2.705$, and $f(1000) = 1.001^{1000} = 2.717$.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

64. We have 1 TB $= 2^{10}$ GB $= 2^{10}(2^{10}$ MB$) = 2^{10}(2^{10}(2^{20}$ bytes$)) = 2^{10+10+20}$ bytes $= 2^{40}$ bytes

65. $\left(28.2\dfrac{\text{ft}}{\text{s}}\right)(9.81\text{ s}) = 276.642$ ft which is rounded to 277 ft.

66. $\left(40.5\dfrac{\text{mi}}{\text{gal}}\right)(3.7\text{ gal}) = 149.85$ mi which is rounded to 150 mi.

67. $\left(7.25\dfrac{\text{m}}{\text{s}^2}\right)\left(\dfrac{1\text{ ft}}{0.3048\text{ m}}\right)\left(\dfrac{60\text{ s}}{1\text{ min}}\right)^2 = 85,629.92\ \dfrac{\text{ft}}{\text{min}^2}$ which is rounded to 85,600 $\dfrac{\text{ft}}{\text{min}^2}$.

68. $\left(238\dfrac{\text{kg}}{\text{m}^3}\right)\left(\dfrac{1000\text{ g}}{1\text{ kg}}\right)\left(\dfrac{1\text{ m}}{100\text{ cm}}\right)^3 = 0.238\ \dfrac{\text{g}}{\text{cm}^3}$.

69. $15.7\text{ qt} = 15.7\text{ qt}\times\left(\dfrac{1\text{ L}}{1.057\text{ qt}}\right) = 14.8533586$ L which is rounded to 14.9 L.

70. $7.50\text{ W} = 7.50\text{ W}\times\left(\dfrac{1\text{ hp}}{746.0\text{ W}}\right) = 0.01005362$ hp which is rounded to 0.0101 hp.

71. $245\text{ cm}^2 = 245\text{ cm}^2\times\left(\dfrac{1\text{ in}}{2.54\text{ cm}}\right)^2 = 37.975076\text{ in}^2$ which is rounded to 38.0 in^2.

72. $85.7\text{ mi}^2 = 85.7\text{ mi}^2\times\left(\dfrac{1\text{ km}}{0.6214\text{ mi}}\right)^2 = 221.941401\text{ km}^2$ which is rounded to 222 km^2.

73. $65.2\dfrac{\text{m}}{\text{s}} = 65.2\dfrac{\text{m}}{\text{s}}\times\left(\dfrac{60\text{ s}}{1\text{ min}}\right)\times\left(\dfrac{1\text{ ft}}{0.3048\text{ m}}\right) = 12834.6457\ \dfrac{\text{ft}}{\text{min}}$ which is rounded to 12800 $\dfrac{\text{ft}}{\text{min}}$.

74. $25.0\dfrac{\text{mi}}{\text{gal}} = 25.0\dfrac{\text{mi}}{\text{gal}}\times\left(\dfrac{1\text{ km}}{0.6214\text{ mi}}\right)\times\left(\dfrac{1\text{ gal}}{3.785\text{ L}}\right) = 10.6292562\ \dfrac{\text{km}}{\text{L}}$ which is rounded to 10.6 $\dfrac{\text{km}}{\text{L}}$.

75. $15.6\text{ in} = 15.6\text{ in}\times\left(\dfrac{2.54\text{ cm}}{1\text{ in}}\right) = 39.624$ cm which is rounded to 39.6 cm.

76. $12,500\text{ mi} = 12,500\text{ mi}\times\left(\dfrac{1\text{ km}}{0.6214\text{ mi}}\right) = 20,115.8674$ km which is rounded to 20,100 km.

77. $575,000\dfrac{\text{gal}}{\text{day}} = 575,000\dfrac{\text{gal}}{\text{day}}\times\left(\dfrac{1\text{ day}}{24\text{ hr}}\right)\times\left(\dfrac{3.785\text{ L}}{1\text{ gal}}\right) = 90,682.2917\ \dfrac{\text{L}}{\text{hr}}$ which is rounded to 90,700 $\dfrac{\text{L}}{\text{hr}}$.

78. $85\dfrac{\text{gal}}{\text{min}} = 85\dfrac{\text{gal}}{\text{min}}\times\left(\dfrac{1\text{ min}}{60\text{ s}}\right)\times\left(\dfrac{3.785\text{ L}}{1\text{ gal}}\right) = 5.3620833\ \dfrac{\text{L}}{\text{s}}$ which is rounded to 5.4 $\dfrac{\text{L}}{\text{s}}$.

79. $1130\dfrac{\text{ft}}{\text{s}} = 1130\dfrac{\text{ft}}{\text{s}}\times\left(\dfrac{60\text{ s}}{1\text{ min}}\right)\times\left(\dfrac{60\text{ min}}{1\text{ hr}}\right)\times\left(\dfrac{0.3084\text{ m}}{1\text{ ft}}\right)\times\left(\dfrac{1\text{ km}}{1000\text{ m}}\right) = 1254.5712\ \dfrac{\text{km}}{\text{hr}}$ which is rounded to 1250 $\dfrac{\text{km}}{\text{hr}}$.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

80. $7200\dfrac{\text{km}}{\text{hr}} = 7200\dfrac{\text{km}}{\text{hr}} \times \left(\dfrac{1\text{ hr}}{60\text{ min}}\right) \times \left(\dfrac{1\text{ min}}{60\text{ s}}\right) \times \left(\dfrac{1000\text{ m}}{1\text{ km}}\right) = 2000\ \dfrac{\text{m}}{\text{s}}.$

81. $14.7\dfrac{\text{lb}}{\text{in}^2} = 14.7\dfrac{\text{lb}}{\text{in}^2} \times \left(\dfrac{4.448\text{ N}}{1\text{ lb}}\right) \times \left(\dfrac{1\text{ in}}{2.54\text{ cm}}\right)^2 \times \left(\dfrac{100\text{ cm}}{1\text{ m}}\right)^2 = 101{,}347.883\ \dfrac{\text{N}}{\text{m}^2}$ which is rounded to 101,000 Pa.

82. $62.4\dfrac{\text{lb}}{\text{ft}^3} = 62.4\dfrac{\text{lb}}{\text{ft}^3} \times \dfrac{1\text{ kg}}{2.205\text{ lb}} \times \left(\dfrac{1\text{ ft}}{0.3048\text{ m}}\right)^3 = 999.381\ \dfrac{\text{kg}}{\text{m}^3}$ which is rounded to 999 $\dfrac{\text{kg}}{\text{m}^3}$.

(The actual value is 1000 $\dfrac{\text{kg}}{\text{m}^3}$.)

1.5 Scientific Notation

1. $8.06\times10^3 = 8060$

2. $750\ 000\ 000\ 000^{-1} = (7.5\times10^{11})^{-1}$
$= 7.5^{-1}\times10^{-11}$
$= 0.1333...\times10^{-11}$
$= 1.33\times10^{-12}$
rounded to 3 significant digits.

3. $4.5\times10^4 = 45{,}000$

4. $6.8\times10^7 = 68{,}000{,}000$

5. $2.01\times10^{-3} = 0.00201$

6. $9.61\times10^{-5} = 0.0000961$

7. $3.23\times10^0 = 3.23\times1 = 3.23$

8. $8\times10^0 = 8\times1 = 8$

9. $1.86\times10 = 18.6$

10. $1\times10^{-1} = 0.1$

11. $4000 = 4\times10^3$

12. $56\ 000 = 5.6\times10^4$

13. $0.0087 = 8.7\times10^{-3}$

14. $0.00074 = 7.4\times10^{-4}$

15. $609{,}000{,}000 = 6.09\times10^8$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

16. $10 = 1\times10^{1}$

17. $0.0528 = 5.28\times10^{-2}$

18. $0.0000908 = 9.08\times10^{-5}$

19. $28{,}000(2{,}000{,}000{,}000) = 2.8\times10^{4}(2\times10^{9}) = 5.6\times10^{13}$

20. $50{,}000(0.006) = 5\times10^{4}(6\times10^{-3}) = 300 = 3\times10^{2}$

21. $\dfrac{88{,}000}{0.0004} = \dfrac{8.8\times10^{4}}{4\times10^{-4}} = 2.2\times10^{8}$

22. $\dfrac{0.00003}{6{,}000{,}000} = \dfrac{3\times10^{-5}}{6\times10^{6}} = 5\times10^{-12}$

23. $35{,}600{,}000 = 35.6\times10^{6}$

24. $0.0000056 = 5.6\times10^{-6}$

25. $0.0973 = 97.3\times10^{-3}$

26. $925{,}000{,}000{,}000 = 925\times10^{9}$

27. $0.000000475 = 475\times10^{-9}$

28. $370{,}000 = 370\times10^{3}$

29. $2\times10^{-35} + 3\times10^{-34} = 0.2\times10^{-34} + 3\times10^{-34} = 3.2\times10^{-34}$

30. $5.3\times10^{12} - 3.7\times10^{10} = 530\times10^{10} - 3.7\times10^{10} = 526.3\times10^{10} = 5.263\times10^{12}$

31. $(1.2\times10^{29})^{3} = 1.2^{3}\times10^{29(3)} = 1.728\times10^{87}$

32. $(2\times10^{-16})^{-5} = 2^{-5}\times10^{-16(-5)} = 0.031\ 25\times10^{80} = 3.125\times10^{78}$

33. $1320(649{,}000)(85.3) = 7.3074804\times10^{10}$
which gets rounded to 7.31×10^{10}.

34. $0.0000569(3{,}190{,}000) = 181.511$
which gets rounded to 1.82×10^{2}.

35. $\dfrac{0.0732(6710)}{0.00134(0.0231)} = \dfrac{491.172}{0.000030954} = 1.5867803\times10^{7}$
which gets rounded to 1.59×10^{7}.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

36. $\dfrac{0.00452}{2430(97{,}100)} = \dfrac{0.00452}{235{,}953{,}000} = 1.915635741\times10^{-11}$

which gets rounded to 1.92×10^{-11}.

37. $(3.642\times10^{-8})(2.736\times10^{5}) = 9.964512\times10^{-3}$

which gets rounded to 9.965×10^{-3}.

38. $\dfrac{(7.309\times10^{-1})^2}{5.9843(2.5036\times10^{-20})} = \dfrac{0.53421481}{1.49787029\times10^{-19}} = 3.56656723394\times10^{18}$

which gets rounded to 3.567×10^{18}.

39. $\dfrac{(3.69\times10^{-7})(4.61\times10^{21})}{0.0504} = \dfrac{1.70109\times10^{15}}{0.0504} = 3.37517857142\times10^{16}$

which gets rounded to 3.38×10^{16}.

40. $\dfrac{(9.907\times10^{7})(1.08\times10^{12})^2}{(3.603\times10^{-5})(2054)} = \dfrac{(9.907\times10^{7})(1.1664\times10^{24})}{0.07400562} = \dfrac{1.01555248\times10^{32}}{0.07400562} = 1.56143882045\times10^{33}$

which gets rounded to 1.56×10^{33}.

41. $500{,}000{,}000 \text{ tweets} = 5\times10^{8} \text{ tweets}$

42. $17{,}200{,}000{,}000 \text{ bytes} = 1.72\times10^{10} \text{ bytes}$

43. $0.000003 \text{ W} = 3\times10^{-6} \text{ W}$

44. $0.0075 \text{ mm} = 7.5\times10^{-3} \text{ mm}$

45. $1{,}200{,}000{,}000 \text{ Hz} = 1.2\times10^{9} \text{ Hz}$

46. $1.84\times10^{12} = 1{,}840{,}000{,}000{,}000$

47. $12{,}000{,}000{,}000 \text{ m}^2 = 1.2\times10^{10} \text{ m}^2$

48. $3.086\times10^{16} \text{ m} = 30{,}860{,}000{,}000{,}000{,}000 \text{ m}$

49. $1.6\times10^{-12} \text{ W} = 0.0000000000016 \text{ W}$

50. $2.4\times10^{-43} = 0.000000000000000000000000000000000000000000024$

51. **(a)** $2300 = 2.3\times10^{3} = 2.3 \text{ kW}$
(b) $0.23 = 230\times10^{-3} = 230 \text{ mW}$
(c) $2{,}300{,}000 = 2.3\times10^{6} = 2.3 \text{ MW}$
(d) $0.00023 = 230\times10^{-6} = 230\ \mu\text{W}$

52. **(a)** $8090000 = 8.09\times10^{6} = 8.09 \text{ M}\Omega$
(b) $809\ 000 = 809\times10^{3} = 809 \text{ k}\Omega$
(c) $0.0809 = 80.9\times10^{-3} = 80.9 \text{ m}\Omega$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

53. **(a)** $\text{googol} = 1\times10^{100} = 10^{100}$

(b) $\text{googolplex} = 10^{\text{googol}} = 10^{10^{100}}$

54. $\text{googol} = 10^{100}$, so to find the ratio $\dfrac{10^{100}}{10^{79}} = 10^{100-79} = 10^{21}$

A googol is 10^{21} times larger than the number of electrons in the universe.

55. $\text{earth's diameter} = \dfrac{\text{sun's diameter}}{110} = \dfrac{1.4\times10^{9}\ \text{m}}{110} = 1.27272\times10^{7}\ \text{m}$ which is rounded to $1.3\times10^{7}\ \text{m}$.

56. $2^{30} = 1,073,741,824 = 1.073741824\times10^{9} \approx 1\times10^{9}$

57. $\dfrac{7.5\times10^{-15}\,\text{s}}{\text{addition}}\times5.6\times10^{6}\,\text{additions} = 4.2\times10^{-8}\,\text{s}$

58. $0.000000039\ \% = 0.00000000039$

$0.00000000039\times0.085\ \text{mg} = 3.315\times10^{-11}\,\text{mg} = 3.3\times10^{-11}\,\text{mg}$

59. $0.078\ \text{s}\times2.998\times10^{8}\,\dfrac{\text{m}}{\text{s}} = 2.3384400\times10^{7}\ \text{m}$ which rounds to $2.3\times10^{7}\ \text{m}$

60. **(a)** $1\ \text{day}\times\dfrac{24\ \text{h}}{\text{day}}\times\dfrac{60\ \text{min}}{\text{h}}\times\dfrac{60\ \text{s}}{\text{min}} = 86400\ \text{s} = 8.64\times10^{4}\ \text{s}$

(b) $100\ \text{year}\times\dfrac{365.25\ \text{day}}{\text{year}}\times\dfrac{24\ \text{h}}{\text{day}}\times\dfrac{60\ \text{min}}{\text{h}}\times\dfrac{60\ \text{s}}{\text{min}} = 3\ 155\ 760\ 000\ \text{s} = 3.155\ 760\ 0\times10^{9}\ \text{s}$

61. $\dfrac{1.66\times10^{-27}\ \text{kg}}{\text{amu}}\times\dfrac{1.6\times10^{1}\ \text{amu}}{\text{oxygen atoms}}\times1.25\times10^{8}\ \text{oxygen atoms} = 3.32\times10^{-18}\ \text{kg}$

62. $W = kT^4$

$W = 5.7\times10^{-8}\ \text{W/K}^4\times(3.03\times10^{2}\ \text{K})^4$

$W = 5.7\times10^{-8}\ \text{W/K}^4\ \times\ 8.428892481\times10^{9}\ \text{K}^4$

$W = 4.80446871417\times10^{2}\ \text{W}$

$W = 4.8\times10^{2}\ \text{W}$

63. $R = \dfrac{k}{d^2} = \dfrac{2.196\times10^{-8}\ \Omega\cdot\text{m}^2}{\left(7.998\times10^{-5}\text{m}\right)^2} = \dfrac{2.196\times10^{-8}\ \Omega\cdot\text{m}^2}{6.396\ 800\ 4\times10^{-9}\text{m}^2} = 3.432\ 966\ 268\ 57\ \Omega = 3.433\ \Omega$

64. $\dfrac{1.496\times10^{8}\,\text{km}}{\text{AU}}\times\dfrac{\text{AU}}{4.99\times10^{2}\,\text{s}} = 2.99799599198\times10^{5}\ \text{km/s} = 2.998\times10^{5}\ \text{km/s}$

This is the same speed mentioned in Question 56 as the speed of radio waves.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

1.6 Roots and Radicals

1. $-\sqrt[3]{64} = -\sqrt[3]{(4)^3} = -4$

2. $\sqrt{(15)(5)}$
Neither 15 nor 5 is a perfect square, so this expression is not as useful. However, if we further factor the 15 to $\sqrt{(3)(5)(5)} = \sqrt{3(5)^2} = 5\sqrt{3}$, the result can still be obtained.

3. $\sqrt{16\times 9} = \sqrt{144} = \sqrt{12^2} = 12$

4. $-\sqrt{-64}$ is still imaginary because an even root (in this case $n = 2$) of a negative number is imaginary, regardless of the numerical factor placed in front of the root.

5. $\sqrt{49} = \sqrt{7^2} = 7$

6. $\sqrt{225} = \sqrt{(25)(9)} = \sqrt{25}\times\sqrt{9} = 5\times 3 = 15$

7. $-\sqrt{121} = -\sqrt{11^2} = -11$

8. $-\sqrt{36} = -\sqrt{6^2} = -6$

9. $-\sqrt{64} = -\sqrt{8^2} = -8$

10. $\sqrt{0.25} = \sqrt{\frac{1}{4}} = \frac{\sqrt{1}}{\sqrt{4}} = \frac{1}{2} = 0.5$

11. $\sqrt{0.09} = \sqrt{\frac{9}{100}} = \frac{\sqrt{9}}{\sqrt{100}} = \frac{3}{10} = 0.3$

12. $-\sqrt{900} = -\sqrt{(9)(100)} = -\sqrt{9}\times\sqrt{100} = -3\times 10 = -30$

13. $\sqrt[3]{125} = \sqrt[3]{5^3} = 5$

14. $\sqrt[4]{16} = \sqrt[4]{2^4} = 2$

15. $\sqrt[4]{81} = \sqrt[4]{3^4} = 3$

16. $-\sqrt[5]{-32} = -\sqrt[5]{(-2)^5} = -(-2) = 2$

17. $\left(\sqrt{5}\right)^2 = \sqrt{5}\times\sqrt{5} = 5$

18. $\left(\sqrt[3]{31}\right)^3 = \sqrt[3]{31}\times\sqrt[3]{31}\times\sqrt[3]{31} = 31$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

19. $\left(-\sqrt[3]{-47}\right)^3 = (-1)^3\left(\sqrt[3]{-47}\right)^3 = (-1)(-47) = 47$

20. $\left(\sqrt[5]{-23}\right)^5 = -23$

21. $\left(-\sqrt[4]{53}\right)^4 = (-1)^4\left(\sqrt[4]{53}\right)^4 = (1)(53) = 53$

22. $\sqrt{75} = \sqrt{(25)(3)} = \sqrt{25}\times\sqrt{3} = 5\sqrt{3}$

23. $\sqrt{18} = \sqrt{(9)(2)} = \sqrt{9}\times\sqrt{2} = 3\sqrt{2}$

24. $-\sqrt{32} = -\sqrt{(16)(2)} = -\sqrt{16}\times\sqrt{2} = -4\sqrt{2}$

25. $\sqrt{1200} = \sqrt{(100)(4)(3)} = \sqrt{100}\times\sqrt{4}\times\sqrt{3} = 10\times 2\times\sqrt{3} = 20\sqrt{3}$

26. $\sqrt{50} = \sqrt{(25)(2)} = \sqrt{25}\times\sqrt{2} = 5\times\sqrt{2} = 5\sqrt{2}$

27. $2\sqrt{84} = 2\sqrt{(4)(21)} = 2\times\sqrt{4}\times\sqrt{21} = 2\times 2\times\sqrt{21} = 4\sqrt{21}$

28. $\dfrac{\sqrt{108}}{2} = \dfrac{\sqrt{(36)(3)}}{2} = \dfrac{\sqrt{36}\times\sqrt{3}}{2} = \dfrac{6\times\sqrt{3}}{2} = 3\sqrt{3}$

29. $\sqrt{\dfrac{80}{|3-7|}} = \sqrt{\dfrac{80}{4}} = \sqrt{20} = \sqrt{4\times 5} = \sqrt{4}\times\sqrt{5} = 2\times\sqrt{5} = 2\sqrt{5}$

30. $\sqrt{81\times 10^2} = \sqrt{81}\times\sqrt{10^2} = 9\times 10 = 90$

31. $\sqrt[3]{-8^2} = \sqrt[3]{-64} = \sqrt[3]{(-4)^3} = -4$

32. $\sqrt[4]{9^2} = \sqrt[4]{81} = \sqrt[4]{3^4} = 3$

33. $\dfrac{7^2\sqrt{81}}{(-3)^2\sqrt{49}} = \dfrac{(49)(9)}{(9)(7)} = \dfrac{(49)\cancel{(9)}}{\cancel{(9)}\,(7)} = 7$

34. $\dfrac{2^5\sqrt[5]{243}}{-3\sqrt{144}} = -\dfrac{32\sqrt[5]{3^5}}{3\sqrt{12^2}} = -\dfrac{(32)\cancel{(3)}}{\cancel{(3)}\,(12)} = -\dfrac{8}{3}$

35. $\sqrt{36+64} = \sqrt{100} = \sqrt{10^2} = 10$

36. $\sqrt{25+144} = \sqrt{169} = \sqrt{13^2} = 13$

37. $\sqrt{3^2+9^2} = \sqrt{9+81} = \sqrt{90} = \sqrt{(9)(10)} = \sqrt{9}\times\sqrt{10} = 3\sqrt{10}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

38. $\sqrt{8^2-4^2} = \sqrt{64-16} = \sqrt{48} = \sqrt{(16)(3)} = \sqrt{16}\times\sqrt{3} = 4\sqrt{3}$

39. $\sqrt{85.4} = 9.24121204171$,which is rounded to 9.24

40. $\sqrt{3762} = 61.3351449007$,which is rounded to 61.34

41. $\sqrt{0.8152} = 0.9028842672$,which is rounded to 0.9029

42. $\sqrt{0.0627} = 0.25039968051$,which is rounded to 0.250

43. **(a)** $\sqrt{1296+2304} = \sqrt{3600} = 60$, which is expressed as 60.00
(b) $\sqrt{1296}+\sqrt{2304} = 36+48 = 84$, which is expressed as 84.00

44. **(a)** $\sqrt{10.6276+2.1609} = \sqrt{12.7885} = 3.57610122899$,which is rounded to 3.57610
(b) $\sqrt{10.6276}+\sqrt{2.1609} = 3.26+1.47 = 4.73$,which is expressed as 4.7300

45. **(a)**
$$\begin{aligned}\sqrt{0.0429^2-0.0183^2} &= \sqrt{0.00184041-0.00033489}\\ &= \sqrt{0.00150552}\\ &= 0.03880103091\\ &= 0.0388\end{aligned}$$
(b)
$$\begin{aligned}\sqrt{0.0429^2}-\sqrt{0.0183^2} &= 0.0429-0.0183\\ &= 0.0246\end{aligned}$$

46. **(a)**
$$\begin{aligned}\sqrt{3.625^2+0.614^2} &= \sqrt{13.140625+0.376996}\\ &= \sqrt{13.517621}\\ &= 3.67663174658\\ &= 3.677\end{aligned}$$
(b)
$$\begin{aligned}\sqrt{3.625^2}+\sqrt{0.614^2} &= 3.625+0.614\\ &= 4.239\end{aligned}$$

47. $\sqrt{24s} = \sqrt{(24)(150)} = \sqrt{3600} = 60$ mi/h

48.
$$\begin{aligned}\sqrt{Z^2-X^2} &= \sqrt{(5.362\ \Omega)^2-(2.875\ \Omega)^2}\\ &= \sqrt{28.751044\ \Omega^2-8.265625\ \Omega^2}\\ &= \sqrt{20.485419\ \Omega^2}\\ &= 4.52608208056\ \Omega\\ &= 4.526\ \Omega\end{aligned}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

49. $$\sqrt{\frac{B}{d}} = \sqrt{\frac{2.18\times10^{9}\ \text{Pa}}{1.03\times10^{3}\ \text{kg/m}^3}}$$
$$= \sqrt{2116504.85436\ \frac{\text{N/m}^2}{\text{kg/m}^3}}$$
$$= \sqrt{2116504.85436\ \frac{\left(\text{kg}\cdot\text{m/s}^2\right)/\text{m}^2}{\text{kg/m}^3}}$$
$$= \sqrt{2116504.85436\ \text{m}^2/\text{s}^2}$$
$$= 1454.82124481\ \text{m/s}$$
$$= 1450\ \text{m/s}$$

50. $$\sqrt{40m} = \sqrt{(40)(75)}$$
$$= \sqrt{3000}$$
$$= 54.7722557505$$
$$= 55\ \text{m/s}$$

51. $$\sqrt{w^2+h^2} = \sqrt{(52.3\ \text{in})^2 + (29.3\ \text{in})^2}$$
$$= \sqrt{2735.29\ \text{in}^2 + 858.49\ \text{in}^2}$$
$$= \sqrt{3593.78\ \text{in}^2}$$
$$= 59.948144258\ \text{in}$$
$$= 59.9\ \text{in}$$

52. $$100\left(1-\sqrt{\frac{V}{C}}\right) = 100\left(1-\sqrt{\frac{24000}{38000}}\right)$$
$$= 100\left(1-\sqrt{0.63157895}\right)$$
$$= 100\left(1-0.79471941589\right)$$
$$= 100(0.2052805841)$$
$$= 20.52805841\ \%$$
$$= 21\ \%$$

53. $$\sqrt{gd} = \sqrt{(9.8)(3500)}$$
$$= \sqrt{34300}$$
$$= 185.20259$$
$$= 190\ \text{m/s}$$

54. $$\sqrt{1.27\times10^4 h + h^2} = \sqrt{1.27\times10^4(9500) + (9500)^2}$$
$$= \sqrt{1.2065\times10^8 + 9.025\times10^7}$$
$$= \sqrt{2.109\times10^8}$$
$$= 14522.3965$$
which is rounded to 15000 km

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

55. $\sqrt{a^2} = a$ is not necessarily true for negative values of a because a^2 will be a positive number, regardless whether a is negative or positive.The principal root calculated is assumed to be positive, but there are always two solutions to a square root, $\sqrt{a^2} = \pm a$ since $(+a)^2 = a^2$ and $(-a)^2 = a^2$ (see the introduction to this chapter section),so it is sometimes true and sometimes false for negative values of a, depending on which root solution is desired. If *only principal roots* are considered, then it will *not* be true for negative values of a. For example, $\sqrt{(-4)^2} = \sqrt{16} = 4 \neq -4$.


56. **(a)** $x > \sqrt{x}$ when $x > 1$. Any number greater than 1 will have a square root that is smaller than itself. For example, $2 > \sqrt{2} = 1.41$

(b) $x = \sqrt{x}$ when $x = 1$ or $x = 0$ because the only numbers that are their own squares are 0 and 1 (i.e., $0^2 = 0$ and $1^2 = 1$).

(c) $x < \sqrt{x}$ when $0 < x < 1$. Any number between 0 and 1 will have a square root larger than itself. For example, $0.25 < \sqrt{0.25} = 0.5$


57. **(a)** $\sqrt[3]{2140} = 12.8865874254$,which is rounded to 12.9

(b) $\sqrt[3]{-0.214} = -0.59814240297$,which is rounded to –0.598



58. **(a)** $\sqrt[7]{0.382} = 0.87155493458$,which is rounded to 0.872

(b) $\sqrt[7]{-382} = -2.33811675837$,which is rounded to –2.34



59. $f = \dfrac{1}{2\pi\sqrt{LC}} = \dfrac{1}{2(3.1416)\sqrt{0.250(40.52 \times 10^{-6})}}$

$= \dfrac{1}{6.2832\sqrt{10.0625 \times 10^{-6}}}$

$= \dfrac{1}{6.2832(0.003172144385)}$

$= \dfrac{1}{0.0199312175998}$

$= 50.172549$

which is rounded to 50.2 Hz

60. standard deviation = $\sqrt{\text{variance}}$

$= \sqrt{80.5 \text{ kg}^2}$

$= 8.972179222$ kg

which is rounded to 8.97 kg

1.7 Addition and Subtraction of Algebraic Expressions

1. $3x + 2y - 5y = 3x - 3y$

2. $3c - (2b - c) = 3c - 2b + c = -2b + 4c$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

3. $3ax-[(ax-5s)-2ax]=3ax-[ax-5s-2ax]$
$=3ax-[-ax-5s]$
$=3ax+ax+5s$
$=4ax+5s$

4. $3a^2b-\{a-[2a^2b-(a+2b)]\}=3a^2b-\{a-[2a^2b-a-2b]\}$
$=3a^2b-\{a-2a^2b+a+2b\}$
$=3a^2b-\{2a-2a^2b+2b\}$
$=3a^2b-2a+2a^2b-2b$
$=5a^2b-2a-2b$

5. $5x+7x-4x=8x$

6. $6t-3t-4t=-t$

7. $2y-y+4x=y+4x$

8. $-4C+L-6C=-10C+L$

9. $3t-4s-3t-s=0t-5s=-5s$

10. $-8a-b+12a+b=4a+0b=4a$

11. $2F-2T-2+3F-T=5F-3T-2$

12. $x-2y-3x-y+z=-2x-3y+z$

13. $a^2b-a^2b^2-2a^2b=-a^2b-a^2b^2$

14. $-xy^2-3x^2y^2+2xy^2=xy^2-3x^2y^2$

15. $2p+(p-6-2p)=2p-6-p=p-6$

16. $5+(3-4n+p)=5+3-4n+p=-4n+p+8$

17. $v-(7-9x+2v)=v-7+9x-2v=-v+9x-7$

18. $-2a-\frac{1}{2}(b-a)=-2a-\frac{1}{2}b+\frac{1}{2}a=-\frac{3}{2}a-\frac{1}{2}b$

19. $2-3-(4-5a)=-1-4+5a=5a-5$

20. $\sqrt{A}+(h-2\sqrt{A})-3\sqrt{A}=\sqrt{A}+h-2\sqrt{A}-3\sqrt{A}=-4\sqrt{A}+h$

21. $(a-3)+(5-6a)=a-3+5-6a=-5a+2$

22. $(4x-y)-(-2x-4y)=4x-y+2x+4y=6x+3y$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

23. $-(t-2u)+(3u-t)=-t+2u+3u-t=-2t+5u$

24. $-2(6x-3y)-(5y-4x)=-12x+6y-5y+4x=-8x+y$

25. $3(2r+s)-(-5s-r)=6r+3s+5s+r=7r+8s$

26. $3(a-b)-2(a-2b)=3a-3b-2a+4b=a+b$

27. $-7(6-3j)-2(j+4)=-42+21j-2j-8=19j-50$

28. $-(5t+a^2)-2(3a^2-2st)=-5t-a^2-6a^2+4st=-7a^2+4st-5t$

29.
$$\begin{aligned}-[(4-6n)-(n-3)] &= -[4-6n-n+3] \\ &= -[-7n+7] \\ &= 7n-7\end{aligned}$$

30.
$$\begin{aligned}-[(A-B)-(B-A)] &= -[A-B-B+A] \\ &= -[2A-2B] \\ &= -2A+2B\end{aligned}$$

31.
$$\begin{aligned}2[4-(t^2-5)] &= 2[4-t^2+5] \\ &= 2[-t^2+9] \\ &= -2t^2+18\end{aligned}$$

32.
$$\begin{aligned}-3[-3-\frac{2}{3}(-a-4)] &= -3[-3+\frac{2}{3}a+\frac{8}{3}] \\ &= -3[\frac{2}{3}a-\frac{1}{3}] \\ &= -2a+1\end{aligned}$$

33.
$$\begin{aligned}-2[-x-2a-(a-x)] &= -2[-x-2a-a+x] \\ &= -2[-3a] \\ &= 6a\end{aligned}$$

34.
$$\begin{aligned}-2[-3(x-2y)+4y] &= -2[-3x+6y+4y] \\ &= -2[-3x+10y] \\ &= 6x-20y\end{aligned}$$

35.
$$\begin{aligned}aZ-[3-(aZ+4)] &= aZ-[3-aZ-4] \\ &= aZ-[-aZ-1] \\ &= aZ+aZ+1 \\ &= 2aZ+1\end{aligned}$$

36.
$$\begin{aligned}9v-[6-(-v-4)+4v] &= 9v-[6+v+4+4v] \\ &= 9v-[5v+10] \\ &= 9v-5v-10 \\ &= 4v-10\end{aligned}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

37. $5z-\{8-[4-(2z+1)]\} = 5z-\{8-[4-2z-1]\}$
$= 5z-\{8-4+2z+1\}$
$= 5z-\{5+2z\}$
$= 5z-5-2z$
$= 3z-5$

38. $7y-\{y-[2y-(x-y)]\} = 7y-\{y-[2y-x+y]\}$
$= 7y-\{y-[3y-x]\}$
$= 7y-\{y-3y+x\}$
$= 7y-\{-2y+x\}$
$= 7y+2y-x$
$= -x+9y$

39. $5p-(q-2p)-[3q-(p-q)] = 5p-q+2p-[3q-p+q]$
$= 5p-q+2p-[4q-p]$
$= 7p-q-4q+p$
$= 8p-5q$

40. $-(4-\sqrt{LC})-[(5\sqrt{LC}-7)-(6\sqrt{LC}+2)] = -4+\sqrt{LC}-[5\sqrt{LC}-7-6\sqrt{LC}-2]$
$= -4+\sqrt{LC}-[-\sqrt{LC}-9]$
$= -4+\sqrt{LC}+\sqrt{LC}+9$
$= 2\sqrt{LC}+5$

41. $-2\{-(4-x^2)-[3+(4-x^2)]\} = -2\{-4+x^2-[3+4-x^2]\}$
$= -2\{-4+x^2-3-4+x^2\}$
$= -2\{2x^2-11\}$
$= -4x^2+22$

42. $-\{-[-(x-2a)-b]-(a-x)\} = -\{-[-x+2a-b]-a+x\}$
$= -\{x-2a+b-a+x\}$
$= -\{-3a+b+2x\}$
$= 3a-b-2x$

43. $5V^2-(6-(2V^2+3)) = 5V^2-(6-2V^2-3)$
$= 5V^2-(-2V^2+3)$
$= 5V^2+2V^2-3$
$= 7V^2-3$

44. $-2F+2((2F-1)-5) = -2F+2(2F-1-5)$
$= -2F+2(2F-6)$
$= -2F+4F-12$
$= 2F-12$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

45. $-(3t-(7+2t-(5t-6))) = -(3t-(7+2t-5t+6))$
$= -(3t-(-3t+13))$
$= -(3t+3t-13)$
$= -(6t-13)$
$= -6t+13$

46. $a^2-2(x-5-(7-2(a^2-2x)-3x)) = a^2-2(x-5-(7-2a^2+4x-3x))$
$= a^2-2(x-5-(7-2a^2+x))$
$= a^2-2(x-5-7+2a^2-x)$
$= a^2-2(2a^2-12)$
$= a^2-4a^2+24$
$= -3a^2+24$

47. $-4[4R-2.5(Z-2R)-1.5(2R-Z)] = -4[4R-2.5Z+5R-3R+1.5Z]$
$= -4[6R-Z]$
$= -24R+4Z$

48. $-3\{2.1e-1.3[-f-2(e-5f)]\} = -3\{2.1e-1.3[-f-2e+10f]\}$
$= -3\{2.1e-1.3[-2e+9f]\}$
$= -3\{2.1e+2.6e-11.7f\}$
$= -3\{4.7e-11.7f\}$
$= -14.1e+35.1f$

49. $3D-(D-d) = 3D-D+d = 2D+d$

50. $i_1-(2-3i_2)+i_2 = i_1-2+3i_2+i_2 = i_1+4i_2-2$

51. $B+\frac{4}{3}\alpha+2\ B-\frac{2}{3}\alpha\ -\ B+\frac{4}{3}\alpha\ -\ B-\frac{2}{3}\alpha = B+\frac{4}{3}\alpha+2B-\frac{4}{3}\alpha\ -\ B+\frac{4}{3}\alpha-B+\frac{2}{3}\alpha$
$= [3B]-\frac{6}{3}\alpha$
$= 3B-2\alpha$

52. Distance $= 30\text{ km/h}\times(t-1)\text{h}+40\text{ km/h}\times(t+2)\text{ h}$
$= 30(t-1)\text{ km}+40(t+2)\text{ km}$
$= (30t-30+40t+80)\text{ km}$
$= (70t+50)\text{ km}$

53. Memory $= x(4\text{ terabytes})+(x+25)(8\text{ terabytes})$
$= (4x+8x+200)\text{ terabytes}$
$= (12x+200)\text{ terabytes}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

54. $\begin{aligned}\text{Difference} &= 2[(2n+1)(\$30)-(n-2)(\$20)]\\ &= \$\ 2[60n+30-20n+40]\\ &= \$\ 2[40n+70]\\ &= \$\ (80n+140)\end{aligned}$

55. **(a)** $\begin{aligned}(2x^2-y+2a)+(3y-x^2-b) &= 2x^2-y+2a+3y-x^2-b\\ &= x^2+2y+2a-b\end{aligned}$

(b) $\begin{aligned}(2x^2-y+2a)-(3y-x^2-b) &= 2x^2-y+2a-3y+x^2+b\\ &= 3x^2-4y+2a+b\end{aligned}$

56. $\begin{aligned}(3a^2+b-c^3)+(2c^3-2b-a^2)-(4c^3-4b+3) &= 3a^2+b-c^3+2c^3-2b-a^2-4c^3+4b-3\\ &= 2a^2+3b-3c^3-3\end{aligned}$

57. The final y should be added and the final 3 should be subtracted. The correct final answer is $-2x-2y+2$.

58. The final occurrence of $2c$ should be added rather than subtracted, resulting in the final answer of $7a-6b-2c$.

59. $\begin{aligned}|a-b| &= |-(-a+b)|\\ &= |-(b-a)|\\ &= |-1\times(b-a)|\\ &= |-1|\times|(b-a)|\\ &= 1\times|b-a|\\ &= |b-a|\end{aligned}$

60. $(a-b)-c = a-b-c$

However, $a-(b-c) = a-b+c$

Since they are not equivalent, subtraction is not associative.

For example, $(10-5)-2 = 5-2 = 3$ is not the same as $10-(5-2) = 10-3 = 7$.

1.8 Multiplication of Algebraic Expressions

1. $\begin{aligned}2s^3(-st^4)^3(4s^2t) &= 2s^3(-1)^3s^3t^{12}(4s^2t)\\ &= -2s^6t^{12}(4s^2t)\\ &= -8s^8t^{13}\end{aligned}$

2. $\begin{aligned}-2ax(3ax^2-4yz) &= (-2ax)(3ax^2)-(-2ax)(4yz)\\ &= (-6a^2x^3)-(-8axyz)\\ &= -6a^2x^3+8axyz\end{aligned}$

3. $\begin{aligned}(x-2)(x-3) &= x(x)+x(-3)+(-2)(x)+(-2)(-3)\\ &= x^2-3x-2x+6\\ &= x^2-5x+6\end{aligned}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

4. $(2a-b)^2 = (2a-b)(2a-b)$
$= (2a)(2a)+(2a)(-b)+(2a)(-b)+(-b)(-b)$
$= 4a^2 - 2ab - 2ab + b^2$
$= 4a^2 - 4ab + b^2$

5. $(a^2)(ax) = a^3x$

6. $(2xy)(x^2y^3) = 2x^3y^4$

7. $-a^2c^2(a^2cx^3) = -a^4c^3x^3$

8. $(-2cs^2)(-4cs)^2 = (-2cs^2)(-4cs)(-4cs)$
$= (-2cs^2)(16c^2s^2)$
$= -32c^3s^4$

9. $(2ax^2)^2(-2ax) = (2ax^2)(2ax^2)(-2ax)$
$= (4a^2x^4)(-2ax)$
$= -8a^3x^5$

10. $(6pq^3)(3pq^2)^2 = (6pq^3)(3pq^2)(3pq^2)$
$= (6pq^3)(9p^2q^4)$
$= 54p^3q^7$

11. $i^2(Ri+2ri) = (i^2)(Ri)+(i^2)(2ri)$
$= i^3R + 2i^3r$

12. $2x(-p-q) = (2x)(-p)-(2x)(q)$
$= -2px - 2qx$

13. $-3s(s^2-5t) = (-3s)(s^2)+(-3s)(-5t)$
$= -3s^3 + 15st$

14. $-3b(2b^2-b) = (-3b)(2b^2)+(-3b)(-b)$
$= -6b^3 + 3b^2$

15. $5m(m^2n+3mn) = (5m)(m^2n)+(5m)(3mn)$
$= 5m^3n + 15m^2n$

16. $a^2bc(2ac-3b^2c) = (a^2bc)(2ac)+(a^2bc)(-3b^2c)$
$= 2a^3bc^2 - 3a^2b^3c^2$

17. $3M(-M-N+2) = (3M)(-M)+(3M)(-N)+(3M)(2)$
$= -3M^2 - 3MN + 6M$

18. $-4c^2(-9gc-2c+g^2) = (-4c^2)(-9cg)+(-4c^2)(-2c)+(-4c^2)(g^2)$
$= 36c^3g + 8c^3 - 4c^2g^2$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

19. $xy(tx^2)(x+y^3) = tx^3y(x+y^3)$
$= (tx^3y)(x) + (tx^3y)(y^3)$
$= tx^4y + tx^3y^4$

20. $-2(-3st^3)(3s-4t) = 6st^3(3s-4t)$
$= (6st^3)(3s) + (6st^3)(-4t)$
$= 18s^2t^3 - 24st^4$

21. $(x-3)(x+5) = (x)(x) + (x)(5) + (-3)(x) + (-3)(5)$
$= x^2 + 5x - 3x - 15$
$= x^2 + 2x - 15$

22. $(a+7)(a+1) = (a)(a) + (a)(1) + (7)(a) + (7)(1)$
$= a^2 + a + 7a + 7$
$= a^2 + 8a + 7$

23. $(x+5)(2x-1) = (x)(2x) + (x)(-1) + (5)(2x) + (5)(-1)$
$= 2x^2 - x + 10x - 5$
$= 2x^2 + 9x - 5$

24. $(4t_1 + t_2)(2t_1 - 3t_2) = (4t_1)(2t_1) + (4t_1)(-3t_2) + (t_2)(2t_1) + (t_2)(-3t_2)$
$= 8t_1^2 - 12t_1t_2 + 2t_1t_2 - 3t_2^2$
$= 8t_1^2 - 10t_1t_2 - 3t_2^2$

25. $(y+8)(y-8) = (y)(y) + (y)(-8) + (8)(y) + (8)(-8)$
$= y^2 - 8y + 8y - 64$
$= y^2 - 64$

26. $(z-4)(z+4) = (z)(z) + (z)(4) + (-4)(z) + (-4)(4)$
$= z^2 + 4z - 4z - 16$
$= z^2 - 16$

27. $(2a-b)(-2b+3a) = (2a)(-2b) + (2a)(3a) + (-b)(-2b) + (-b)(3a)$
$= -4ab + 6a^2 + 2b^2 - 3ab$
$= 6a^2 - 7ab + 2b^2$

28. $(-3+4w^2)(3w^2-1) = (-3)(3w^2) + (-3)(-1) + (4w^2)(3w^2) + (4w)(-1)$
$= -9w^2 + 3 + 12w^4 - 4w^2$
$= 12w^4 - 13w^2 + 3$

29. $(2s+7t)(3s-5t) = (2s)(3s) + (2s)(-5t) + (7t)(3s) + (7t)(-5t)$
$= 6s^2 - 10st + 21st - 35t^2$
$= 6s^2 + 11st - 35t^2$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

30. $(5p-2q)(p+8q) = (5p)(p)+(5p)(8q)+(-2q)(p)+(-2q)(8q)$
$= 5p^2+40pq-2pq-16q^2$
$= 5p^2+38pq-16q^2$

31. $(x^2-1)(2x+5) = (x^2)(2x)+(x^2)(5)+(-1)(2x)+(-1)(5)$
$= 2x^3+5x^2-2x-5$

32. $(3y^2+2)(2y-9) = (3y^2)(2y)+(3y^2)(-9)+(2)(2y)+(-9)(2)$
$= 6y^3-27y^2+4y-18$

33. $(x-2y-4)(x-2y+4)$
$= (x)(x)+(x)(-2y)+(x)(4)+(-2y)(x)+(-2y)(-2y)+(-2y)(4)+(-4)(x)+(-4)(-2y)+(-4)(4)$
$= x^2-2xy+4x-2xy+4y^2-8y-4x+8y-16$
$= x^2+4y^2-4xy-16$

34. $(2a+3b+1)(2a+3b-1)$
$= (2a)(2a)+(2a)(3b)+(2a)(-1)+(3b)(2a)+(3b)(3b)+(3b)(-1)+(1)(2a)+(1)(3b)+(1)(-1)$
$= 4a^2+6ab-2a+6ab+9b^2-3b+2a+3b-1$
$= 4a^2+9b^2+12ab-1$

35. $2(a+1)(a-9) = 2[(a)(a)+(a)(-9)+(1)(a)+(-9)(1)]$
$= 2[a^2-9a+a-9]$
$= 2[a^2-8a-9]$
$= 2a^2-16a-18$

36. $-5(y-3)(y+6) = -5[(y)(y)+(y)(6)+(-3)(y)+(-3)(6)]$
$= -5[y^2+6y-3y-18]$
$= -5[y^2+3y-18]$
$= -5y^2-15y+90$

37. $-3(3-2T)(3T+2) = -3[(3)(3T)+(3)(2)+(-2T)(3T)+(-2T)(2)]$
$= -3[-6T^2+9T-4T+6]$
$= -3[-6T^2+5T+6]$
$= 18T^2-15T-18$

38. $2n(-n+5)(6n+5) = 2n[(-n)(6n)+(-n)(5)+(5)(6n)+(5)(5)]$
$= 2n[-6n^2-5n+30n+25]$
$= 2n[-6n^2+25n+25]$
$= -12n^3+50n^2+50n$

39. $2L(L+1)(4-L) = 2L[(L)(4)+(L)(-L)+(1)(4)+(1)(-L)]$
$= 2L[-L^2+4L-L+4]$
$= 2L[-L^2+3L+4]$
$= -2L^3+6L^2+8L$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

40. $ax(x+4)(7-x^2) = ax[(x)(7)+(x)(-x^2)+(4)(7)+(4)(-x^2)]$
$= ax[-x^3-4x^2+7x+28]$
$= -ax^4-4ax^3+7ax^2+28ax$

41. $(3x-7)^2 = (3x-7)(3x-7)$
$= (3x)(3x)+(3x)(-7)+(-7)(3x)+(-7)(-7)$
$= 9x^2-21x-21x+49$
$= 9x^2-42x+49$

42. $(x-3y)^2 = (x-3y)(x-3y)$
$= (x)(x)+(x)(-3y)+(-3y)(x)+(-3y)(-3y)$
$= x^2-3xy-3xy+9y^2$
$= x^2-6xy+9y^2$

43. $(x_1+3x_2)^2 = (x_1+3x_2)(x_1+3x_2) = (x_1)(x_1)+(x_1)(3x_2)+(3x_2)(x_1)+(3x_2)(3x_2)$
$= x_1^2+3x_1x_2+3x_1x_2+9x_2^2$
$= x_1^2+6x_1x_2+9x_2^2$

44. $(-7m-1)^2 = (-7m-1)(-7m-1)$
$= (-7m)(-7m)+(-7m)(-1)+(-1)(-7m)+(-1)(-1)$
$= 49m^2+7m+7m+1$
$= 49m^2+14m+1$

45. $(xyz-2)^2 = (xyz-2)(xyz-2)$
$= (xyz)(xyz)+(xyz)(-2)+(-2)(xyz)+(-2)(-2)$
$= x^2y^2z^2-2xyz-2xyz+4$
$= x^2y^2z^2-4xyz+4$

46. $(-6x^2+b)^2 = (-6x^2+b)(-6x^2+b)$
$= (-6x^2)(-6x^2)+(-6x^2)(b)+(b)(-6x^2)+(b)(b)$
$= 36x^4-6bx^2-6bx^2+b^2$
$= 36x^4-12bx^2+b^2$

47. $2(x+8)^2 = 2[(x+8)(x+8)]$
$= 2[(x)(x)+(x)(8)+(8)(x)+(8)(8)]$
$= 2[x^2+8x+8x+64]$
$= 2[x^2+16x+64]$
$= 2x^2+32x+128$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

48. $3(3R-4)^2 = 3[(3R-4)(3R-4)]$
$= 3[(3R)(3R)+(3R)(-4)+(-4)(3R)+(-4)(-4)]$
$= 3[9R^2-12R-12R+16]$
$= 3[9R^2-24R+16]$
$= 27R^2-72R+48$

49. $(2+x)(3-x)(x-1) = [(6-2x+3x-x^2)](x-1)$
$= (x-1)[-x^2+x+6]$
$= (x)(-x^2)+(x)(x)+(6)(x)+(-1)(-x^2)+(-1)(x)+(-1)(6)$
$= -x^3+x^2+6x+x^2-x-6$
$= -x^3+2x^2+5x-6$

50. $(-c^2+3x)^3 = (-c^2+3x)(-c^2+3x)(-c^2+3x)$
$= [(3x)(3x)-3c^2x-3c^2x+c^4)](-c^2+3x)$
$= (-c^2+3x)[9x^2-6c^2x+c^4]$
$= (-c^2)(9x^2)+(-c^2)(-6c^2x)+(-c^2)(+c^4)+(3x)(9x^2)+(3x)(-6c^2x)+(3x)(+c^4)$
$= -9c^2x^2+6c^4x-c^6+27x^3-18c^2x^2+3c^4x$
$= -c^6+9c^4x-27c^2x^2+27x^3$

51. $3T(T+2)(2T-1) = 3T[(T)(2T)+(T)(-1)+(2)(2T)+(2)(-1)]$
$= 3T[2T^2-T+4T-2]$
$= 3T[2T^2-T+4T-2]$
$= 3T[2T^2+3T-2]$
$= 6T^3+9T^2-6T$

52. $[(x-2)^2(x+2)]^2$
$= [(x-2)(x-2)(x+2)][(x-2)(x-2)(x+2)]$
$= [(x-2)[(x)(x)+(-2)(x)+(2)(x)+(-2)(2)]][(x-2)[(x)(x)+(-2)(x)+(2)(x)+(-2)(2)]]$
$= [(x-2)[x^2-2x+2x-4]][(x-2)[x^2-2x+2x-4]]$
$= [(x-2)[x^2-4]][(x-2)[x^2-4]]$
$= [(x)(x^2)+(-4)(x)+(-2)(x^2)+(-2)(-4)][(x)(x^2)+(-4)(x)+(-2)(x^2)+(-2)(-4)]$
$= [x^3-2x^2-4x+8][x^3-2x^2-4x+8]$
$= (x^3)(x^3)+(x^3)(-2x^2)+(x^3)(-4x)+(x^3)(8)+(-2x^2)(x^3)+(-2x^2)(-2x^2)+(-2x^2)(-4x)+(-2x^2)(8)$
$+(-4x)(x^3)+(-4x)(-2x^2)+(-4x)(-4x)+(-4x)(8)+(8)(x^3)+(8)(-2x^2)+(8)(-4x)+(8)(8)$
$= x^6-2x^5-4x^4+8x^3-2x^5+4x^4+8x^3-16x^2-4x^4+8x^3+16x^2-32x+8x^3-16x^2-32x+64$
$= x^6-4x^5-4x^4+32x^3-16x^2-64x+64$

53. **(a)** $(x+y)^2 = (3+4)^2 = 7^2 = 49$
$x^2+y^2 = 3^2+4^2 = 9+16 = 25$
$(x+y)^2 \neq x^2+y^2$
$49 \neq 25$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

(b) $(x-y)^2 = (3-4)^2 = (-1)^2 = 1$

$x^2 - y^2 = 3^2 - 4^2 = 9 - 16 = -7$

$(x-y)^2 \neq x^2 - y^2$

$1 \neq -7$

54. One can write $(x+3)^5 = (x+3)(x+3)(x+3)(x+3)(x+3)$ and then perform the multiplications using the rightmost pair of terms at each step.

55.
$$\begin{aligned}(x+y)^3 &= (x+y)(x+y)(x+y)\\ &= (x+y)[(x)(x)+(x)(y)+(y)(x)+(y)(y)]\\ &= (x+y)[x^2+xy+xy+y^2]\\ &= (x+y)[x^2+2xy+y^2]\\ &= (x)(x^2)+(x)(2xy)+(x)(y^2)+(y)(x^2)+(y)(2xy)+(y)(y^2)\\ &= x^3+2x^2y+y^2x+x^2y+2y^2x+y^3\\ &= x^3+3x^2y+3y^2x+y^3\end{aligned}$$

This differs from x^3+y^3 by the presence of $3x^2y+3y^2x$ in the center.

56.
$$\begin{aligned}&(x+y)(x^2-xy+y^2)\\ &= (x)(x^2)+(x)(-xy)+(x)(y^2)+(y)(x^2)+(y)(-xy)+(y)(y^2)\\ &= x^3-x^2y+y^2x+x^2y-y^2x+y^3\\ &= x^3+y^3\end{aligned}$$

57.
$$\begin{aligned}P(1+0.01r)^2 &= P(1+0.01r)(1+0.01r)\\ &= P[(1)(1)+(1)(0.01r)+(0.01r)(1)+(0.01r)(0.01r)]\\ &= P[1+0.01r+0.01r+0.0001r^2]\\ &= 0.0001r^2P+0.02rP+P\end{aligned}$$

58.
$$\begin{aligned}1000(1+0.0025r)^2 &= 1000(1+0.0025r)(1+0.0025r)\\ &= 1000[(1)(1)+(1)(0.0025r)+(0.0025r)(1)+(0.0025r)(0.0025r)]\\ &= 1000[1+0.0025r+0.0025r+0.0000625r^2]\\ &= 1000+2.5r+0.0625r^2\end{aligned}$$

59. The room will be $5+w+5 = w+10$ feet wide and $5+2w+5 = 2w+10$ feet long. Its area is

$$\begin{aligned}(w+10)(2w+10) &= (w)(2w)+(w)(10)+(10)(2w)+(10)(10)\\ &= 2w^2+10w+20w+100\\ &= 2w^2+30w+100\end{aligned}$$

60.
$$\begin{aligned}R &= xp\\ &= x(30-0.01x)\\ &= 30x-0.01x^2\end{aligned}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

61. $(2R-X)^2-(R^2+X^2)=(2R-X)(2R-X)-(R^2+X^2)$
$=[(2R)(2R)+(2R)(-X)+(2R)(-X)+(-X)(-X)]-(R^2+X^2)$
$=4R^2-2RX-2RX+X^2-R^2-X^2$
$=3R^2-4RX$

62. $(2T^3+3)(T^2-T-3)=(2T^3)(T^2)+(2T^3)(-T)+(2T^3)(-3)+(3)(T^2)+(3)(-T)+(3)(-3)$
$=2T^5-2T^4-6T^3+3T^2-3T-9$

63. Number of switches for n elements $=n^2$
Number of switches for $n+100$ elements $=(n+100)^2$
$=(n+100)(n+100)$
$=(n)(n)+(n)(100)+(100)(n)+(100)(100)$
$=n^2+100n+100n+10000$
$=n^2+200n+10000$

64. $(T^2-100)(T-10)(T+10)=(T^2-100)\ \ T^2+10T-10T-100$
$=(T^2-100)\ \ T^2-100$
$=T^4-100T^2-100T^2+10\ 000$
$=T^4-200T^2+10\ 000$

65. $(R_1+R_2)^2-2R_2(R_1+R_2)=\left[(R_1+R_2)(R_1+R_2)\right]-2R_2(R_1+R_2)$
$=\left[(R_1)(R_1)+(R_1)(R_2)+(R_2)(R_1)+(R_2)(R_2)\right]-2R_1R_2-2R_2{}^2$
$=\ R_1{}^2+R_1R_2+R_1R_2+R_2{}^2\ -2R_1R_2-2R_2{}^2$
$=R_1{}^2-R_2{}^2$

66. $27x^2-24(x-6)^2-(x-12)^3$
$=27x^2-24(x-6)(x-6)-(x-12)(x-12)(x-12)$
$=27x^2-24\ \ x^2-6x-6x+36\ \ -(x-12)\ \ x^2-12x-12x+144$
$=27x^2-24\ \ x^2-12x+36\ \ -(x-12)\ \ x^2-24x+144$
$=27x^2-24x^2+288x-864-\ \ (x)(x^2)+(x)(-24x)+(x)(144)+(-12)(x^2)+(-12)(-24x)+(-12)(144)$
$=3x^2+288x-864-x^3+24x^2-144x+12x^2-288x+1728$
$=-x^3+39x^2-144x+864$

1.9 Division of Algebraic Expressions

1. $\dfrac{-6a^2xy^2}{-2a^2xy^5}=\left(\dfrac{-6}{-2}\right)\dfrac{a^{2-2}x^{1-1}}{y^{5-2}}=\dfrac{3}{y^3}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

2. $$\frac{4x^3y-8x^3y^2+2x^2y}{2xy^2}=\frac{4x^3y}{2xy^2}-\frac{8x^3y^2}{2xy^2}+\frac{2x^2y}{2xy^2}$$
$$=\frac{2x^{3-1}}{y^{2-1}}-4x^{3-1}y^{2-2}+\frac{x^{2-1}}{y^{2-1}}$$
$$=\frac{2x^2}{y}-4x^2+\frac{x}{y}$$

3.
$$\begin{array}{r} 3x-2 \\ 2x-1\overline{)\,6x^2-7x+2} \\ \underline{6x^2-3x} \\ -4x+2 \\ \underline{-4x+2} \\ 0 \end{array}$$

4.
$$\begin{array}{r} 2x-1 \\ 4x^2-1\overline{)\,8x^3-4x^2+0x+3} \\ \underline{8x^3\qquad -2x} \\ -4x^2+2x+3 \\ \underline{-4x^2\qquad 1} \\ 2x+2 \end{array}$$
$$\frac{8x^3-4x^2+3}{4x^2-1}=2x-1+\frac{2x+2}{4x^2-1}$$

5. $$\frac{8x^3y^2}{-2xy}=-4x^{3-1}y^{2-1}=-4x^2y$$

6. $$\frac{-18b^7c^3}{bc^2}=-18b^{7-1}c^{3-2}=-18b^6c$$

7. $$\frac{-16r^3t^5}{-4r^5t}=\frac{4t^{5-1}}{r^{5-3}}=\frac{4t^4}{r^2}$$

8. $$\frac{51mn^5}{17m^2n^2}=\frac{3n^{5-2}}{m^{2-1}}=\frac{3n^3}{m}$$

9. $$\frac{(15x^2y)(2xz)}{10xy}=\frac{30x^3yz}{10xy}=3x^{3-1}y^{1-1}z=3x^2z$$

10. $$\frac{(5sT)(8s^2T^3)}{10s^3T^2}=\frac{40s^3T^4}{10s^3T^2}=4s^{3-3}T^{4-2}=4T^2$$

11. $$\frac{(4a^3)(2x)^2}{(4ax)^2}=\frac{4a^3(4x^2)}{16a^2x^2}=1a^{3-2}x^{2-2}=a$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

12. $\dfrac{12a^2b}{(3ab^2)^2} = \dfrac{12a^2b}{9a^2b^4} = \dfrac{4a^{2-2}}{3b^{4-1}} = \dfrac{4}{3b^3}$

13. $\dfrac{3a^2x+6xy}{3x} = \dfrac{3a^2x}{3x} + \dfrac{6xy}{3x} = \dfrac{3a^2x^{1-1}}{3} + \dfrac{6x^{1-1}y}{3} = a^2 + 2y$

14. $\dfrac{2m^2n-6mn}{-2m} = \dfrac{2m^2n}{-2m} - \dfrac{6mn}{-2m} = -m^{2-1}n + 3m^{1-1}n = -mn + 3n$

15. $\dfrac{3rst-6r^2st^2}{3rs} = \dfrac{3rst}{3rs} - \dfrac{6r^2st^2}{3rs} = r^{1-1}s^{1-1}t - 2r^{2-1}s^{1-1}t^2 = -2rt^2 + t$

16. $\dfrac{-5a^2n-10an^2}{5an} = \dfrac{-5a^2n}{5an} - \dfrac{10an^2}{5an} = -a^{2-1}n^{1-1} - 2a^{1-1}n^{2-1} = -a - 2n$

17.
$$\begin{aligned}\frac{4pq^3+8p^2q^2-16pq^5}{4pq^2} &= \frac{4pq^3}{4pq^2} + \frac{8p^2q^2}{4pq^2} - \frac{16pq^5}{4pq^2} \\ &= p^{1-1}q^{3-2} + 2p^{2-1}q^{2-2} - 4p^{1-1}q^{5-2} \\ &= -4q^3 + 2p + q\end{aligned}$$

18.
$$\begin{aligned}\frac{a^2x_1x_2^2 + ax_1^3 - ax_1}{ax_1} &= \frac{a^2x_1x_2^2}{ax_1} + \frac{ax_1^3}{ax_1} - \frac{ax_1}{ax_1} \\ &= a^{2-1}x_1^{1-1}x_2^2 + a^{1-1}x_1^{3-1} - a^{1-1}x_1^{1-1} \\ &= ax_2^2 + x_1^2 - 1\end{aligned}$$

19.
$$\begin{aligned}\frac{2\pi fL - \pi fR^2}{\pi fR} &= \frac{2\pi fL}{\pi fR} - \frac{\pi fR^2}{\pi fR} \\ &= \frac{2f^{1-1}L}{R} - f^{1-1}R^{2-1} \\ &= \frac{2L}{R} - R\end{aligned}$$

20.
$$\begin{aligned}\frac{9(aB)^4 - 6aB^4}{-3aB^3} &= \frac{9(aB)^4}{-3aB^3} - \frac{6aB^4}{-3aB^3} \\ &= -\frac{9a^4B^4}{3aB^3} + \frac{6aB^4}{3aB^3} \\ &= -3a^{4-1}B^{4-3} + 2a^{1-1}B^{4-3} \\ &= -3a^3B + 2B\end{aligned}$$

21.
$$\begin{aligned}\frac{-7a^2b+14ab^2-21a^3}{14a^2b^2} &= -\frac{7a^2b}{14a^2b^2} + \frac{14ab^2}{14a^2b^2} - \frac{21a^3}{14a^2b^2} \\ &= -\frac{a^{2-2}}{2b^{2-1}} + \frac{b^{2-2}}{a^{2-1}} - \frac{3}{2}a^{3-2}b^{-2} \\ &= -\frac{1}{2b} + \frac{1}{a} - \frac{3a}{2b^2}\end{aligned}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

22. $$\frac{2x^{n+2}+4ax^n}{2x^n}=\frac{2x^{n+2}}{2x^n}+\frac{4ax^n}{2x^n}$$
$$=x^{n-n+2}+2ax^{n-n}$$
$$=x^2+2a$$

23. $$\frac{6y^{2n}-4ay^{n+1}}{2y^n}=\frac{6y^{2n}}{2y^n}-\frac{4ay^{n+1}}{2y^n}$$
$$=3y^{2n-n}-2ay^{n-n+1}$$
$$=3y^n-2ay$$

24. $$\frac{3a(F+T)b^2-(F+T)}{a(F+T)}=\frac{3a(F+T)b^2}{a(F+T)}-\frac{(F+T)}{a(F+T)}$$
$$=\frac{3a^{1-1}\cancel{(F+T)}b^2}{\cancel{(F+T)}}-\frac{\cancel{(F+T)}}{a\cancel{(F+T)}}$$
$$=3b^2-\frac{1}{a}$$

25.
$$\begin{array}{r} x+5 \\ x+4\overline{)\,x^2+9x+20} \\ \underline{x^2+4x} \\ 5x+20 \\ \underline{5x+20} \\ 0 \end{array}$$
$$\frac{x^2+9x+20}{x+4}=x+5$$

26.
$$\begin{array}{r} x+9 \\ x-2\overline{)\,x^2+7x-18} \\ \underline{x^2-2x} \\ 9x-18 \\ \underline{9x-18} \\ 0 \end{array}$$
$$\frac{x^2+7x-18}{x-2}=x+9$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

27.
$$x+3\overline{)2x^2+7x+3}$$
$$\begin{array}{r} 2x+1 \\ x+3\overline{)2x^2+7x+3} \\ \underline{2x^2+6x} \\ x+3 \\ \underline{x+3} \\ 0 \end{array}$$
$$\frac{2x^2+7x+3}{x+3}=2x+1$$

28.
$$\begin{array}{r} 3t-4 \\ t-1\overline{)3t^2-7t+4} \\ \underline{3t^2-3t} \\ -4t+4 \\ \underline{-4t+4} \\ 0 \end{array}$$
$$\frac{3t^2-7t+4}{t-1}=3t-4$$

29.
$$\begin{array}{r} x-1 \\ x-2\overline{)x^2-3x+2} \\ \underline{x^2-2x} \\ -x+2 \\ \underline{-x+2} \\ 0 \end{array}$$
$$\frac{x^2-3x+2}{x-2}=x-1$$

30.
$$\begin{array}{r} 2x-7 \\ x+1\overline{)2x^2-5x-7} \\ \underline{2x^2+2x} \\ -7x-7 \\ \underline{-7x-7} \\ 0 \end{array}$$
$$\frac{2x^2-5x-7}{x+1}=2x-7$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

31.
$$\begin{array}{r} 4x^2 - x - 1 \\ 2x-3\overline{)\,8x^3 - 14x^2 + x + 0} \\ \underline{8x^3 - 12x^2} \\ -2x^2 + x \\ \underline{-2x^2 + 3x} \\ -2x + 0 \\ \underline{-2x + 3} \\ -3 \end{array}$$

$$\frac{8x^3 - 14x^2 + x}{2x-3} = 4x^2 - x - 1 - \frac{3}{2x-3}$$

32.
$$\begin{array}{r} 3y + 2 \\ 2y+1\overline{)\,6y^2 + 7y + 6} \\ \underline{6y^2 + 3y} \\ 4y + 6 \\ \underline{4y + 2} \\ 4 \end{array}$$

$$\frac{6y^2 + 7y + 6}{2y+1} = 3y + 2 + \frac{4}{2y+1}$$

33.
$$\begin{array}{r} Z - 2 \\ 4Z+3\overline{)\,4Z^2 - 5Z - 7} \\ \underline{4Z^2 + 3Z} \\ -8Z - 7 \\ \underline{-8Z - 6} \\ -1 \end{array}$$

$$\frac{4Z^2 - 5Z - 7}{4Z+3} = Z - 2 - \frac{1}{4Z+3}$$

34.
$$\begin{array}{r} 2x + 1 \\ 3x-4\overline{)\,6x^2 - 5x - 9} \\ \underline{6x^2 - 8x} \\ 3x - 9 \\ \underline{3x - 4} \\ -5 \end{array}$$

$$\frac{6x^2 - 5x - 9}{3x-4} = 2x + 1 - \frac{5}{3x-4}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

35.
$$\begin{array}{r} x^2+x-6 \\ x+2\overline{)\,x^3+3x^2-4x-12} \\ \underline{x^3+2x^2} \\ x^2-4x \\ \underline{x^2+2x} \\ -6x-12 \\ \underline{-6x-12} \\ 0 \end{array}$$

$$\frac{x^3+3x^2-4x-12}{x+2}=x^2+x-6$$

36.
$$\begin{array}{r} x^2+7x+9 \\ 3x-2\overline{)\,3x^3+19x^2+13x-20} \\ \underline{3x^3-2x^2} \\ 21x^2+13x \\ \underline{21x^2-14x} \\ 27x-20 \\ \underline{27x-18} \\ -2 \end{array}$$

$$\frac{3x^3+19x^2+13x-20}{3x-2}=x^2+7x+9-\frac{2}{3x-2}$$

37.
$$\begin{array}{r} 2a^2+8 \\ a^2-2\overline{)\,2a^4+0a^3+4a^2+0a-16} \\ \underline{2a^4-4a^2} \\ 8a^2-16 \\ \underline{8a^2-16} \\ 0 \end{array}$$

$$\frac{2a^4+4a^2-16}{a^2-2}=2a^2+8$$

38.
$$\begin{array}{r} 2T+1 \\ 3T^2-T+2\overline{)\,6T^3+T^2+0T+2} \\ \underline{6T^3-2T^2+4\mathrm{T}} \\ 3T^2-4T+2 \\ \underline{3T^2-T+2} \\ -3T \end{array}$$

$$\frac{6T^3+T^2+2}{3T^2-T+2}=2T+1-\frac{3T}{3T^2-T+2}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

39.
$$y+3\overline{)y^3+0y^2+0y+27}$$
$$\begin{array}{r} y^2-3y+9 \\ \hline y^3+3y^2 \\ \quad -3y^2+0y \\ \quad \underline{-3y^2-9y} \\ \qquad\quad 9y+27 \\ \qquad\quad \underline{9y+27} \\ \qquad\qquad 0 \end{array}$$

$$\frac{y^3+27}{y+3}=y^2-3y+9$$

40.
$$D-1\overline{)D^3+0D^2+0D-1}$$
$$\begin{array}{r} D^2+D+1 \\ \hline D^3-1D^2 \\ \quad D^2+0D \\ \quad \underline{D^2-1D} \\ \qquad\quad D-1 \\ \qquad\quad \underline{D-1} \\ \qquad\qquad 0 \end{array}$$

$$\frac{D^3-1}{D-1}=D^2+D+1$$

41.
$$x-y\overline{)x^2-2xy+y^2}$$
$$\begin{array}{r} x-y \\ \hline x^2-xy \\ \quad -xy+y^2 \\ \quad \underline{-xy+y^2} \\ \qquad\quad 0 \end{array}$$

$$\frac{x^2-2xy+y^2}{x-y}=x-y$$

42.
$$r-3R\overline{)3r^2-5rR+2R^2}$$
$$\begin{array}{r} 3r+4R \\ \hline 3r^2-9rR \\ \quad 4rR+2R^2 \\ \quad \underline{4rR-12R^2} \\ \qquad\quad 14R^2 \end{array}$$

$$\frac{3r^2-5rR+2R^2}{r-3R}=3r+4R+\frac{14R^2}{r-3R}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

43.
$$\begin{array}{r} t-2 \\ t^2+2t+4\overline{)\,t^3+0t^2+0t-8} \\ \underline{t^3+2t^2+4t} \\ -2t^2-4t-8 \\ \underline{-2t^2-4t-8} \\ 0 \end{array}$$

$$\frac{t^3-8}{t^2+2t+4}=t-2$$

44.
$$\begin{array}{r} a^2+2ab+2b^2 \\ a^2-2ab+2b^2\overline{)\,a^4+0a^3b+0a^2b^2+0ab^3+b^4} \\ \underline{a^4-2a^3b+2a^2b^2} \\ 2a^3b-2a^2b^2+0ab^3 \\ \underline{2a^3b-4a^2b^2+4ab^3} \\ 2a^2b^2-4ab^3+b^4 \\ \underline{2a^2b^2-4ab^3+4b^4} \\ -3b^4 \end{array}$$

$$\frac{a^4+b^4}{a^2-2ab+2b^2}=a^2+2ab+2b^2-\frac{3b^4}{a^2-2ab+2b^2}$$

45. We know that $2x+1$ multiplied by $x+c$ will give us $2x^2-9x-5$, so $2x^2-9x-5$ divided by $2x+1$ will give us $x+c$:

$$\begin{array}{r} x-5 \\ 2x+1\overline{)\,2x^2-9x-5} \\ \underline{2x^2+x} \\ -10x-5 \\ \underline{-10x-5} \\ 0 \end{array}$$

$$x+c=x-5$$
$$c=-5$$

46.
$$\begin{array}{r} 2x-3 \\ 3x+4\overline{)\,6x^2-x+k} \\ \underline{6x^2+8x} \\ -9x+k \\ \underline{-9x-12} \\ 0 \end{array}$$

$$k-(-12)=0$$
$$k+12=0$$
$$k=-12$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

47.
$$\begin{array}{r} x^3 - x^2 + x - 1 \\ x+1\overline{)\,x^4 + 0x^3 + 0x^2 + 0x + 1} \\ \underline{x^4 + x^3} \\ -x^3 + 0x^2 \\ \underline{-x^3 - x^2} \\ x^2 + 0x \\ \underline{x^2 + x} \\ -x + 1 \\ \underline{-x - 1} \\ 2 \end{array}$$

$$\frac{x^4+1}{x+1} = x^3 - x^2 + x - 1 + \frac{2}{x+1} \neq x^3$$

48.
$$\begin{array}{r} x^2 - xy + y^2 \\ x+y\overline{)\,x^3 + 0x^2y + 0y^2x + 0x + y^3} \\ \underline{x^3 + x^2\text{y}} \\ -x^2\text{y} + 0y^2x \\ \underline{-x^2y - y^2x} \\ y^2x + y^3 \\ \underline{y^2x + y^3} \\ 0 \end{array}$$

$$\frac{x^3+y^3}{x+y} = x^2 - xy + y \neq x^2 + y^2$$

49.
$$\begin{aligned} V_1\left(1 + \frac{T_2 - T_1}{T_1}\right) &= V_1\left(1 + \frac{T_2}{T_1} - \frac{T_1}{T_1}\right) \\ &= V_1\left(1 + \frac{T_2}{T_1} - 1\right) \\ &= V_1\frac{T_2}{T_1} \\ &= \frac{V_1T_2}{T_1} \end{aligned}$$

50.
$$\begin{array}{r} 3x + 2 \\ 2x+5\overline{)\,6x^2 + 19x + 10} \\ \underline{6x^2 + 15x} \\ 4x + 10 \\ \underline{4x + 10} \\ 0 \end{array}$$

The width is $3x + 2$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

51. $$\frac{8A^5+4A^3\mu^2E^2-A\mu^4E^4}{8A^4}=\frac{8A^5}{8A^4}+\frac{4A^3\mu^2E^2}{8A^4}-\frac{A\mu^4E^4}{8A^4}$$
$$=A^{5-4}+\frac{\mu^2E^2}{2A^{4-3}}-\frac{\mu^4E^4}{8A^{4-1}}$$
$$=A+\frac{\mu^2E^2}{2A}-\frac{\mu^4E^4}{8A^3}$$

52. $$\frac{6R_1+6R_2+R_1R_2}{6R_1R_2}=\frac{6R_1}{6R_1R_2}+\frac{6R_2}{6R_1R_2}+\frac{R_1R_2}{6R_1R_2}$$
$$=\frac{\cancel{R_1}}{\cancel{R_1}R_2}+\frac{\cancel{R_2}}{R_1\cancel{R_2}}+\frac{\cancel{R_1R_2}}{6\cancel{R_1R_2}}$$
$$=\frac{1}{R_1}+\frac{1}{R_2}+\frac{1}{6}$$

53. $$\frac{GMm[(R+r)-(R-r)]}{2rR}=\frac{GMm[R+r-R+r]}{2rR}$$
$$=\frac{GMm[2r]}{2rR}$$
$$=\frac{GMm\cancel{[2r]}}{\cancel{2r}R}$$
$$=\frac{GMm}{R}$$

54.
$$\begin{array}{r} 3T^2-2T-4 \\ T-2\overline{)\,3T^3-8T^2+0T+8} \\ \underline{3T^3-6T^2} \\ -2T^2+0T \\ \underline{-2T^2+4T} \\ -4T+8 \\ \underline{-4T+8} \\ 0 \end{array}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

55. $\left(\dfrac{s^2-2s-2}{s^4+4}\right)^{-1} = \dfrac{s^4+4}{s^2-2s-2}$

$$
\begin{array}{r}
s^2+2s+6 \\
s^2-2s-2\overline{)\,s^4+0s^3+0s^2+0s+4} \\
\underline{s^4-2s^3-2s^2} \\
2s^3+2s^2+0s \\
\underline{2s^3-4s^2-4s} \\
6s^2+4s+4 \\
\underline{6s^2-12s-12} \\
16s+16
\end{array}
$$

$$\frac{s^4+4}{s^2-2s-2} = s^2+2s+6+\frac{16s+16}{s^2-2s-2}$$

56.
$$
\begin{array}{r}
t^2-3t+5 \\
2t+100\overline{)\,2t^3+94t^2-290t+500} \\
\underline{2t^3+100t^2} \\
-6t^2-290t \\
\underline{-6t^2-300t} \\
10t+500 \\
\underline{10t+500} \\
0
\end{array}
$$

1.10 Solving Equations

1. **(a)** $x-3=-12$

$x-3+3=-12+3$

$x=-9$

(b) $x+3=-12$

$x+3-3=-12-3$

$x=-15$

(c) $\dfrac{x}{3}=-12$

$3\left(\dfrac{x}{3}\right)=3(-12)$

$x=-36$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

(d) $3x = -12$

$$\frac{3x}{3} = \frac{-12}{3}$$

$$x = -4$$

2. $7 - 2t = 9$

$7 - 7 - 2t = 9 - 7$

$-2t = 2$

$$\frac{-2t}{-2} = \frac{2}{-2}$$

$t = -1$

Check:

$7 - 2t = 9$

$7 - 2(-1) = 9$

$7 - (-2) = 9$

$9 = 9$

3. $x - 7 = 3x - (8 - 6x)$

$x - 7 = 3x - 8 + 6x$

$x - 7 = 9x - 8$

$-8x = -1$

$$x = \frac{1}{8}$$

4. $$\frac{L}{3.80} = \frac{7}{4}$$

$$3.80\ \frac{L}{3.80} = 3.80\ \frac{7}{4}$$

$L = 6.65$ m

5. $x - 2 = 7$

$x = 7 + 2$

$x = 9$

6. $x - 4 = -1$

$x = -1 + 4$

$x = 3$

7. $x + 5 = 4$

$x = 4 - 5$

$x = -1$

8. $s + 6 = -3$

$s = -3 - 6$

$s = -9$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

9. $\frac{t}{2} = -5$

$t = 2(-5)$

$t = -10$

10. $\frac{x}{-4} = 2$

$x = -4(2)$

$x = -8$

11. $\frac{y-8}{3} = 4$

$y - 8 = 4(3)$

$y = 12 + 8$

$y = 20$

12. $\frac{7-r}{6} = 3$

$7 - r = 6(3)$

$-r = 18 - 7$

$-r = 11$

$r = -11$

13. $4E = -20$

$E = \frac{-20}{4}$

$E = -5$

14. $2x = 12$

$x = \frac{12}{2}$

$x = 6$

15. $5t + 9 = -1$

$5t = -1 - 9$

$t = \frac{-10}{5}$

$t = -2$

16. $5D - 2 = 13$

$5D = 13 + 2$

$D = \frac{15}{5}$

$D = 3$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

17. $5-2y=-3$

$-2y=-3-5$

$y=\dfrac{-8}{-2}$

$y=4$

18. $-5t+8=18$

$-5t=18-8$

$t=\dfrac{10}{-5}$

$t=-2$

19. $3x+7=x$

$x-3x=7$

$-2x=7$

$x=-\dfrac{7}{2}$

20. $6+4L=5-3L$

$4L+3L=5-6$

$7L=-1$

$L=-\dfrac{1}{7}$

21. $2(3q+4)=5q$

$6q+8=5q$

$6q-5q=-8$

$q=-8$

22. $3(4-n)=-n$

$-n=12-3n$

$-n+3n=12$

$2n=12$

$n=\dfrac{12}{2}$

$n=6$

23. $-(r-4)=6+2r$

$-r+4=6+2r$

$-r-2r=2$

$-3r=2$

$r=-\dfrac{2}{3}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

24. $-(x+2)+5=5x$

$5x=5-x-2$

$5x+x=3$

$6x=3$

$x=\frac{3}{6}=\frac{1}{2}$

25. $8(y-5)=-2y$

$8y-40=-2y$

$8y+2y=40$

$10y=40$

$y=\frac{40}{10}$

$y=4$

26. $4(7-F)=-7$

$28-4F=-7$

$-4F=-7-28$

$F=\frac{-35}{-4}=\frac{35}{4}$

27. $0.1x-0.5(x-2)=2$

$x-5(x-2)=2(10)$

$x-5x+10=20$

$-4x=20-10$

$x=\frac{10}{-4}=-\frac{5}{2}$

28. $1.5x-0.3(x-4)=6$

$15x-3(x-4)=6(10)$

$15x-3x+12=60$

$12x=60-12$

$x=\frac{48}{12}$

$x=4$

29. $-4-3(1-2p)=-7+2p$

$-4-3+6p=-7+2p$

$-7+6p-2p=-7$

$4p=-7+7$

$p=\frac{0}{4}$

$p=0$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

30. $3-6(2-3t)=t-5$

$3-12+18t=t-5$

$-9+18t-t=-5$

$17t=4$

$t=\frac{4}{17}$

31. $\frac{4x-2(x-4)}{3}=8$

$4x-2x+8=3(8)$

$2x=24-8$

$x=\frac{16}{2}$

$x=8$

32. $2x=\frac{-5(7-3x)+2}{4}$

$4(2x)=-35+15x+2$

$8x-15x=-33$

$-7x=-33$

$x=\frac{-33}{-7}=\frac{33}{7}$

33. $|x|-9=2$

$|x|=2+9=11$

$x=11$ or $x=-11$

34. $2-|x|=4$

$-|x|=4-2$

$|x|=\frac{2}{-1}$

$|x|=-2$

There is no real solution for x.

35. $|2x-3|=5$

$2x-3=5$ or $2x-3=-5$

$2x-3=5$ $\quad 2x-3=-5$

$2x=5+3$ $\quad 2x=-5+3$

$2x=8$ $\quad 2x=-2$

$x=\frac{8}{2}$ $\quad x=\frac{-2}{2}$

$x=4$ or $x=-1$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

36. $|7-x|=1$

$7-x=1$ or $7-x=-1$

$-x=1-7 \qquad -x=-1-7$

$-x=-6 \qquad -x=-8$

$x=6 \qquad$ or $\qquad x=8$

37. $5.8-0.3(x-6.0)=0.5x$

$0.5x=5.8-0.3x+1.8$

$0.5x+0.3x=7.6$

$0.8x=7.6$

$x=\dfrac{7.6}{0.8}$

$x=9.5$

38. $1.9t=0.5(4.0-t)-0.8$

$1.9t=2.0-0.5t-0.8$

$1.9t+0.5t=1.2$

$2.4t=1.2$

$t=\dfrac{1.2}{2.4}$

$t=0.50$

39. $-0.24(C-0.50)=0.63$

$-0.24C+0.12=0.63$

$-0.24C=0.63-0.12$

$-0.24C=0.51$

$C=\dfrac{0.51}{-0.24}$

$C=-2.125$

$C=-2.1$

40. $27.5(5.17-1.44x)=73.4$

$142.175-39.6x=73.4$

$-39.6x=73.4-142.175$

$-39.6x=-68.775$

$x=\dfrac{-68.775}{-39.6}$

$x=1.736742424$

$x=1.74$

41. $\dfrac{x}{2.0}=\dfrac{17}{6.0}$

$x=2.0\ \dfrac{17}{6.0}$

$x=5.6666666...$

$x=5.7$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

42. $\frac{3.0}{7.0} = \frac{R}{42}$

$R = 42\ \frac{3.0}{7.0}$

$R = 18$

43. $\frac{165}{223} = \frac{13V}{15}$

$\frac{15}{13}\ \frac{165}{223} = \frac{15}{13}\ \frac{13V}{15}$

$V = \frac{2475}{2899}$

$V = 0.85374267$

$V = 0.85$

44. $\frac{276x}{17.0} = \frac{1360}{46.4}$

$276x = 17\ \frac{1360}{46.4}$

$x = \frac{498.2758621}{276}$

$x = 1.805347326$

$x = 1.81$

45. **(a)** $2x + 3 = 3 + 2x$

$2x + 3 = 2x + 3$

Is an identity, since it is true for all values of x.

(b) $2x - 3 = 3 - 2x$

$4x = 6$

$x = \frac{6}{4} = \frac{3}{2}$

Is conditional as x has one answer only.


46. There are no values of a that result in a conditional equation. If $a = 0$, then the identity $2x = 2x$ results. If $a \neq 0$, then a contradiction results.


47. $x - 7 = 3x - (6x - 8)$


$0 = 3x - 6x + 8 - x + 7$

$0 = -4x + 15$

$x = 3.75$







Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual


48. $0.0595 - 0.525i - 8.85(i + 0.0316) = 0$


$0.595 - 0.525i - 8.85i - 0.27966 = 0$

$-9.375i + 0.31534 = 0$

$i = 0.033636266$

$i = 0.0336$





49. $0.03x + 0.06(2000 - x) = 96$

$0.03x + 120 - 0.06x = 96$

$-0.03x = 96 - 120$

$-0.03x = -24$

$$x = \frac{-24}{-0.03}$$

$x = \$800$

50. $15(5.5 + v) = 24(5.5 - v)$

$82.5 + 15v = 132 - 24v$

$15v + 24v = 132 - 82.5$

$39v = 49.5$

$$v = \frac{49.5}{39}$$

$v = 1.269230769$ km/h

$v = 1.3$ km/h

51. $$1.1 = \frac{(T - 76)}{40}$$

$40(1.1) = T - 76$

$44 = T - 76$

$T = 44 + 76$

$T = 120$ °C

52. $1.12V - 0.67(10.5 - V) = 0$

$1.12V - 7.035 + 0.67V = 0$

$1.79V - 7.035 = 0$

$1.79V = 7.035$

$$V = \frac{7.035}{1.79}$$

$V = 3.930167598$ V

$V = 3.9$ V

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

53. $0.14n + 0.06(2000 - n) = 0.09(2000)$

$$0.14n + 120 - 0.06n = 180$$
$$0.14n - 0.06n = 180 - 120$$
$$0.08n = 60$$
$$n = \frac{60}{0.08}$$
$$n = 750 \text{ L}$$

54. $$210(3x) = 55.3x + 38.5(8.25 - 3x)$$
$$630x = 55.3x + 317.625 - 115.5x$$
$$630x - 55.3x + 115.5x = 317.625$$
$$690.2x = 317.625$$
$$x = \frac{317.625}{690.2}$$
$$x = 0.4601927 \text{ m}$$
$$x = 0.460 \text{ m}$$

55. $$\frac{x}{350 \text{ mi}} = \frac{30 \text{ kW} \cdot \text{h}}{107 \text{ mi}}$$
$$x = 350 \text{ mi} \times \frac{30 \text{ kW} \cdot \text{h}}{107 \text{ mi}}$$
$$x = 98 \text{ kW} \cdot \text{h}$$

56. $$\frac{20 \text{ min}}{250 \text{ cal}} = \frac{x}{400 \text{ cal}}$$
$$x = 400 \text{ cal} \ \frac{20 \text{ min}}{250 \text{ cal}}$$
$$x = 32 \text{ min}$$

1.11 Formulas and Literal Equations

1. $$v = v_0 + at$$
$$v - v_0 = at$$
$$a = \frac{v - v_0}{t}$$

2. $$W = \frac{L(wL + 2P)}{8}$$
$$8W = L(wL + 2P)$$
$$8W = wL^2 + 2LP$$
$$wL^2 = 8W - 2LP$$
$$w = \frac{8W - 2LP}{L^2}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

3.
$$V = V_0[1 + b(T - T_0)]$$
$$V = V_0[1 + bT - bT_0]$$
$$V = V_0 + bTV_0 - bT_0V_0$$
$$bT_0V_0 = V_0 + bTV_0 - V$$
$$T_0 = \frac{V_0 + bTV_0 - V}{bV_0}$$

4.
$$V = V_0 + V_0\beta T$$
$$V - V_0 = V_0\beta T$$
$$\beta = \frac{V - V_0}{V_0T}$$

5.
$$E = IR$$
$$R = \frac{E}{I}$$

6.
$$pV = nRT$$
$$T = \frac{pV}{nR}$$

7.
$$rL = g_2 - g_1$$
$$g_1 + rL = g_2$$
$$g_1 = g_2 - rL$$

8.
$$W = S_dT - Q$$
$$Q + W = S_dT$$
$$Q = S_dT - W$$

9.
$$B = \frac{nTWL}{12}$$
$$12B = nTWL$$
$$n = \frac{12B}{TWL}$$

10.
$$P = 2\pi Tf$$
$$T = \frac{P}{2\pi f}$$

11.
$$p = p_a + dgh$$
$$p - p_a = dgh$$
$$h = \frac{p - p_a}{dg}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

12. $2Q = 2I + A + S$

$2I = 2Q - A - S$

$I = \dfrac{2Q - A - S}{2}$

13. $F_c = \dfrac{mv^2}{r}$

$rF_c = mv^2$

$r = \dfrac{mv^2}{F_c}$

14. $P = \dfrac{4F}{\pi D^2}$

$P\pi D^2 = 4F$

$F = \dfrac{P\pi D^2}{4}$

15. $S_T = \dfrac{A}{5T} + 0.05d$

$S_T - 0.05d = \dfrac{A}{5T}$

$A = 5T\left(S_T - 0.05d\right)$

16. $u = -\dfrac{eL}{2m}$

$eL = -2mu$

$L = -\dfrac{2mu}{e}$

17. $ct^2 = 0.3t - ac$

$ac + ct^2 = 0.3t$

$ac = 0.3t - ct^2$

$a = \dfrac{-ct^2 + 0.3t}{c}$

18. $2p + dv^2 = 2d(C - W)$

$2p + dv^2 = 2Cd - 2dW$

$2Cd = dv^2 + 2p + 2dW$

$C = \dfrac{dv^2 + 2dW + 2p}{2d}$

19. $T = \dfrac{c + d}{v}$

$c + d = Tv$

$d = Tv - c$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

20.
$$B = \frac{\mu_0 I}{2\pi R}$$
$$BR = \frac{\mu_0 I}{2\pi}$$
$$R = \frac{\mu_0 I}{2\pi B}$$

21.
$$\frac{K_1}{K_2} = \frac{m_1 + m_2}{m_1}$$
$$K_2(m_1 + m_2) = K_1 m_1$$
$$K_2 m_1 + K_2 m_2 = K_1 m_1$$
$$K_2 m_2 = K_1 m_1 - K_2 m_1$$
$$m_2 = \frac{K_1 m_1 - K_2 m_1}{K_2}$$

22.
$$f = \frac{F}{d - F}$$
$$f(d - F) = F$$
$$fd - fF = F$$
$$fd = F + fF$$
$$d = \frac{F + fF}{f}$$

23.
$$a = \frac{2mg}{M + 2m}$$
$$a(M + 2m) = 2gm$$
$$aM + 2am = 2gm$$
$$aM = 2gm - 2am$$
$$M = \frac{2gm - 2am}{a}$$

24.
$$v = \frac{V(m + M)}{m}$$
$$mv = mV + MV$$
$$MV = mv - mV$$
$$M = \frac{mv - mV}{V}$$

25.
$$C_0^2 = C_1^2(1 + 2V)$$
$$C_0^2 = C_1^2 + 2C_1^2 V$$
$$2C_1^2 V = C_0^2 - C_1^2$$
$$V = \frac{C_0^2 - C_1^2}{2C_1^2}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

26.
$$A_1 = A(M+1)$$
$$A_1 = AM + A$$
$$AM = A_1 - A$$
$$M = \frac{A_1 - A}{A}$$

27.
$$N = r(A - s)$$
$$N = Ar - rs$$
$$rs + N = Ar$$
$$rs = Ar - N$$
$$s = \frac{Ar - N}{r}$$

28.
$$T = 3(T_2 - T_1)$$
$$T = 3T_2 - 3T_1$$
$$3T_1 + T = 3T_2$$
$$3T_1 = 3T_2 - T$$
$$T_1 = \frac{3T_2 - T}{3}$$

29.
$$T_2 = T_1 - \frac{h}{100}$$
$$100T_2 = 100T_1 - h$$
$$h + 100T_2 = 100T_1$$
$$h = 100T_1 - 100T_2$$

30.
$$p_2 = p_1 + rp_1(1 - p_1)$$
$$p_2 - p_1 = rp_1(1 - p_1)$$
$$r = \frac{p_2 - p_1}{p_1(1 - p_1)}$$

31.
$$Q_1 = P(Q_2 - Q_1)$$
$$Q_1 = PQ_2 - PQ_1$$
$$PQ_2 = Q_1 + PQ_1$$
$$Q_2 = \frac{Q_1 + PQ_1}{P}$$

32.
$$p - p_a = dg(y_2 - y_1)$$
$$y_2 - y_1 = \frac{p - p_a}{dg}$$
$$-y_1 = \frac{p - p_a}{dg} - y_2$$
$$y_1 = y_2 - \frac{p - p_a}{dg}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

33.
$$N = N_1T - N_2(1-T)$$
$$N_1T = N + N_2(1-T)$$
$$N_1 = \frac{N + N_2 - N_2T}{T}$$

34.
$$t_a = t_c + (1-h)t_m$$
$$t_a = t_c + t_m - ht_m$$
$$t_a + ht_m = t_c + t_m$$
$$ht_m = t_c + t_m - t_a$$
$$h = \frac{t_c + t_m - t_a}{t_m}$$

35.
$$L = \pi(r_1 + r_2) + 2x_1 + 2x_2$$
$$L = \pi r_1 + \pi r_2 + 2x_1 + 2x_2$$
$$\pi r_1 = L - \pi r_2 - 2x_1 - 2x_2$$
$$r_1 = \frac{L - \pi r_2 - 2x_1 - 2x_2}{\pi}$$

36.
$$I = \frac{VR_2 + VR_1(1+\mu)}{R_1R_2}$$
$$IR_1R_2 = VR_2 + VR_1 + VR_1\mu$$
$$VR_1\mu = IR_1R_2 - VR_2 + VR_1$$
$$\mu = \frac{IR_1R_2 - VR_2 + VR_1}{VR_1}$$

37.
$$P = \frac{V_1(V_2 - V_1)}{gJ}$$
$$gJP = V_1V_2 - V_1^2$$
$$V_1V_2 = V_1^2 + gJP$$
$$V_2 = \frac{V_1^2 + gJP}{V_1}$$

38.
$$W = T(S_1 - S_2) - Q$$
$$W + Q = TS_1 - TS_2$$
$$TS_2 = TS_1 - W - Q$$
$$S_2 = \frac{TS_1 - W - Q}{T}$$

39.
$$C = \frac{2eAk_1k_2}{d(k_1 + k_2)}$$
$$Cd(k_1 + k_2) = 2eAk_1k_2$$
$$e = \frac{Cd(k_1 + k_2)}{2Ak_1k_2}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

40.
$$d = \frac{3LPx^2 - Px^3}{6EI}$$
$$6dEI = 3LPx^2 - Px^3$$
$$3LPx^2 = 6dEI + Px^3$$
$$L = \frac{6dEI + Px^3}{3Px^2}$$

41.
$$V = C\left(1 - \frac{n}{N}\right)$$
$$V = C - \frac{Cn}{N}$$
$$V + \frac{Cn}{N} = C$$
$$\frac{Cn}{N} = C - V$$
$$Cn = CN - NV$$
$$n = \frac{CN - NV}{C}$$

42.
$$\frac{p}{P} = \frac{AI}{B + AI}$$
$$p(B + AI) = AIP$$
$$pB + AIp = AIP$$
$$pB = AIP - AIp$$
$$B = \frac{AIP - AIp}{p}$$

43.
$$p(C - n) + n = A$$
$$pC - pn + n = A$$
$$(-p + 1)n = A - pC$$
$$(1 - p)n = A - pC$$
$$n = \frac{A - pC}{1 - p}$$
$$n = \frac{13.0\text{ L} - 0.25(15.0\text{ L})}{1 - 0.25}$$
$$T_1 = \frac{13.0\text{ L} - 3.75\text{ L}}{0.75}$$
$$T_1 = \frac{9.25\text{ L}}{0.75}$$
$$T_1 = 12.333333\text{ L}$$
$$T_1 = 12\text{ L}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

44. $P_t = P_c(1+0.500m^2)\dfrac{\pi}{2}$

$$P_c = \frac{P_t}{1+0.500m^2}$$

$$P_c = \frac{680\text{ W}}{1+0.500(0.925)^2}$$

$$P_c = \frac{680\text{ W}}{1+0.500(0.855625)}$$

$$P_c = \frac{680\text{ W}}{1+0.4278125}$$

$$P_c = \frac{680\text{ W}}{1.4278125}$$

$$P_c = 476.253009\text{ W}$$

$$P_c = 476\text{ W}$$

45. $$F = \frac{9}{5}C + 32$$

$$90.2 = \frac{9}{5}C + 32$$

$$\frac{5}{9}(90.2 - 32) = C$$

$$C = \frac{5}{9} \times 58.2$$

$$C = 32.3^\circ\text{C}$$

46. $$V = \frac{1}{2}L(B+b)$$

$$2V = BL + bL$$

$$bL = 2V - BL$$

$$b = \frac{2V - BL}{L}$$

$$b = \frac{2(38.6\text{ ft}^3) - (2.63\text{ ft}^2)(16.1\text{ ft})}{16.1\text{ ft}}$$

$$b = \frac{77.2\text{ ft}^3 - 42.343\text{ ft}^3}{16.1\text{ ft}}$$

$$b = \frac{34.857\text{ ft}^3}{16.1\text{ ft}}$$

$$b = 2.16503106\text{ ft}^2$$

$$b = 2.16\text{ ft}^2$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

47. $V_1 = \dfrac{VR_1}{R_1 + R_2}$

$V_1(R_1 + R_2) = VR_1$

$R_1 + R_2 = \dfrac{VR_1}{V_1}$

$R_2 = \dfrac{VR_1}{V_1} - R_1$

$R_2 = \dfrac{(12.0\text{ V})(3.56\ \Omega)}{6.30\text{ V}} - (3.56\ \Omega)$

$R_2 = 6.780952381\ \Omega - 3.56\ \Omega$

$R_2 = 3.220952381\ \Omega$

$R_2 = 3.22\ \Omega$

48. $\eta = \dfrac{1}{q + p(1-q)}$

$\eta\left[q + p(1-q)\right] = 1$

$\eta q + \eta p(1-q) = 1$

$\eta p(1-q) = 1 - \eta q$

$p = \dfrac{1 - \eta q}{\eta(1-q)}$

$p = \dfrac{1 - (0.66)(0.83)}{0.66(1-0.83)}$

$p = \dfrac{1 - 0.5478}{0.66(0.17)}$

$p = \dfrac{0.4522}{0.1122}$

$p = 4.03030303$

$p = 4$ processors

49. $d = v_2 t_2 + v_1 t_1$

$d = v_2(4\text{ h}) + v_1(t + 2\text{ h})$

$tv_1 + v_1(2\text{ h}) = d - v_2(4\text{ h})$

$tv_1 = d - v_2(4\text{ h}) - v_1(2\text{ h})$

$t = \dfrac{d - v_2(4\text{ h}) - v_1(2\text{ h})}{v_1}$

50. $C = x + 15y$

$15y = C - x$

$y = \dfrac{C - x}{15}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

1.12 Applied Word Problems

1. Let x = the number of 25 W lights.
Let $31-x$ = the number of 40 W lights.

$$25x+40(31-x)=1000$$
$$25x+1240-40x=1000$$
$$-15x=1000-1240$$
$$-15x=-240$$
$$x=16$$

There are 16 of the 25 W lights and$(31-16)=15$ of the 40 W lights.
Check:

$$25\cdot 16+40(31-16)=1000$$
$$400+40(15)=1000$$
$$400+600=1000$$
$$1000=1000$$

2. Let$x$ = the number of slides with 5 mg.
Let $x-3$ = the number of slides with 6 mg.

$$(5\text{ mg})x=(6\text{ mg})(x-3)$$
$$(5\text{ mg})x=(6\text{ mg})x-18\text{ mg}$$
$$-x=-18$$
$$x=18\text{ slides}$$

There are 18 slides with 5 mg and $(18-3)=15$ slides with 6 mg.
Check:

$$5\text{ mg}(18)=6\text{ mg}(15)$$
$$90\text{ mg}=90\text{ mg}$$

3. Let t = the time for the shuttle to reach the satellite.

$$(29\ 500\text{ km/h})t=6000\text{ km}+(27\ 100\text{ km/h})t$$
$$(2400\text{ km/h})t=6000\text{ km}$$
$$t=\frac{6000\text{ km}}{2400\text{ km/h}}$$
$$t=2.500\text{ h}$$

It will take the shuttle 2.500 h to reach the satellite.
Check:

$$(29500\text{ km/h})(2.500\text{ h})=6000\text{ km}+(27100\text{ km/h})(2.500\text{ h})$$
$$73750\text{ km}=6000\text{ km}+67750\text{ km}$$
$$73750\text{ km}=73750\text{ km}$$

4. Let x = the number of litres of 50% methanol blend that must be added.

$$0.0600(7250\text{ L})+0.500(x)=0.100(7250\text{ L}+x)$$
$$435\text{ L}+0.500(x)=725\text{ L}+0.100x$$
$$0.400(x)=290\text{ L}$$
$$x=\frac{290\text{ L}}{0.400}$$
$$x=725\text{ L}$$

725 L of the 50% methanol blend must be added.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

Check:

$$0.0600(7250\text{ L}) + 0.500(725\text{ L}) = 0.100(7250\text{ L} + 725\text{ L})$$
$$435\text{ L} + 362.5\text{ L} = 0.1(7975\text{ L})$$
$$797.5\text{ L} = 797.5\text{ L}$$

5. Let x = the cost of the car 6 years ago.
Let x + \$5000 = the cost of the car model today.

$$x + (x + \$5000) = \$49\ 000$$
$$2x = \$44\ 000$$
$$x = \frac{\$44\ 000}{2}$$
$$x = \$22\ 000$$

The cost of the car 6 years ago was \$22 000, and the cost of the today's model is (\$22 000 + 5000) = \$27 000.
Check:

$$\$22\ 000 + (\$22\ 000 + \$5000) = \$49\ 000$$
$$\$22\ 000 + \$27\ 000 = \$49\ 000$$
$$\$49\ 000 = \$49\ 000$$

6. Let x = the flow from the first stream in m^3/s.
Let $x - 1700\ ft^3/s$ = the flow from the second streamin m^3/s.

$$x + (x - 1700\ \text{ft}^3/\text{s}) = \frac{1.98\times10^7\ \text{ft}^3}{3600\ \text{s}}$$
$$2x - 1700\ \text{ft}^3/\text{s} = 5500\ \text{ft}^3/\text{s}$$
$$2x = 7200\ \text{ft}^3/\text{s}$$
$$x = \frac{7200\ \text{ft}^3/\text{s}}{2}$$
$$x = 3600\ \text{ft}^3/\text{s}$$

The first stream flows $3600\ ft^3/s$ and the second stream flows $3600\ ft^3/s - 1700\ ft^3/s = 1900\ ft^3/s$.
Check:

$$3600\ \text{ft}^3/\text{s} + (3600\ \text{ft}^3/\text{s} - 1700\ \text{ft}^3/\text{s}) = \frac{1.98\times10^7\ \text{ft}^3}{3600\ \text{s}}$$
$$7200\ \text{ft}^3/\text{s} - 1700\ \text{ft}^3/\text{s} = 5500\ \text{ft}^3/\text{s}$$
$$5500\ \text{ft}^3/\text{s} = 5500\ \text{ft}^3/\text{s}$$

7. Let x = the number of cars recycled the first year.
Let x+500000 = the number of cars recycled the second year.

$$x + (x + 500000\text{ cars}) = 6900000\text{ cars}$$
$$2x + 500000\text{ cars} = 6900000\text{ cars}$$
$$2x = 6400000\text{ cars}$$
$$x = \frac{6400000\text{ cars}}{2}$$
$$x = 3200000\text{ cars}$$

The first year,3.2×10^6 cars were recycled, and the second year (3200000 + 500000)= 3.7×10^6 cars were recycled.
Check:

$$3200000\text{ cars} + (3200000\text{ cars} + 500000\text{ cars}) = 6900000\text{ cars}$$
$$3200000\text{ cars} + 3700000\text{ cars} = 6900000\text{ cars}$$
$$6900000\text{ cars} = 6900000\text{ cars}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

8. Let x = the number of hits to the website on the first day.
Let $1/2\,x$ = the number of hits on the second day.

$$x + 1/2\,x = 495000 \text{ hits}$$
$$3/2\,x = 495000 \text{ hits}$$
$$x = \frac{495000 \text{ hits}}{3/2}$$
$$x = 330000 \text{ hits}$$

The first day there were 330000 hits, the second day there were 1/2(33000 hits= 165000 hits.
Check:

$$330000 \text{ hits} + 1/2(330000 \text{ hits}) = 495000 \text{ hits}$$
$$330000 \text{ hits} + 165000 \text{ hits} = 495000 \text{ hits}$$
$$495000 \text{ hits} = 495000 \text{ hits}$$

9. Let x = the number acres of land leased for \$200 per acre.
Let $140 - x$ = the number of acres of land leased for \$300 per acre.

$$\$200/\text{acre}\,x + \$300/\text{acre}(140 \text{ acre} - x) = \$37\,000$$
$$-\$100/\text{acre}\,(x) = -\$5\,000$$
$$x = \frac{-\$5000}{-\$100/\text{acre}}$$
$$x = 50 \text{ acres}$$

There are 50 acres leased at \$200 per acre and (140 acres – 50 acres) = 90 hectares leased for \$300 per hectare.
Check:

$$\$200/\text{acre}\,(50 \text{ acres}) + \$300/\text{acre}(140 \text{ acres} - 50 \text{ acres}) = \$37\,000$$
$$\$10\,000 + \$27\,000 = \$37\,000$$
$$\$37\,000 = \$37\,000$$

10. Let x = the first dose in mg.
Let x + 660 mg = the second dose in mg.

$$x + x + 660 \text{ mg} = 2000 \text{ mg}$$
$$2x = 1340 \text{ mg}$$
$$x = \frac{1340 \text{ mg}}{2}$$
$$x = 670 \text{ mg}$$

The first dose is 670 mg, and the second dose is (670 mg + 660 mg) = 1130 mg.
Check:

$$670 \text{ mg} + 670 \text{ mg} + 660 \text{ mg} = 2000 \text{ mg}$$
$$670 \text{ mg} + 1330 \text{ mg} = 2000 \text{ mg}$$
$$2000 \text{ mg} = 2000 \text{ mg}$$

11. Let x = the amount of pollutant after modification in ppm/h.

$$(5 \text{ h})x = (3 \text{ h})150 \text{ ppm/h}$$
$$x = \frac{450 \text{ ppm}}{5 \text{ h}}$$
$$x = 90 \text{ ppm/h}$$

The amount of pollutant after modification is 90 ppm/h. The device reduced emissions by
(150 ppm/h – 90 ppm/h) = 60 ppm/h.
Check:

$$(5 \text{ h})90 \text{ ppm/h} = (3 \text{ h})150 \text{ ppm/h}$$
$$450 \text{ ppm} = 450 \text{ ppm}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

12. Let $x - 13$ = the number of teeth that the first meshed spur has.
Let x = the number of teeth that the second meshed spur has.
Let $x + 15$ = the number of teeth that the third meshed spur has.

$$x - 13 \text{ teeth} + x + x + 15 \text{ teeth} = 107 \text{ teeth}$$
$$3x + 2 = 107 \text{ teeth}$$
$$3x = 105 \text{ teeth}$$
$$x = \frac{105 \text{ teeth}}{3}$$
$$x = 35 \text{ teeth}$$

The first spur has (35 – 13)= 22 teeth, the second spur has 35 teeth, and the third spur has (35 + 15) = 50 teeth.
Check:

$$35 \text{ teeth} - 13 \text{ teeth} + 35 \text{ teeth} + 35 \text{ teeth} + 15 \text{ teeth} = 107 \text{ teeth}$$
$$107 \text{ teeth} = 107 \text{ teeth}$$

13. Let x = amount paid per month for first six months.
Let $x + 10$ = amount paid per month for final four months.

$$(6 \text{ mo})x + (4 \text{ mo})(x + \$10 / \text{mo}) = \$890$$
$$(10 \text{ mo})x + \$40 = \$890$$
$$(10 \text{ mo})x = \$850$$
$$x = \frac{\$850}{10 \text{ mo}}$$
$$x = \$85 / \text{mo}$$

The bill was \$85/mo for the first six months and \$95/mo for the next four months.
Check:

$$(6 \text{ mo})\$85/\text{mo} + (4 \text{ mo})(\$85 / \text{mo} + \$10 / \text{mo}) = \$890$$
$$\$510 + (4 \text{ mo})(\$95 / \text{mo}) = \$890$$
$$\$510 + \$380 = \$890$$
$$\$890 = \$890$$

14. Let x = amount paid per month for first year.
Let $x + 30$ = amount paid per month for next two years.
Let $(x + 30)+20 = x + 50$= amount paid per month for final two years.

$$(12 \text{ mo})x + (24 \text{ mo})(x + \$30 / \text{mo}) + (24 \text{ mo})(x + \$50 / \text{mo}) = \$7320$$
$$(12 \text{ mo})x + (24 \text{ mo})x + \$720 + (24 \text{ mo})x + \$1200 = \$7320$$
$$(60 \text{ mo})x + \$1920 = \$7320$$
$$(60 \text{ mo})x = \$5400$$
$$x = \frac{\$5400}{60 \text{ mo}}$$
$$x = \$90 / \text{mo}$$

For the first year, the bill was \$90/mo, during years 2 and 3, the bill was \$120/mo, and during years 4 and 5, the bill was \$140/mo.
Check:

$$(12 \text{ mo})(\$90 / \text{mo}) + (24 \text{ mo})(\$90 / \text{mo} + \$30 / \text{mo}) + (24 \text{ mo})(\$90 / \text{mo} + \$50 / \text{mo}) = \$7320$$
$$\$1080 + (24 \text{ mo})(\$120 / \text{mo}) + (24 \text{ mo})(\$140 / \text{mo}) = \$7320$$
$$\$1080 + \$2880 + \$3360 = \$7320$$
$$\$7320 = \$7320$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

15. Let x = the first current in μA .
Let $2x$ = the second current in μA .
Let $x + 9.2\ \mu$A = the third current in μA

$$x + 2x + x + 9.2\ \mu\text{A} = 0\ \mu\text{A}$$
$$4x = -9.2\ \mu\text{A}$$
$$x = \frac{-9.2\ \mu\text{A}}{4}$$
$$x = -2.3\ \mu\text{A}$$

The first current is $-2.3\ \mu$A ,the second current is $2(-2.3\ \mu\text{A}) = -4.6\ \mu$A , and the third current is $(-2.3\ \mu\text{A} + 9.2\ \mu\text{A}) = 6.9\ \mu$A .
Check:

$$-2.3\ \mu\text{A} + 2(-2.3\ \mu\text{A}) + (-2.3)\ \mu\text{A} + 9.2\ \mu\text{A} = 0\ \mu\text{A}$$
$$-2.3\ \mu\text{A} - 4.6\ \mu\text{A} - 2.3\mu\text{A} + 9.2\ \mu\text{A} = 0\ \mu\text{A}$$
$$0\ \mu\text{A} = 0\ \mu\text{A}$$

16. Let x = the number of trucks in the first fleet.
Let $x + 5$ = the number of trucks in the second fleet.

$$(8\text{ h})x + (6\text{ h})(x + 5) = 198\text{ h}$$
$$(8\text{ h})x + (6\text{ h})x + 30\text{ h} = 198\text{ h}$$
$$(14\text{ h})x = 168\text{ h}$$
$$x = \frac{168\text{ h}}{14\text{ h}}$$
$$x = 12\text{ trucks}$$

There are 12 trucks in the first fleet and (12 trucks + 5 trucks) = 17 trucks in the second fleet.
Check:

$$(8\text{ h})(12) + (6\text{ h})(12 + 5) = 198\text{ h}$$
$$96\text{ h} + (6\text{ h})(17) = 198\text{ h}$$
$$96\text{ h} + 102\text{ h} = 198\text{ h}$$
$$198\text{ h} = 198\text{ h}$$

17. Let x = the length of the first pipeline in km.
Let $x + 2.6$ km = the length of the 3 other pipelines.

$$x + 3(x + 2.6\text{ km}) = 35.4\text{ km}$$
$$x + 3x + 7.8\text{ km} = 35.4\text{ km}$$
$$4x = 27.6\text{ km}$$
$$x = \frac{27.6\text{ km}}{4}$$
$$x = 6.9\text{ km}$$

The first pipeline is 6.9 km long, and the other three pipelines are each (6.9 km + 2.6 km) = 9.5 km long.
Check:

$$6.9\text{ km} + 3(6.9\text{ km} + 2.6\text{ km}) = 35.4\text{ km}$$
$$6.9\text{ km} + 3(9.5\text{ km}) = 35.4\text{ km}$$
$$6.9\text{ km} + 28.5\text{ km} = 35.4\text{ km}$$
$$35.4\text{ km} = 35.4\text{ km}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

18. Let x = the power of the first generator in MW.
Let 750 MW$-x$ = the power of the second generator in MW.

$$0.65x + 0.75(750 \text{ MW} - x) = 530 \text{ MW}$$
$$0.65x + 562.5 \text{ MW} - 0.75x = 530 \text{ MW}$$
$$-0.1x = -32.5 \text{ MW}$$
$$x = \frac{-32.5 \text{ MW}}{-0.1}$$
$$x = 325 \text{ MW}$$

The first generator produces 325 MW of power, and the second generator produces (750 MW – 325 MW) = 425 MW of power.
Check:

$$0.65(325 \text{ MW}) + 0.75(750 \text{ MW} - (325 \text{ MW})) = 530 \text{ MW}$$
$$211.25 \text{ MW} + 0.75(425 \text{ MW}) = 530 \text{ MW}$$
$$211.25 \text{ MW} + 318.75 \text{ MW} = 530 \text{ MW}$$
$$530 \text{ MW} = 530 \text{ MW}$$

19. Let x = the number of deluxe systems.
Let $2x$ = the number economy systems.
Let x+75= the number of econo-plus systems.

$$\$140x + \$40(2x) + \$80(x + 75) = \$42000$$
$$\$140x + \$80x + \$80x + \$6000 = \$42000$$
$$\$300x = \$36000$$
$$x = 120 \text{ systems}$$

There are 120 deluxe systems, 240 economy systems, and 195 econo-plus systems sold.
Check:

$$\$140(120) + \$40(240) + \$80(195) = \$42000$$
$$\$16800 + \$9600 + \$15600 = \$42000$$
$$\$42000 = \$42000$$

20. The amount of lottery winnings after taxes is \$20 000 ×(1−0.25) = \$15 000.
Let x = the amount of money invested at a 40% gain.
Let \$15 000− x = the amount of money invested at a 10% loss.

$$0.40x - 0.10(\$15\ 000 - x) = \$2000$$
$$0.40x - \$1500 + 0.10x = \$2000$$
$$0.50x = \$3500$$
$$x = \frac{\$3500}{0.50}$$
$$x = \$7000$$

The 40% gain investment had \$7000 invested, and the 10% loss investment had (\$15 000− \$7000) = \$8000 invested.
Check:

$$0.40(\$7000) - 0.10(\$15\ 000 - \$7000) = \$2000$$
$$\$2800 - \$1500 + 0.10(\$7000) = \$2000$$
$$\$2800 - \$1500 + \$700 = \$2000$$
$$\$2000 = \$2000$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

21. Let x = the amount of time in seconds between when the start of the trains pass each other to when the end of the trains pass each other.
The total distance the ends must travel in this time is 960 feet. We first convert mi/hr into ft/sec.

$$1 \text{ mi/hr} = \frac{5280 \text{ ft}}{3600 \text{ s}} = \frac{22 \text{ ft}}{15 \text{ s}} = \frac{22}{15} \text{ ft/s}$$

Therefore, train A travels at 60(22/15)=88 ft/s and train B travels at 40(22/15)=176/3 ft/s.

$$(88 \text{ ft/s})x + (176/3 \text{ ft/s})x = 960 \text{ ft}$$
$$(440/3 \text{ ft/s})x = 960 \text{ ft}$$
$$x = \frac{960 \text{ ft}}{440/3 \text{ ft/s}}$$
$$x = \frac{72}{11} \text{ s}$$

The trains completely pass each other in about 6.55 seconds.
Check:

$$(88 \text{ ft/s})\frac{72}{11} \text{ s} + (176/3 \text{ ft/s})\frac{72}{11} \text{ s} = 960 \text{ ft}$$
$$576 \text{ ft} + 384 \text{ ft} = 960 \text{ ft}$$
$$960 \text{ ft} = 960 \text{ ft}$$

22. Let x = the mortgage payment and $x/0.23$=the monthly income.

$$x/0.23 - x = \$3850$$
$$x\left(\frac{1}{0.23} - 1\right) = \$3850$$
$$x\left(\frac{1}{0.23} - \frac{0.23}{0.23}\right) = \$3850$$
$$x\left(\frac{0.77}{0.23}\right) = \$3850$$
$$x = \$3850 \cdot \frac{0.23}{0.77}$$
$$x = \$1150$$

The mortgage payment is \$1150 and the monthly income is \$5000.
Check:

$$\$1150/0.23 - \$1150 = \$3850$$
$$\$5000 - \$1150 = \$3850$$
$$\$3850 = \$3850$$

23. Let x = the amount of time the skier spends on the ski lift in minutes.
Let 24 minutes $-x$ = the amount of time the skier spends skiing down the hill in minutes.

$$(50 \text{ m/min})x = (150 \text{ m/min})(24 \text{ min} - x)$$
$$(50 \text{ m/min})x = 3600 \text{ m} - (150 \text{ m/min})x$$
$$(200 \text{ m/min})x = 3600 \text{ m}$$
$$x = \frac{3600 \text{ m}}{200 \text{ m/min}}$$
$$x = 18 \text{ min}$$

The length of the slope is 18 minutes × 50 m/minute = 900m.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

Check:

$$(50 \text{ m/min})18 \text{ min} = (150 \text{ m/min})(24 \text{ min} - 18 \text{ min})$$
$$900 \text{ m} = 3600 \text{ m} - (150 \text{ m/min})(18 \text{ min})$$
$$900 \text{ m} = 3600 \text{ m} - 2700 \text{ m}$$
$$900 \text{ m} = 900 \text{ m}$$

24. Let x = the speed of sound.
Let $x - 120$ mi/h = speed travelled for 1 h.
Let $x + 410$ mi/h = the speed travelled for 3 h.

$$1 \text{ h}(x - 120 \text{ mi/h}) + 3 \text{ h}(x + 410 \text{ mi/h}) = 3990 \text{ mi}$$
$$(1 \text{ h})x - (1 \text{ h})(120 \text{ mi/h}) + (3 \text{ h})x + (3 \text{ h})(410 \text{ mi/h}) = 3990 \text{ mi}$$
$$(1 \text{ h})x - 120 \text{ mi} + (3 \text{ h})x + 1230 \text{ mi} = 3990 \text{ mi}$$
$$(4 \text{ h})x = 2880 \text{ mi}$$
$$x = \frac{2880 \text{ mi}}{4 \text{ h}}$$
$$x = 720 \text{ mi/h}$$

The speed of sound is 720 mi/h.
Check:

$$1 \text{ h}(720 \text{ mi/h} - 120 \text{ mi/h}) + 3 \text{ h}(720 \text{ mi/h} + 410 \text{ mi/h}) = 3990 \text{ mi}$$
$$(1 \text{ h})(600 \text{ mi/h}) + (3 \text{ h})(1130 \text{ mi/h}) = 3990 \text{ mi}$$
$$600 \text{ mi} + 3390 \text{ mi} = 3990 \text{ mi}$$
$$3990 \text{ mi} = 3990 \text{ mi}$$

25. Let x = the speed the train leaving England in km/h.
Let $x + 8$ km/h = speed of the train leaving France in km/h.
The distance travelled by each train is speed ×time.

$$x\left(\frac{17 \text{ min}}{60 \text{ min/ h}}\right) + (x + 8 \text{ km/h})\left(\frac{17 \text{ min}}{60 \text{ min/ h}}\right) = 50 \text{ km}$$
$$(0.28333 \text{ h})x + (x + 8 \text{ km/h})(0.28333 \text{ h}) = 50 \text{ km}$$
$$(0.28333 \text{ h})x + (0.28333 \text{ h})x + 2.26667 \text{ km} = 50 \text{ km}$$
$$(0.56666 \text{ h})x = 47.73333 \text{ km}$$
$$x = \frac{47.73333 \text{ km}}{0.56666 \text{ h}}$$
$$x = 84.23529421 \text{ km/h}$$
$$x = 84.2 \text{ km/h}$$

The train leaving England was travelling at 84.2 km/h, and the train leaving France was travelling at (84.2 km/h + 8 km/h) = 92.2 km/h.
Check:

$$84.23529421 \text{ km/h}\left(\frac{17 \text{ min}}{60 \text{ min/ h}}\right) + (84.23529421 \text{ km/h} + 8 \text{ km/h})\left(\frac{17 \text{ min}}{60 \text{ min/ h}}\right) = 50 \text{ km}$$
$$23.86666 \text{ km} + (92.23529421 \text{ km/h})\left(\frac{17 \text{ min}}{60 \text{ min/ h}}\right) = 50 \text{ km}$$
$$23.86666 \text{ km} + 26.13333 \text{ km} = 50 \text{ km}$$
$$50 \text{ km} = 50 \text{ km}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

26. Let x = time left until the appointment.
Let x– 10.0 min = time taken to get to the appointment travelling at 60.0 mi/h.
Let x– 5.0 min = time taken to get to the appointment travelling at 45.0 mi/h.
The distance travelled by the executive in each scenario is the same. Distance = speed × time

$$60.0 \text{ mi/h}\left(x - \frac{10.0 \text{ min}}{60 \text{ min/h}}\right) = 45 \text{ mi/h}\left(x - \frac{5.0 \text{ min}}{60 \text{ min/h}}\right)$$

$$(60.0 \text{ mi/h})x - 60 \text{ mi/h}\left(\frac{10.0 \text{ min}}{60 \text{ min/h}}\right) = (45 \text{ mi/h})x - 45 \text{ mi/h}\left(\frac{5.0 \text{ min}}{60 \text{ min/h}}\right)$$

$$(60.0 \text{ mi/h})x - 10 \text{ mi} = (45.0 \text{ mi/h})x - 3.75 \text{ mi}$$

$$(15.0 \text{ mi/h})x = 6.25 \text{ mi}$$

$$x = \frac{6.25 \text{ mi}}{15.0 \text{ mi/h}}$$

$$x = 0.416666667 \text{ h}$$

$$x = 0.416666667 \text{ h x } 60 \text{ min/h}$$

$$x = 25 \text{ min}$$

There is 25 minutes left until the executive's appointment.
Check:

$$60.0 \text{ mi/h}\left(0.41667 \text{ h} - \frac{10.0 \text{ min}}{60 \text{ min/h}}\right) = 45 \text{ mi/h}\left(0.41667 \text{ h} - \frac{5.0 \text{ min}}{60 \text{ min/h}}\right)$$

$$60.0 \text{ mi/h}(0.25 \text{ h}) = 45 \text{ mi/h}(0.33333 \text{ h})$$

$$15 \text{ mi} = 15 \text{ mi}$$

27. Let x – 30.0 s = time since the first car started moving in the race in seconds.
Let x= time since the second car started the race in seconds.
The distance travelled by each car will be the same at the point where the first car overtakes the second car.
Distance = speed × time.

$$260.0 \text{ ft/s}(x - 30.0 \text{ s}) = 240.0 \text{ ft/s}(x)$$

$$(260.0 \text{ ft/s})x - (260.0 \text{ ft/s})(30.0 \text{ s}) = (240.0 \text{ ft/s})x$$

$$(260.0 \text{ ft/s})x - 7800 \text{ ft} = (240.0 \text{ ft/s})x$$

$$(20.0 \text{ ft/s})x = 7800 \text{ ft}$$

$$x = \frac{7800 \text{ ft}}{20.0 \text{ ft/s}}$$

$$x = 390 \text{ s}$$

The first car will overtake the second car after 390 s. The first car travels 260 ft/s × (390 s – 30 s) = 93600 ft by this point. 8 laps around the track is 2.5 mi/lap. 8 laps × 5280 ft/mi = 105,600 ft, so the first car will already be in the lead at the end of the 8th lap.
Check:

$$260.0 \text{ ft/s}(390.0 \text{ s} - 30.0 \text{ s}) = 240.0 \text{ ft/s}(390 \text{ s})$$

$$260.0 \text{ ft/s}(360.0 \text{ s}) = 240.0 \text{ ft/s}(390 \text{ s})$$

$$93,600 \text{ ft} = 93,600 \text{ ft}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual


28. Let x = the number of the first chips that is defective 0.50%.
Let $6100 - x$ = the number of the second chips that is defective 0.80%.

$$0.0050(x) + 0.0080(6100 \text{ chips} - x) = 38 \text{ chips}$$
$$(0.0050)x + 48.8 \text{ chips} - (0.0080)x = 38 \text{ chips}$$
$$-(0.0030)x = -10.8 \text{ chips}$$
$$x = \frac{-10.8 \text{ chips}}{-0.0030}$$
$$x = 3600 \text{ chips}$$

There are 3600 chips that are 0.50% defective and (6100 chips – 3600 chips) = 2500 chips that are defective 0.80%.
Check:

$$0.0050(3600 \text{ chips}) + 0.0080(6100 \text{ chips} - 3600 \text{ chips}) = 38 \text{ chips}$$
$$18 \text{ chips} + 0.0080(2500 \text{ chips}) = 38 \text{ chips}$$
$$18 \text{ chips} + 20 \text{ chips} = 38 \text{ chips}$$
$$38 \text{ chips} = 38 \text{ chips}$$

29.




Assuming that the customer is located between the two gasoline distributors:
Let x = the distance in km to the first gasoline distributor that costs \$2.90/gal.
Let $228 \text{ mi} - x$= the distance in km to the second gasoline distributor that costs \$2.70/gal.

$$\$2.90 + \$0.002(x) = \$2.70 + \$0.002(228 - x)$$
$$\$2.90 + \$0.002(x) = \$2.70 + \$0.456 - \$0.002(x)$$
$$\$0.004(x) = \$0.256$$
$$x = \frac{\$0.256}{\$0.004}$$
$$x = 64 \text{ mi}$$

The customer is 64 mi away from the first gas distributor (\$2.90/gal) and (228 mi – 64 mi)= 164 mi away from the second gas distributor (\$2.70).
Check:

$$\$2.90 + \$0.002(64) = \$2.70 + \$0.002(228 - 64)$$
$$\$2.90 + \$0.128 = \$2.70 + \$0.002(164)$$
$$\$3.028 = \$2.70 + \$0.328$$
$$\$3.028 = \$3.028$$

30.



(100 % Gas) 8.0 L gas can (needs to be full of 93.75% gas/oil mixture)

A 15:1 gas/oil mixture is 15/16 gasoline = 93.75%.
Let x = the amount of 100% gasoline added in L.
Let $8.0 \text{ L} - x$ = the amount of 75% gasoline mixture in L.

Copyright © 2018 Pearson Education, Inc.


Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

$$1.00(x)+0.75(8.0\text{ L}-x)=0.9375(8.0\text{ L})$$
$$1.00(x)+6.0\text{ L}-0.75(x)=7.5\text{ L}$$
$$0.25(x)=1.5\text{ L}$$
$$x=\frac{1.5\text{ L}}{0.25}$$
$$x=6.0\text{ L}$$

6.0 L of 100% gasoline must be added to the 75% gas/oil mixture to make 8 L of 15:1 gasoline/ oil.
Check:

$$1.00(6.0\text{ L})+0.75(8.0\text{ L}-6.0\text{ L})=0.9375(8.0\text{ L})$$
$$6\text{ L}+0.75(2.0\text{ L})=7.5\text{ L}$$
$$6\text{ L}+1.5\text{ L}=7.5\text{ L}$$
$$7.5\text{ L}=7.5\text{ L}$$

31.



100% Antifreeze 12.0 L radiator (needs to be filled with 50% mixture)

Let x = the amount in L of 25% antifreeze left in radiator
Let 12.0 L – x = the amount of 100% antifreeze added in L.


$$0.25(x)+1.00(12.0\text{ L}-x)=0.5(12.0\text{ L})$$
$$0.25(x)+12.0\text{ L}-1.00(x)=6.0\text{ L}$$
$$-0.75(x)=-6.0\text{ L}$$
$$x=\frac{-6.0\text{ L}}{-0.75}$$
$$x=8.0\text{ L}$$

There needs to be 8L of 25% antifreeze left in radiator, so (12.0 L – 8.0 L) = 4.0 L must be drained.

Check:

$$0.25(8.0\text{ L})+1.00(12.0\text{ L}-8.0\text{ L})=0.5(12.0\text{ L})$$
$$2\text{ .0L}+1.00(4.0\text{ L})=6.0\text{ L}$$
$$2.0\text{ L}+4.0\text{ L}=6.0\text{ L}$$
$$6.0\text{ L}=6.0\text{ L}$$

32.



Let x = the amount of sand added.
Let 250 lb + x = the amount in lb of the final 25% sand mixture.

Copyright © 2018 Pearson Education, Inc.


Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

$$1.00(x)+0.22(250\text{ lb})=0.25(250\text{ lb}+x)$$
$$1.00(x)+55\text{ lb}=62.5\text{ lb}+0.25(x)$$
$$0.75(x)=7.5\text{ lb}$$
$$x=\frac{7.5\text{ lb}}{0.75}$$
$$x=10\text{ lb}$$

Check:

$$1.00(10\text{ lb})+0.22(250\text{ lb})=0.25(250\text{ lb}+10\text{ lb})$$
$$10\text{ lb}+55\text{ lb}=62.5\text{ lb}+2.5\text{ lb}$$
$$65\text{ lb}=65\text{ lb}$$

33.




Let x = the speed the car needs to travel in km/h to pass the semi in 10 s.
Speed = distance/time. 10 s is 10s/3600 s/h = 0.002777777 h.

$$x=\frac{\text{distance needed to pass truck + distance travelled by truck in 10s}}{10s}$$
$$x=\frac{0.025\text{ km}+70\text{ km/h}\left(0.0027777\text{ h}\right)}{0.0027777\text{ h}}$$
$$x=\frac{0.025\text{ km}+0.19444\text{ km}}{0.0027777\text{ h}}$$
$$x=\frac{2.19444\text{ km}}{0.0027777\text{ h}}$$
$$x=79\text{ km/h}$$

The car needs to travel at a speed of 79 km/h to pass the semitrailer in 10s.
Check:

$$79\text{ km/h}=\frac{0.025\text{ km}+70\text{ km/h}\left(0.0027777\text{ h}\right)}{0.0027777\text{ h}}$$
$$79\text{ km/h}=\frac{0.025\text{ km}+0.19444\text{ km}}{0.0027777\text{ h}}$$
$$79\text{ km/h}=79\text{ km/h}$$

34.



Let x = the time the first wave takes to travel to the seismic station in s.
Let x + 120 s = the time the first wave takes to travel to the seismic station in s.
Distance =speed × time. The distances travelled by both waves to the seismic station are the same. 2.0 min is (2.0 min × 60 s/min) = 120 s.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

$$8.0 \text{ km/s}(x) = 5.0 \text{ km/s}(x + 120 \text{ s})$$
$$8.0 \text{ km/s}(x) = 5.0 \text{ km/s}(x) + (5 \text{ km/s})(120 \text{ s})$$
$$3.0 \text{ km/s}(x) = 600 \text{ km}$$
$$x = \frac{600 \text{ km}}{3.0 \text{ km/s}}$$
$$x = 200 \text{ s}$$

The distance to the seismic station is (200 s × 8.0 km/s) = 1600 km.
Check:

$$8.0 \text{ km/s}(200 \text{ s}) = 5.0 \text{ km/s}(200 \text{ s} + 120 \text{ s})$$
$$1600 \text{ km} = 5.0 \text{ km/s}(320 \text{ s})$$
$$1600 \text{ km} = 1600 \text{ km}$$

Review Exercises

1. False, because $|0| = 0$ which is not a positive value.

2. True. The order of operations dictates performing the division first, then the subtraction.

3. False. The reported answer should have only two significant digits.

4. False. Had the problem been given as $(2a)^3 = 8a^3$, then it would be true.

5. True.

6. False. The left-hand side, $-\sqrt{-4}$, is not a real number, in fact.

7. False. The left-hand side simplifies to $4x - (2x + 3) = 4x - 2x - 3 = 2x - 3$.

8. True.

9. False. The left-hand side simplifies to $\frac{6x + 2}{2} = \frac{6x}{2} + \frac{2}{2} = 3x + 1$.

10. True.

11. False. Solving for c yields

$$a - bc = d$$
$$-bc = d - a$$
$$c = \frac{d - a}{-b}$$
$$c = -\frac{d - a}{b}$$

12. False. It is likely that one should set up a phrase such as ‘let x be the number of gears of the first type…’

13. $(-2) + (-5) - 3 = -2 - 5 - 3 = -10$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

14. $6-8-(-4)=6-8+4=2$

15. $\dfrac{(-5)(6)(-4)}{(-2)(3)}=\dfrac{(20)\cancel{(6)}}{-\cancel{(6)}}=-20$

16. $\dfrac{(-9)(-12)(-4)}{24}=\dfrac{108(-4)}{24}=\dfrac{-432}{24}=-18$

17. $-5-\left|2(-6)\right|+\dfrac{-15}{3}=-5-\left|-12\right|+(-5)=-5-12-5=-22$

18. $3-5\left|-3-2\right|-\dfrac{12}{-4}=3-5\left|-5\right|-(-3)=3-5(5)+3=6-25=-19$

19. $\dfrac{18}{3-5}-(-4)^2=\dfrac{18}{-2}-(-4)(-4)=-9-16=-25$

20. $-(-3)^2-\dfrac{-8}{(-2)-\left|-4\right|}=-(-3)(-3)-\dfrac{-8}{(-2)-4}=-9-\dfrac{-8}{-6}=-\dfrac{27}{3}-\dfrac{4}{3}=-\dfrac{31}{3}$

21. $\sqrt{16}-\sqrt{64}=\sqrt{(4)(4)}-\sqrt{(8)(8)}=4-8=-4$

22. $-\sqrt{81+144}=-\sqrt{225}=-\sqrt{(5)(5)(3)(3)}=-(3)(5)=-15$

23. $(\sqrt{7})^2-\sqrt[3]{8}=(\sqrt{7})(\sqrt{7})-\sqrt[3]{(2)(2)(2)}=7-2=5$

24. $-\sqrt[4]{16}+(\sqrt{6})^2=-\sqrt[4]{(2)(2)(2)(2)}+(\sqrt{6})(\sqrt{6})=-2+6=4$

25. $(-2rt^2)^2=(-2)^2r^2t^{2x2}=4r^2t^4$

26. $(3a^0b^{-2})^3=(3)^3(1)^3b^{-2\times3}=27(1)b^{-6}=\dfrac{27}{b^6}$

27. $-3mn^{-5}t(8m^{-3}n^4)=-(3)(8)m^{1-3}n^{-5+4}t=-24m^{-2}n^{-1}t=-\dfrac{24t}{m^2n}$

28. $\dfrac{15p^4q^2r}{5pq^5r}=\dfrac{3p^{4-1}\cancel{r}}{q^{5-2}\cancel{r}}=\dfrac{3p^3}{q^3}$

29. $\dfrac{-16N^{-2}(NT^2)}{-2N^0T^{-1}}=\dfrac{8N^{-2+1}T^{2+1}}{(1)}=\dfrac{8N^{-1}T^3}{(1)}=\dfrac{8T^3}{N}$

30. $\dfrac{-35x^{-1}y(x^2y)}{5xy^{-1}}=\dfrac{-7y^{1+1+1}x^2}{x^{1+1}}=\dfrac{-7y^3\cancel{x^2}}{\cancel{x^2}}=-7y^3$

31. $\sqrt{45}=\sqrt{(5)(3)(3)}=3\sqrt{5}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

32. $\sqrt{9+36} = \sqrt{45} = \sqrt{(5)(3)(3)} = 3\sqrt{5}$

33. 8000 has 1 significant digit. Rounded to 2 significant digits, it is $8\bar{0}00$.

34. 21490 has 4 significant digits. Rounded to 2 significant digits, it is 21000.

35. 9.050 has 4 significant digits. Rounded to 2 significant digits, it is 9.0.

36. 0.7000 has 4 significant digits. Rounded to 2 significant digits, it is 0.70.

37. $37.3 - 16.92(1.067)^2 = 37.3 - 16.92(1.138489)$
$= 37.3 - 19.26323388$
$= 18.03676612$
which rounds to 18.0.

38. $\dfrac{8.896 \times 10^{-12}}{-3.5954 - 6.0449} = \dfrac{8.896 \times 10^{-12}}{-9.6403}$
$= -9.227928591 \times 10^{-13}$
which rounds to -9.228×10^{-13}.

39. $\dfrac{\sqrt{0.1958 + 2.844}}{3.142(65)^2} = \dfrac{\sqrt{3.0398}}{3.142(4225)}$
$= \dfrac{1.743502223}{13274.95}$
$= 0.000131337$
which rounds to 1.3×10^{-4}.

40. $\dfrac{1}{0.03568} + \dfrac{37\ 466}{29.63^2} = 28.02690583 + \dfrac{37\ 466}{877.9369}$
$= 28.02690583 + 42.67504874$
$= 70.70195457$
which rounds to 70.70, assuming that the 1 is exact.

41. $875 \text{ Btu} = 875 \text{ Btu} \times \dfrac{778.2 \text{ ft} \cdot \text{lb}}{1 \text{ Btu}} \times \dfrac{1.356 \text{ J}}{1 \text{ ft} \cdot \text{lb}}$
$= 923,334.3 \text{ J}$
which rounds to 923,000 J.

42. $18.4 \text{ in} = 18.4 \text{ in} \times \dfrac{2.54 \text{ cm}}{1 \text{ in}} \times \dfrac{1 \text{ m}}{100 \text{ cm}}$
$= 0.46736 \text{ m}$
which rounds to 0.467 m.

43. $65 \dfrac{\text{km}}{\text{h}} = 65 \dfrac{\text{km}}{\text{h}} \times \dfrac{1 \text{ hr}}{3600 \text{ s}} \times \dfrac{0.6214 \text{ mi}}{1 \text{ km}} \times \dfrac{5280 \text{ ft}}{1 \text{ mi}}$
$= 59.2401333 \dfrac{\text{ft}}{\text{s}}$
which rounds to 59 ft/s.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

44. $12.25\ \dfrac{\text{g}}{\text{L}} = 12.25\ \dfrac{\text{g}}{\text{L}} \times \dfrac{28.32\ \text{L}}{1\ \text{ft}^3} \times \dfrac{1\ \text{kg}}{1000\ \text{g}} \times \dfrac{2.205\ \text{lb}}{1\ \text{kg}}$

$= 0.7649586\ \dfrac{\text{lb}}{\text{ft}^3}$

which rounds to $0.7650\ \text{lb/ft}^3$.

45. $225\ \text{hp} = 225\ \text{hp} \times \dfrac{550\ \text{ft} \cdot \text{lb/s}}{1\ \text{hp}} \times \dfrac{1.356\ \text{J}}{1\ \text{ft} \cdot \text{lb}} \times \dfrac{60\ \text{s}}{1\ \text{min}}$

$= 10068300\ \dfrac{\text{J}}{\text{min}}$

which rounds to $10100000 = 1.01 \times 10^7\ \text{J/min}$.

46. $89.7\ \dfrac{\text{lb}}{\text{in}^2} = 89.7\ \dfrac{\text{lb}}{\text{in}^2} \times \dfrac{4.448\ \text{N}}{1\ \text{lb}} \times \left(\dfrac{1\ \text{in}}{2.54\ \text{cm}}\right)^2$

$= 61.8428917\ \dfrac{\text{N}}{\text{cm}^2}$

which rounds to $61.8\ \text{N/cm}^2$.

47. $a - 3ab - 2a + ab = -2ab - a$

48. $xy - y - 5y - 4xy = -3xy - 6y$

49. $6LC - (3 - LC) = 6LC - 3 + LC = 7LC - 3$

50. $-(2x - b) - 3(-x - 5b) = -2x + b + 3x + 15b = 16b + x$

51. $(2x - 1)(5 + x) = (2x)(5) + (2x)(x) + (-1)(5) + (-1)(x)$

$= 10x + 2x^2 - 5 - x$

$= 2x^2 + 9x - 5$

52. $(C - 4D)(D - 2C) = (C)(D) + (C)(-2C) + (-4D)(D) + (-4D)(-2C)$

$= CD - 2C^2 - 4D^2 + 8CD$

$= -2C^2 + 9CD - 4D^2$

53. $(x + 8)^2 = (x + 8)(x + 8)$

$= (x)(x) + (x)(8) + (8)(x) + (8)(8)$

$= x^2 + 8x + 8x + 64$

$= x^2 + 16x + 64$

54. $(2r - 9s)^2 = (2r - 9s)(2r - 9s)$

$= (2r)(2r) + (2r)(-9s) + (-9s)(2r) + (-9s)(-9s)$

$= 4r^2 - 18rs - 18rs + 81s^2$

$= 4r^2 - 36rs + 81s^2$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

55. $$\frac{2h^3k^2 - 6h^4k^5}{2h^2k} = \frac{2h^3k^2}{2h^2k} - \frac{6h^4k^5}{2h^2k}$$
$$= h^{3-2}k^{2-1} - 3h^{4-2}k^{5-1}$$
$$= -3h^2k^4 + hk$$

56. $$\frac{4a^2x^3 - 8ax^4}{-2ax^2} = \frac{4a^2x^3}{-2ax^2} - \frac{8ax^4}{-2ax^2}$$
$$= -2a^{2-1}x^{3-2} + \frac{4\cancel{a}x^{4-2}}{\cancel{a}}$$
$$= 4x^2 - 2ax$$

57. $$4R - [2r - (3R - 4r)] = 4R - [2r - 3R + 4r]$$
$$= 4R - [6r - 3R]$$
$$= 4R - 6r + 3R$$
$$= 7R - 6r$$

58. $$-3b - [3a - (a - 3b)] + 4a = 4a - 3b - [3a - a + 3b]$$
$$= 4a - 3b - [2a + 3b]$$
$$= 4a - 3b - 2a - 3b$$
$$= 2a - 6b$$

59. $$2xy - \{3z - [5xy - (7z - 6xy)]\} = 2xy - \{3z - [5xy - 7z + 6xy)]\}$$
$$= 2xy - \{3z - [11xy - 7z]\}$$
$$= 2xy - \{3z - 11xy + 7z\}$$
$$= 2xy - \{10z - 11xy\}$$
$$= 2xy - 10z + 11xy$$
$$= 13xy - 10z$$

60. $$x^2 + 3b + [(b - y) - 3(2b - y + z)] = x^2 + 3b + [b - y - 6b + 3y - 3z)]$$
$$= x^2 + 3b + [-5b + 2y - 3z]$$
$$= x^2 + 3b - 5b + 2y - 3z$$
$$= x^2 - 2b + 2y - 3z$$

61. $$(2x + 1)(x^2 - x - 3) = (2x)(x^2) + (2x)(-x) + (2x)(-3) + (1)(x^2) + (1)(-x) + (1)(-3)$$
$$= 2x^3 - 2x^2 - 6x + x^2 - x - 3$$
$$= 2x^3 - x^2 - 7x - 3$$

62. $$(x - 3)(2x^2 + 1 - 3x) = (x)(2x^2) + (x)(1) + (x)(-3x) + (-3)(2x^2) + (-3)(1) + (-3)(-3x)$$
$$= 2x^3 + x - 3x^2 - 6x^2 - 3 + 9x$$
$$= 2x^3 - 9x^2 + 10x - 3$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

63. $-3y(x-4y)^2 = -3y(x-4y)(x-4y)$
$= -3y[(x)(x)+(x)(-4y)+(-4y)(x)+(-4y)(-4y)]$
$= -3y[x^2-4xy-4xy+16y^2]$
$= -3y[x^2-8xy+16y^2]$
$= -3x^2y+24xy^2-48y^3$

64. $-s(4s-3t)^2 = -s(4s-3t)(4s-3t)$
$= -s[(4s)(4s)+(4s)(-3t)+(-3t)(4s)+(-3t)(-3t)]$
$= -s[16s^2-12st-12st+9t^2]$
$= -s[16s^2-24st+9t^2]$
$= -16s^3+24s^2t-9st^2$

65. $3p[(q-p)-2p(1-3q)] = 3p[q-p-2p+6pq]$
$= 3p[q-3p+6pq]$
$= 18p^2q-9p^2+3pq$

66. $3x[2y-r-4(x-2r)] = 3x[2y-r-4x+8r]$
$= 3x[2y+7r-4x]$
$= 21rx-12x^2+6xy$

67. $\dfrac{12p^3q^2-4p^4q+6pq^5}{2p^4q} = \dfrac{12p^3q^2}{2p^4q}-\dfrac{4p^4q}{2p^4q}+\dfrac{6pq^5}{2p^4q}$
$= \dfrac{6q^{2-1}}{p^{4-3}}-\dfrac{2\cancel{p^4q}}{\cancel{p^4q}}+\dfrac{3q^{5-1}}{p^{4-1}}$
$= \dfrac{3q^4}{p^3}+\dfrac{6q}{p}-2$

68. $\dfrac{27s^3t^2-18s^4t+9s^2t}{-9s^2t} = \dfrac{27s^3t^2}{-9s^2t}-\dfrac{18s^4t}{-9s^2t}+\dfrac{9s^2t}{-9s^2t}$
$= -3s^{3-2}t^{2-1}+\dfrac{2s^{4-2}\cancel{t}}{\cancel{t}}-\dfrac{\cancel{9s^2t}}{\cancel{9s^2t}}$
$= 2s^2-3st-1$

69.
$$\begin{array}{r} 2x-5 \\ x+6\overline{)\,2x^2+7x-30} \\ \underline{2x^2+12x} \\ -5x-30 \\ \underline{-5x-30} \\ 0 \end{array}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

70.
$$\begin{array}{r} 2x-7 \\ 2x+7\overline{)4x^2+0x-41} \\ \underline{4x^2+14x} \\ -14x-41 \\ \underline{-14x-49} \\ 8 \end{array}$$
$$\frac{4x^2-41}{2x+7}=2x-7+\frac{8}{2x+7}$$

71.
$$\begin{array}{r} x^2-2x+3 \\ 3x-1\overline{)3x^3-7x^2+11x-3} \\ \underline{3x^3-x^2} \\ -6x^2+11x \\ \underline{-6x^2+2x} \\ 9x-3 \\ \underline{9x-3} \\ 0 \end{array}$$

72.
$$\begin{array}{r} w^2-w+4 \\ w-3\overline{)w^3-4w^2+7w-12} \\ \underline{w^3-3w^2} \\ -w^2+7w \\ \underline{-w^2+3w} \\ 4w-12 \\ \underline{4w-12} \\ 0 \end{array}$$

73.
$$\begin{array}{r} 4x^3-2x^2+6x \\ x+3\overline{)4x^4+10x^3+0x^2+18x-1} \\ \underline{4x^4+12x^3} \\ -2x^3+0x^2 \\ \underline{-2x^3-6x^2} \\ 6x^2+18x \\ \underline{6x^2+18x} \\ 0x-1 \end{array}$$
$$\frac{4x^4+10x^3+18x-1}{x+3}=4x^3-2x^2+6x-\frac{1}{x+3}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

74.
$$2x+3\overline{)8x^3+0x^2-14x+3}$$
$$\begin{array}{r} 4x^2-6x+2 \\ 8x^3+12x^2 \\ \hline -12x^2-14x \\ -12x^2-18x \\ \hline 4x+3 \\ 4x+6 \\ \hline -3 \end{array}$$
$$\frac{8x^3-14x+3}{2x+3}=4x^2-6x+2-\frac{3}{2x+3}$$

75.
$$\begin{aligned} -3\{(r+s-t)-2[(3r-2s)-(t-2s)]\} &= -3\{r+s-t-2[3r-2s-t+2s)]\} \\ &= -3\{r+s-t-2[3r-t]\} \\ &= -3\{r+s-t-6r+2t]\} \\ &= -3\{-5r+s+t\} \\ &= 15r-3s-3t \end{aligned}$$

76.
$$\begin{aligned} &(1-2x)(x-3)-(x+4)(4-3x) \\ &=[(1)(x)+(1)(-3)+(-2x)(x)+(-2x)(-3)]-[(x)(4)+(x)(-3x)+(4)(4)+(4)(-3x)] \\ &=[x-3-2x^2+6x]-[4x+-3x^2+16+-12x] \\ &=[-2x^2+7x-3]-[-3x^2-8x+16] \\ &=-2x^2+7x-3+3x^2+8x-16 \\ &=x^2+15x-19 \end{aligned}$$

77.
$$2y-1\overline{)2y^3+9y^2-7y+5}$$
$$\begin{array}{r} y^2+5y-1 \\ 2y^3+9y^2 \\ \hline 10y^2-7y \\ 10y^2-5y \\ \hline -2y+5 \\ -2y+1 \\ \hline 4 \end{array}$$
$$\frac{2y^3+9y^2-7y+5}{2y-1}=y^2+5y-1+\frac{4}{2y-1}$$

78.
$$2x-y\overline{)6x^2+5xy-4y^2}$$
$$\begin{array}{r} 3x+4y \\ 6x^2-3xy \\ \hline 8xy-4y^2 \\ 8xy-4y^2 \\ \hline 0 \end{array}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

79. $3x+1=x-8$

$2x=-9$

$x=-\frac{9}{2}$

80. $4y-3=5y+7$

$-y=10$

$y=-10$

81. $\frac{5x}{7}=\frac{3}{2}$

$2(5x)=3(7)$

$10x=21$

$x=\frac{21}{10}$

82. $\frac{2(N-4)}{3}=\frac{5}{4}$

$\frac{2N-8}{3}=\frac{5}{4}$

$4(2N-8)=3(5)$

$8N-32=15$

$8N=47$

$N=\frac{47}{8}$

83. $-6x+5=-3(x-4)$

$-6x+5=-3x+12$

$-3x=7$

$x=-\frac{7}{3}$

84. $-2(-4-y)=3y$

$8+2y=3y$

$y=8$

85. $2s+4(3-s)=6$

$2s+12-4s=6$

$-2s=-6$

$s=\frac{-6}{-2}$

$s=3$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

86. $2|x|-1=3$

$2|x|=4$

$|x|=\frac{4}{2}$

$|x|=2$

$x=-2$ and 2

87. $3t-2(7-t)=5(2t+1)$

$3t-14+2t=10t+5$

$5t-14=10t+5$

$-5t=19$

$t=-\frac{19}{5}$

88. $-(8-x)=x-2(2-x)$

$-8+x=x-4+2x$

$-8+x=3x-4$

$-2x=4$

$x=-\frac{4}{2}$

$x=-2$

89. $2.7+2.0(2.1x-3.4)=0.1$

$2.7+4.2x-6.8=0.1$

$4.2x-4.1=0.1$

$4.2x=4.2$

$x=\frac{4.2}{4.2}$

$x=1.0$

90. $0.250(6.721-2.44x)=2.08$

$1.68025-0.610x=2.08$

$-0.610x=0.39975$

$x=-\frac{0.39975}{0.610}$

$x=0.655327868$

$x=0.655$

91. 60,000,000,000,000 bytes = 6×10^{13} bytes

92. 25,000 mi/h = 2.5×10^{4} mi/h

93. 15,400,000,000 km = 1.54×10^{10} km

94. 1.02×10^{9} Hz = 1,020,000,000 Hz

95. 2.53×10^{13} mi = 25,300,000,000,000 mi

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

96. 10^7 ft^2 = 10,000,000 ft^2

97. 10^{-12} W/m^2 = 0.000000000001 W/m^2

98. 0.00000015 m = 1.5×10^{-7} m

99. 1.5×10^{-1} Bq/L = 0.15 Bq/L

100. 0.00000018 m = 1.8×10^{-7} m

101. $V = \pi r^2 L$

$$L = \frac{V}{\pi r^2}$$

102. $$R = \frac{2GM}{c^2}$$

$$c^2 R = 2GM$$

$$G = \frac{c^2 R}{2M}$$

103. $$P = \frac{\pi^2 EI}{L^2}$$

$$L^2 P = \pi^2 EI$$

$$E = \frac{L^2 P}{\pi^2 I}$$

104. $$f = p(c-1) - c(p-1)$$

$$f = cp - p - cp + c$$

$$f - c = -p$$

$$p = c - f$$

105. $Pp + Qq = Rr$

$$Qq = Rr - Pp$$

$$q = \frac{Rr - Pp}{Q}$$

106. $V = IR + Ir$

$$IR = V - Ir$$

$$R = \frac{V - Ir}{I}$$

107. $$d = (n-1)A$$

$$d = An - A$$

$$d + A = An$$

$$n = \frac{d + A}{A}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

108.
$$mu = (m + M)v$$
$$mu = mv + Mv$$
$$mu - mv = Mv$$
$$M = \frac{mu - mv}{v}$$

109.
$$N_1 = T(N_2 - N_3) + N_3$$
$$N_1 - N_3 = N_2T - N_3T$$
$$N_2T = N_1 - N_3 + N_3T$$
$$N_2 = \frac{N_1 - N_3 + N_3T}{T}$$

110.
$$Q = \frac{kAt(T_2 - T_1)}{L}$$
$$QL = kAt(T_2 - T_1)$$
$$T_2 - T_1 = \frac{QL}{kAt}$$
$$-T_1 = \frac{QL}{kAt} - T_2$$
$$T_1 = T_2 - \frac{QL}{kAt}$$

111.
$$R = \frac{A(T_2 - T_1)}{H}$$
$$HR = AT_2 - AT_1$$
$$AT_2 = HR + AT_1$$
$$T_2 = \frac{HR + AT_1}{A}$$

112.
$$Z^2\left(1 - \frac{\lambda}{2a}\right) = k$$
$$Z^2 - \frac{Z^2\lambda}{2a} = k$$
$$Z^2 - k = \frac{Z^2\lambda}{2a}$$
$$2a(Z^2 - k) = Z^2\lambda$$
$$\lambda = \frac{2aZ^2 - 2ak}{Z^2}$$

113.
$$d = kx^2[3(a + b) - x]$$
$$d = kx^2[3a + 3b - x]$$
$$d = 3akx^2 + 3bkx^2 - kx^3$$
$$3akx^2 = d - 3bkx^2 + kx^3$$
$$a = \frac{d - 3bkx^2 + kx^3}{3kx^2}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

114.
$$V = V_0[1 + 3a(T_2 - T_1)]$$
$$V = V_0[1 + 3aT_2 - 3aT_1]$$
$$V = V_0 + 3aT_2V_0 - 3aT_1V_0$$
$$3aT_2V_0 = V - V_0 + 3aT_1V_0$$
$$T_2 = \frac{V - V_0 + 3aT_1V_0}{3aV_0}$$

115. $\dfrac{5.25\times10^{13}\text{ bytes}}{6.4\times10^{4}\text{ bytes}} = 8.203125\times10^{8}$
which rounds to 8.2×10^8. The newer computer's memory is 8.2×10^8 larger.

116. $t = 0.25\sqrt{66} = 2.0310096$ s
which rounds to 2.0 s. It would take the person 2.0 s to fall 66 ft.

117. $\dfrac{0.553\text{ km}}{0.442\text{ km}} = 1.25113122$
which rounds to 1.25. The CN Tower is 1.25 times taller than the Sears tower.

118. $t = \left(\dfrac{4.8\times10^{3}\text{ cells}}{2650}\right)^2 = (1.8113207)^2 = 3.280882876$ s
which rounds to 3.28 s. It would take the computer 3.28 s to check 4800 memory cells.

119.
$$\frac{R_1R_2}{R_1 + R_2} = \frac{(0.0275\ \Omega)(0.0590\ \Omega)}{0.0275\ \Omega + 0.0590\ \Omega}$$
$$= \frac{0.0016225\ \Omega^2}{0.0865\ \Omega}$$
$$= 0.018757225\ \Omega$$
which rounds to $0.0188\,\Omega$. The combined electric resistance is $0.0188\ \Omega$.

120.
$$1.5\times10^{11}\sqrt{\frac{m}{M}} = 1.5\times10^{11}\sqrt{\frac{5.98\times10^{24}\text{ kg}}{1.99\times10^{30}\text{ kg}}}$$
$$= 1.5\times10^{11}\sqrt{0.000003005}$$
$$= 1.5\times10^{11}(0.0017335)$$
$$= 260\ 025\ 124.4\text{ m}$$
which rounds to 2.6×10^8 m. The distance the space craft will be from the earth is 2.6×10^8 m.

121.
$$(x - 2a) + 3\text{ ft/yd} \cdot (x + 2a) = x - 2a + 3x + 3(2a)$$
$$= x - 2a + 3x + 6a$$
$$= 4x + 4a$$
The sum of their length is $4x + 4a$ ft.

122.
$$(Ai - R)(1 + i)^2 = (Ai - R)(1 + i)(1 + i)$$
$$= (Ai - R)[(1)(1) + (1)(i) + (i)(1) + (i)(i)]$$
$$= (Ai - R)[i^2 + 2i + 1]$$
$$= (Ai)(i^2) + (Ai)(2i) + (Ai)(1) + (-R)(i^2) + (-R)(2i) + (-R)(1)$$
$$= Ai^3 + 2Ai^2 + Ai - i^2R - 2iR - R$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

123. $4(t+h)-2(t+h)^2 = 4t+4h-2(t+h)(t+h)$
$= 4t+4h-2[(t)(t)+(t)(h)+(h)(t)+(h)(h)]$
$= 4t+4h-2[t^2+2ht+h^2]$
$= 4t+4h-2t^2-4ht-2h^2$
$= -2t^2-2h^2-4ht+4t+4h$

124. $\dfrac{k^2r-2h^2k+h^2rv^2}{k^2r} = \dfrac{k^2r}{k^2r}-\dfrac{2h^2k}{k^2r}+\dfrac{h^2rv^2}{k^2r}$
$= \dfrac{\cancel{k^2}\cancel{r}}{\cancel{k^2}\cancel{r}}-\dfrac{2h^2}{k^{2-1}r}+\dfrac{h^2\cancel{r}v^2}{k^2\cancel{r}}$
$= 1-\dfrac{2h^2}{kr}+\dfrac{h^2v^2}{k^2}$

125. $3\times18\div(9-6) = 54\div(3) = 18$
$3\times18\div9-6 = 54\div9-6 = 6-6 = 0$
Yes, the removal of the parentheses does affect the answer.

126. $(3\times18)\div9-6 = 54\div9-6 = 6-6 = 0$
$3\times18\div9-6 = 54\div9-6 = 6-6 = 0$
No, the removal of the parentheses does not affect the answer.

127. $x-(3-x) = 2x-3$
$x-3+x = 2x-3$
$2x-3 = 2x-3$
The equation is valid for all values of the unknown, so the equation is an identity.

128. $7-(2-x) = x+2$
$7-2+x = x+2$
$x+5 = x+2$
$5 = 2$
The equation has no values of the unknown for which it is valid, so the equation is a contradiction.

129. (a) $2|2|-2|4| = 4-8$
$= -4$

(b) $2|2-(-4)| = 2|6|$
$= 12$

130. For $a<0$, $|a|=-a$.

131. Given $3-x=0$,
$|3-x|+7 = 2x$
$-(3-x)+7 = 2x$
$-3+x+7 = 2x$
$4 = x$
This is consistent with $3-x=0$, so $x=4$.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

132. $|x-4|+6=3x$

$|x-4|=3x-6$

$x-4=3x-6$ or $-(x-4)=3x-6$

$2=2x$ $\quad$ $4-x=3x-6$

$x=1$ $\quad$ $10=4x$

and so the only possible solutions are

$x=1$ or $x=5/2$.

The first possibility, $x=1$, yields $|-3|+6=3$ or $9=3$, which is false.

The second possibility, $x=5/2$, yields $|-3/2|+6=15/2$ or $15/2=15/2$, which is true, and so the only solution is $x=5/2$.

133. $(x-y)^3=(x-y)(x-y)(x-y)$

$=(-(y-x))(-(y-x))(-(y-x))$

$=-(y-x)(y-x)(y-x)$

$=-(y-x)^3$

134. Generally, $(a\div b)\div c\neq a\div(b\div c)$

We demonstrate this using $a=8, b=4, c=2$:

$(8\div 4)\div 2=2\div 2=1$

$8\div(4\div 2)=8\div 2=4$

Division is not associative.

135. $\dfrac{8\times10^{-3}}{2\times10^{4}}=4\times10^{-7}$

136. $\dfrac{\sqrt{4+36}}{\sqrt{4}}=\dfrac{\sqrt{(2)(2)(10}}{2}=\dfrac{2\sqrt{10}}{2}=\sqrt{10}$

137. $250\text{ hp}=250\text{ hp}\times\dfrac{746.0\text{ W}}{1\text{ hp}}\times\dfrac{1\text{ kW}}{1000\text{ W}}$

$=186.5\text{ kW}$

This is rounded to 190 kW.

138. $32\ \dfrac{\text{lb}}{\text{in}^2}=32\ \dfrac{\text{lb}}{\text{in}^2}\times\dfrac{4.448\text{ N}}{1\text{ lb}}\times\left(\dfrac{1\text{ in}}{2.54\text{ cm}}\right)^2\times\left(\dfrac{100\text{ cm}}{1\text{ m}}\right)^2$

$=220{,}621.241\ \dfrac{\text{N}}{\text{m}^2}$

This is rounded to 220,000 N/m^2.

139. $110\text{ N}\cdot\text{m}=110\text{ N}\cdot\text{m}\times\dfrac{1\text{ lb}}{4.448\text{ N}}\times\dfrac{1\text{ ft}}{0.3048\text{ m}}$

$=81.1358787\text{ ft}\cdot\text{lb}$

This is rounded to 81 foot pounds.

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

140. $1.2\times10^{6}\ \dfrac{\text{A}}{\text{m}^2}=1.2\times10^{6}\ \dfrac{\text{A}}{\text{m}^2}\times\dfrac{1000\text{ mA}}{1\text{ A}}\times\left(\dfrac{1\text{ m}}{100\text{ cm}}\right)^2$

$$=1.2\times10^{5}\ \frac{\text{mA}}{\text{cm}^2}$$

141. Let x = the cost of the first computer program.
Let $x + \$72$ = the cost of the second computer program.

$$x+(x+\$72)=\$190$$
$$2x+\$72=\$190$$
$$2x=\$118$$
$$x=\frac{\$118}{2}$$
$$x=\$59$$

The cost of the first computer program is \$59, and the other program costs (\$59 + \$72) = \$131.
Check: \$59 + \$131 = 190

142. Let x = the cost to run the commercial on the first station.
Let $x + \$1100$ = the cost to run the commercial on the second station.

$$x+(x+\$1100)=\$9500$$
$$2x+\$1100=\$9500$$
$$2x=\$8400$$
$$x=\frac{\$8400}{2}$$
$$x=\$4200$$

The cost of the run the commercial on the first station is \$4200, and the cost for the other station is (\$4200 + \$1100) = \$5300.
Check: \$4200 + \$5300 = \$9500

143. Let $2x$ = the amount of oxygen produced in cm^3 by the first reaction.
Let x = the amount of oxygen produced in cm^3 by the second reaction.
Let $4x$ = the amount of oxygen produced in cm^3 by the third reaction.

$$2x+x+4x=560\text{ cm}^3$$
$$7x=560\text{ cm}^3$$
$$x=\frac{560\text{ cm}^3}{7}$$
$$x=80\text{ cm}^3$$

The first reaction produces $(2\times80\text{ cm}^3)=160\text{ cm}^3$ of oxygen, the second reaction produces 80 cm^3 of oxygen, and the third reaction produces $(4\times80\text{ cm}^3)=320\text{ cm}^3$ of oxygen.
Check: $160\text{ cm}^3+80\text{ cm}^3+320\text{ cm}^3=560\text{ cm}^{3*}$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

144. Let x = the speed that the river is flowing in mi/h.
Let $x + 5.5$ mi/h = the speed that the boat travels downstream.
Let $-x + 5.5$ mi/h = the speed that the boat travels upstream.
The distance that the boat travelled is the same in both experiments. Distance = speed × time.

$$(x+5.5 \text{ mi/h})(5.0 \text{ h}) = (-x+5.5 \text{ mi/h})(8.0 \text{ h})$$
$$(5.0 \text{ h})(x) + (5.5 \text{ mi/h})(5.0 \text{ h}) = (8.0 \text{ h})(-x) + (5.5 \text{ mi/h})(8.0 \text{ h})$$
$$(5.0 \text{ h})(x) + (27.5 \text{ mi}) = (-8.0 \text{ h})(x) + (44 \text{ mi})$$
$$(13.0 \text{ h})x = 16.5 \text{ mi}$$
$$x = \frac{16.5 \text{ mi}}{13 \text{ h}}$$
$$x = 1.269230769 \text{ mi/h}$$

which rounds to 1.3 mi/h. The polluted stream is flowing at 1.3 mi/h.
Check:

$$(1.269230769 \text{ mi/h} + 5.5 \text{ mi/h})(5.0 \text{ h}) = (-1.269230769 \text{ mi/h} + 5.5 \text{ mi/h})(8.0 \text{ h})$$
$$(6.769230769 \text{ mi/h})(5.0 \text{ h}) = (4.2 \text{ mi/h})(8.0 \text{ h})$$
$$(33.8 \text{ mi}) = (33.8 \text{ mi})$$

145. Let x = the resistance in the first resistor in Ω.
Let $x + 1200\,\Omega$ = the resistance in the second resistor in Ω.
Voltage = current × resistance. $2.4\ \mu A = 2.3 \times 10^{-6}$ A. 12 mV = 0.0120 V

$$(2.4\times10^{-6}\text{A})(x) + (2.4\times10^{-6}\text{A})(x + 1200\ \Omega) = 0.0120 \text{ V}$$
$$(2.4\times10^{-6}\text{A})(x) + (2.4\times10^{-6}\text{A})(x) + (2.4\times10^{-6}\text{A})(1200\ \Omega) = 0.0120 \text{ V}$$
$$(4.8\times10^{-6}\text{A})(x) + (0.00288\text{V}) = 0.0120 \text{ V}$$
$$(4.0\times10^{-6}\text{A})(x) = 0.00912 \text{ V}$$
$$x = \frac{0.00912 \text{ V}}{4.8\times10^{-6}\text{A}}$$
$$x = 1900\ \Omega$$

The first resistor's resistance is $1900\ \Omega$ and the second resistor's is $(1900\ \Omega + 1200\ \Omega) = 3100\ \Omega$.
Check:

$$(2.4\times10^{-6}\text{A})(1900\ \Omega) + (2.4\times10^{-6}\text{A})(1900\ \Omega + 1200\ \Omega) = 0.0120 \text{ V}$$
$$0.00456 \text{ V} + 0.00744 \text{ V} = 0.0120 \text{ V}$$
$$0.0120 \text{ V} = 0.0120 \text{ V}$$

146. Let x = the concentration of the first pollutant in ppm.
Let $4x$ = the concentration of the second pollutant in ppm.

$$x + 4x = 4.0 \text{ ppm}$$
$$5x = 4.0 \text{ ppm}$$
$$x = \frac{4.0 \text{ ppm}}{5}$$
$$x = 0.8 \text{ ppm}$$

The concentration of the first pollutant is 0.8 ppm, and the concentration of the second is (4 × 0.8 ppm) = 3.2 ppm.
Check:

$$0.8 \text{ ppm} + 4(0.8 \text{ ppm}) = 4.0 \text{ ppm}$$
$$0.8 \text{ ppm} + 3.2 \text{ ppm} = 4.0 \text{ ppm}$$
$$4.0 \text{ ppm} = 4.0 \text{ ppm}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

147. Let x = the time taken in hours for the crew to build 250 m of road.
The crew works at a rate of 450 m/12 h, which is 37.5 m/h. Time = distance / speed.

$$x = \frac{250 \text{ m}}{37.5 \text{ m/h}}$$
$$x = 6.666666667 \text{ h}$$

which rounds to 6.7 h.


148. Let x = the amount of oil in L in the mixture.
Let $15x$ = the amount of gas in L in the mixture.

$$x + 15x = 6.6 \text{ L}$$
$$16x = 6.6 \text{ L}$$
$$x = \frac{6.6 \text{ L}}{16}$$
$$x = 0.4125 \text{ L}$$

which rounds to 0.41 L. There is 0.41 L of oil in the mixture and (15 × 0.41 L) = 6.2 L of gas.
Check:

$$0.4125 \text{ L} + 15(0.4125 \text{ L}) = 6.6 \text{ L}$$
$$0.4125 \text{ L} + 6.1875 \text{ L} = 6.6 \text{ L}$$
$$6.6 \text{ L} = 6.6 \text{ L}$$

149.



Let x = the time taken by the second ship in hours.
Let $x + 2$ h = the amount time taken by the first ship in hours.
The distance travelled adds up to 634km. Distance = speed × time.

$$21.8 \text{ km/h}(x) + 17.4 \text{ km/h}(x + 2 \text{ h}) = 634 \text{ km}$$
$$21.8 \text{ km/h}(x) + 17.4 \text{ km/h}(x) + 17.4 \text{ km/h}(2 \text{ h}) = 634 \text{ km}$$
$$39.2 \text{ km/h}(x) + 34.8 \text{ km} = 634 \text{ km}$$
$$39.2 \text{ km/h}(x) = 599.2 \text{ km}$$
$$x = \frac{599.2 \text{ km}}{39.2 \text{ km/h}}$$
$$x = 15.2857 \text{ h}$$

which rounds to 15.2 h. The ships will pass 15.2 h after the second ship enters the canal.
Check:

$$21.8 \text{ km/h}(15.2857 \text{ h}) + 17.4 \text{ km/h}(15.2857 \text{ h} + 2 \text{ h}) = 634 \text{ km}$$
$$333.23 \text{ km} + 300.77 \text{ km} = 634 \text{ km}$$
$$634 \text{ km} = 634 \text{ km}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

150. Let x = the time take in h for the helicopter to travel from the pond to the fire.
Let 0.5 h $-x$ = the time take in h for the helicopter to travel from the fire to the pond.

30 min / 60 min/h = 0.5 h. The distance travelled by the helicopter is the same for both trips. Distance = speed × time.

$$105 \text{ mi/h}(0.5 \text{ h} - x) = 70 \text{ mi/h}(x)$$
$$52.5 \text{ mi} - 105 \text{ mi/h}(x) = 70 \text{ mi/h}(x)$$
$$52.5 \text{ mi} = 175 \text{ mi/h}(x)$$
$$x = \frac{52.5 \text{ mi}}{175 \text{ mi/h}}$$
$$x = 0.3 \text{ h}$$

which is reported as 0.30 h to two significant digits. It will take the helicopter 0.30 h to fly from the pond to the fire.
Check:

$$105 \text{ mi/h}(0.5 \text{ h} - 0.3 \text{ h}) = 70 \text{ mi/h}(0.3 \text{ h})$$
$$105 \text{ mi/h}(0.2 \text{ h}) = 70 \text{ mi/h}(0.3 \text{ h})$$
$$21 \text{ mi} = 21 \text{ mi}$$

151. Let x = the number of litres of 0.50% grade oil used.
Let 1000L $- x$ the number of litres of 0.75% grade oil used.

$$0.005(x) + 0.0075(1000 \text{ L} - x) = 0.0065(1000 \text{ L})$$
$$0.005(x) + 7.5 \text{ L} - 0.0075(x) = 6.5 \text{ L}$$
$$-0.0025(x) = -1.0 \text{ L}$$
$$x = \frac{-1.0 \text{ L}}{-0.0025}$$
$$x = 400 \text{ L}$$

It will take 400 L of the 0.50% grade oil and (1000 L – 400 L) = 600 L of the 0.75% grade oil to make 1000 L of 0.65% grade oil.
Check:

$$0.005(400 \text{ L}) + 0.0075(1000 \text{ L} - 400 \text{ L}) = 0.0065(1000 \text{ L})$$
$$2 \text{ L} + 4.5 \text{ L} = 6.5 \text{ L}$$
$$6.5 \text{ L} = 6.5 \text{ L}$$

152. Let x = the amount of rock containing 72 L/Mg of oil .
Let $18000 - x$ = the remaining amount of rock containing 150 L/Mg of oil.

$$(72 \text{ L/Mg})(x) + (150 \text{ L/Mg})(18000 \text{ Mg} - x) = (120 \text{ L/Mg})(18000 \text{ Mg})$$
$$72 \text{ L/Mg}(x) + 2700000 \text{ L} - 150\text{L/Mg}(x) = 2160000 \text{ L}$$
$$-78 \text{ L/Mg}(x) = -540000 \text{ L}$$
$$x = \frac{-540000 \text{ L}}{-78 \text{ L/Mg}}$$
$$x = 6923.07692 \text{ Mg}$$

which rounds to 6900 Mg. It will take 6900 Mg of 72 L/Mg rock and 11100 Mg of 150 L/Mg rock to make the 18000 Mg of 120 L/Mg rock.
Check:

$$(72 \text{ L/Mg})(6923.07692 \text{ Mg}) + (150 \text{ L/Mg})(18000 \text{ Mg} - 6923.07692 \text{ Mg}) = (120 \text{ L/Mg})(18000 \text{ Mg})$$
$$498461.538 \text{ L} + 2700000 \text{ L} - 1038461.538 \text{ L} = 2160000 \text{ L}$$
$$2160000 \text{ L} = 2160000 \text{ L}$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual

153. Let x = the area of space in ft^2 in the kitchen and bath.

$$\frac{\text{ft}^2 \text{ of tile in the house}}{\text{ft}^2 \text{ in the house}} = 0.25$$

$$\frac{x + 0.15(2200 \text{ ft}^2)}{(x + 2200 \text{ ft}^2)} = 0.25$$

$$x + 330 \text{ ft}^2 = 0.25(x) + (0.25)(2200 \text{ ft}^2)$$

$$x + 330 \text{ ft}^2 = 0.25(x) + 550 \text{ ft}^2$$

$$0.75x = 220 \text{ ft}^2$$

$$x = \frac{220 \text{ ft}^2}{0.75}$$

$$x = 293.33333333 \text{ ft}^2$$

which rounds to 290 ft^2. The kitchen and bath area is 290 ft^2.

Check:

$$\frac{293.33333333 \text{ ft}^2 + 0.15(2200 \text{ ft}^2)}{(293.33333333 \text{ ft}^2 + 2200 \text{ ft}^2)} = 0.25$$

$$\frac{623.3333333 \text{ ft}^2}{2493.3333333 \text{ ft}^2} = 0.25$$

$$0.25 = 0.25$$

154. Let x = the number of grams of 9-karat gold.

Let 200 g – x = the number of grams of 18-karat gold. 9-karat gold is 9/24 gold = 0.375, 18-karat gold is 18/24 gold= 0.75, and 14-karat gold is 14/24 gold = 0.583333333.

$$0.375(x) + 0.75(200 \text{ g} - x) = 0.583333333(200 \text{ g})$$

$$0.375(x) + 150 \text{ g} - 0.75(x) = 116.6666666 \text{ g}$$

$$-0.375(x) = -33.3333334 \text{ g}$$

$$x = \frac{-33.3333334 \text{ g}}{-0.375}$$

$$x = 88.88888907 \text{ g}$$

which rounds to 89 g. There is 89 g of 9-karat gold and (200 g – 89 g) = 111 g of 18-karat gold needed to make 200 g of 14-karat gold.

Check:

$$0.375(88.88888907 \text{ g}) + 0.75(200 \text{ g} - 88.88888907 \text{ g}) = 0.583333333(200 \text{ g})$$

$$33.3333334 \text{ g} + 83.3333332 \text{ g} = 116.6666666 \text{ g}$$

$$116.6666666 \text{ g} = 116.6666666 \text{ g}$$

155.

$$P = P_0 + P_0 rt$$

$$P - P_0 = P_0 rt$$

$$r = \frac{P - P_o}{P_0 t}$$

$$r = \frac{\$7625 - \$6250}{\$6250(4.000 \text{ years})}$$

$$r = \frac{\$1375}{25\,000}$$

$$r = 0.055$$

The rate is equal to 5.500%.

On the calculator type:

$$(7625 - 6250) / (6250 \times 4.000)$$

Copyright © 2018 Pearson Education, Inc.

Basic Technical Mathematics with Calculus 11th edition Washington Solutions Manual