Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


SKU: 9780134686981-SOLUTIONS

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Chapter 2
Linear and Quadratic Functions

Section 2.1

1. From the equation $y = 2x - 3$, we see that the y-intercept is -3. Thus, the point $(0, -3)$ is on the graph. We can obtain a second point by choosing a value for x and finding the corresponding value for y. Let $x = 1$, then $y = 2(1) - 3 = -1$. Thus, the point $(1, -1)$ is also on the graph. Plotting the two points and connecting with a line yields the graph below.



2. $m = \dfrac{y_2 - y_1}{x_2 - x_1} = \dfrac{3-5}{-1-2} = \dfrac{-2}{-3} = \dfrac{2}{3}$

3. $f(2) = 3(2)^2 - 2 = 10$

$f(4) = 3(4)^2 - 2 = 46$

$\dfrac{\Delta y}{\Delta x} = \dfrac{f(4) - f(2)}{4-2} = \dfrac{46-10}{4-2} = \dfrac{36}{2} = 18$

4. $60x - 900 = -15x + 2850$

$75x - 900 = 2850$

$75x = 3750$

$x = 50$

The solution set is $\{50\}$.

5. $f(-2) = (-2)^2 - 4 = 4 - 4 = 0$

6. True

7. slope; y-intercept

8. positive

9. True

10. False. The y-intercept is 8. The average rate of change is 2 (the slope).


11. a

12. d

13. $f(x) = 2x + 3$

a. Slope = 2; y-intercept = 3

b. Plot the point (0, 3). Use the slope to find an additional point by moving 1 unit to the right and 2 units up.




c. average rate of change = 2

d. increasing

14. $g(x) = 5x - 4$

a. Slope = 5; y-intercept = –4

b. Plot the point $(0, -4)$. Use the slope to find an additional point by moving 1 unit to the right and 5 units up.



c. average rate of change = 5

d. increasing


Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

15. $h(x) = -3x + 4$

a. Slope = -3; y-intercept = 4

b. Plot the point (0, 4). Use the slope to find an additional point by moving 1 unit to the right and 3 units down.




c. average rate of change = -3

d. decreasing

16. $p(x) = -x + 6$

a. Slope = -1; y-intercept = 6

b. Plot the point (0, 6). Use the slope to find an additional point by moving 1 unit to the right and 1 unit down.




c. average rate of change = -1

d. decreasing

17. $f(x) = \frac{1}{4}x - 3$

a. Slope = $\frac{1}{4}$; y-intercept = -3

b. Plot the point $(0, -3)$. Use the slope to find an additional point by moving 4 units to the right and 1 unit up.




c. average rate of change = $\frac{1}{4}$

d. increasing

18. $h(x) = -\frac{2}{3}x + 4$

a. Slope = $-\frac{2}{3}$; y-intercept = 4

b. Plot the point (0, 4). Use the slope to find an additional point by moving 3 units to the right and 2 units down.




c. average rate of change = $-\frac{2}{3}$

d. decreasing

19. $F(x) = 4$

a. Slope = 0; y-intercept = 4

b. Plot the point (0, 4) and draw a horizontal line through it.



c. average rate of change = 0

d. constant


Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

20. $G(x) = -2$

a. Slope = 0; y-intercept = -2

b. Plot the point $(0,-2)$ and draw a horizontal line through it.



c. average rate of change = 0


d. constant

21. $g(x) = 2x - 8$

a. zero: $0 = 2x - 8$: y-intercept = -8

$x = 4$

b. Plot the points $(4,0),(0,-8)$.




22. $g(x) = 3x + 12$

a. zero: $0 = 3x + 12$: y-intercept = 12

$x = -4$

b. Plot the points $(-4,0),(0,12)$.




23. $f(x) = -5x + 10$

a. zero: $0 = -5x + 10$: y-intercept = 10

$x = 2$

b. Plot the points 1 unit to the right and 5 units down.




24. $f(x) = -6x + 12$

a. zero: $0 = -6x + 12$: y-intercept = 12

$x = 2$

b. Plot the points $(2,0),(0,12)$.




25. $H(x) = -\frac{1}{2}x + 4$

a. zero: $0 = -\frac{1}{2}x + 4$: y-intercept = 4

$x = 8$

b. Plot the points $(8,0),(0,4)$.



Copyright © 2019 Pearson Education, Inc.


Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

26. $G(x)=\frac{1}{3}x-4$

a. zero: $0=\frac{1}{3}x-4$: y-intercept = -4

$x=12$

b. Plot the points $(12,0),(0,-4)$.



27.

x	y	Avg. rate of change $=\frac{\Delta y}{\Delta x}$
-2	4	
-1	1	$\frac{1-4}{-1-(-2)}=\frac{-3}{1}=-3$
0	-2	$\frac{-2-1}{0-(-1)}=\frac{-3}{1}=-3$
1	-5	$\frac{-5-(-2)}{1-0}=\frac{-3}{1}=-3$
2	-8	$\frac{-8-(-5)}{2-1}=\frac{-3}{1}=-3$

Since the average rate of change is constant at -3, this is a linear function with slope = -3. The y-intercept is $(0,-2)$, so the equation of the line is $y=-3x-2$.

28.

x	y	Avg. rate of change $=\frac{\Delta y}{\Delta x}$
-2	$\frac{1}{4}$	
-1	$\frac{1}{2}$	$\frac{\left(\frac{1}{2}-\frac{1}{4}\right)}{-1-(-2)}=\frac{\frac{1}{4}}{1}=\frac{1}{4}$
0	1	$\frac{\left(1-\frac{1}{2}\right)}{0-(-1)}=\frac{\frac{1}{2}}{1}=\frac{1}{2}$
1	2	
2	4	

Since the average rate of change is not constant, this is not a linear function.

29.

x	y	Avg. rate of change $=\frac{\Delta y}{\Delta x}$
-2	-8	
-1	-3	$\frac{-3-(-8)}{-1-(-2)}=\frac{5}{1}=5$
0	0	$\frac{0-(-3)}{0-(-1)}=\frac{3}{1}=3$
1	1	
2	0	

Since the average rate of change is not constant, this is not a linear function.

30.

x	y	Avg. rate of change $=\frac{\Delta y}{\Delta x}$
-2	-4	
-1	0	$\frac{0-(-4)}{-1-(-2)}=\frac{4}{1}=4$
0	4	$\frac{4-0}{0-(-1)}=\frac{4}{1}=4$
1	8	$\frac{8-4}{1-0}=\frac{4}{1}=4$
2	12	$\frac{12-8}{2-1}=\frac{4}{1}=4$

Since the average rate of change is constant at 4, this is a linear function with slope = 4. The y-intercept is $(0,4)$, so the equation of the line is $y=4x+4$.

31.

x	y	Avg. rate of change $=\frac{\Delta y}{\Delta x}$
-2	-26	
-1	-4	$\frac{-4-(-26)}{-1-(-2)}=\frac{22}{1}=22$
0	2	$\frac{2-(-4)}{0-(-1)}=\frac{6}{1}=6$
1	-2	
2	-10	

Since the average rate of change is not constant, this is not a linear function.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

32.

x	y	Avg. rate of change $= \dfrac{\Delta y}{\Delta x}$
-2	-4	
-1	-3.5	$\dfrac{-3.5-(-4)}{-1-(-2)} = \dfrac{0.5}{1} = 0.5$
0	-3	$\dfrac{-3-(-3.5)}{0-(-1)} = \dfrac{0.5}{1} = 0.5$
1	-2.5	$\dfrac{-2.5-(-3)}{1-0} = \dfrac{0.5}{1} = 0.5$
2	-2	$\dfrac{-2-(-2.5)}{2-1} = \dfrac{0.5}{1} = 0.5$

Since the average rate of change is constant at 0.5, this is a linear function with slope = 0.5. The y-intercept is $(0,-3)$, so the equation of the line is $y = 0.5x - 3$.

33.

x	y	Avg. rate of change $= \dfrac{\Delta y}{\Delta x}$
-2	8	
-1	8	$\dfrac{8-8}{-1-(-2)} = \dfrac{0}{1} = 0$
0	8	$\dfrac{8-8}{0-(-1)} = \dfrac{0}{1} = 0$
1	8	$\dfrac{8-8}{1-0} = \dfrac{0}{1} = 0$
2	8	$\dfrac{8-8}{2-1} = \dfrac{0}{1} = 0$

Since the average rate of change is constant at 0, this is a linear function with slope = 0. The y-intercept is $(0, 8)$, so the equation of the line is $y = 0x + 8$ or $y = 8$.

34.

x	y	Avg. rate of change $= \dfrac{\Delta y}{\Delta x}$
-2	0	
-1	1	$\dfrac{1-0}{-1-(-2)} = \dfrac{1}{1} = 1$
0	4	$\dfrac{4-1}{0-(-1)} = \dfrac{3}{1} = 3$
1	9	
2	16	

Since the average rate of change is not constant, this is not a linear function.

35. $f(x) = 4x - 1; \quad g(x) = -2x + 5$

a. $f(x) = 0$

$4x - 1 = 0$

$x = \dfrac{1}{4}$

b. $f(x) > 0$

$4x - 1 > 0$

$x > \dfrac{1}{4}$

The solution set is $\left\{x \middle| x > \dfrac{1}{4}\right\}$ or $\left(\dfrac{1}{4}, \infty\right)$.

c. $f(x) = g(x)$

$4x - 1 = -2x + 5$

$6x = 6$

$x = 1$

d. $f(x) \le g(x)$


$4x - 1 \le -2x + 5$

$6x \le 6$

$x \le 1$

The solution set is $\{x | x \le 1\}$ or $(-\infty, 1]$.

e.



36. $f(x) = 3x + 5; \quad g(x) = -2x + 15$

a. $f(x) = 0$

$3x + 5 = 0$

$x = -\dfrac{5}{3}$

b. $f(x) < 0$

$3x + 5 < 0$

$x < -\dfrac{5}{3}$

The solution set is $\left\{x \middle| x < -\dfrac{5}{3}\right\}$ or $\left(-\infty, -\dfrac{5}{3}\right)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

c. $f(x) = g(x)$

$3x + 5 = -2x + 15$

$5x = 10$

$x = 2$

d. $f(x) \geq g(x)$


$3x + 5 \geq -2x + 15$

$5x \geq 10$

$x \geq 2$

The solution set is $\{x | x \geq 2\}$ or $[2, \infty)$.

e.



37. a. The point (40, 50) is on the graph of $y = f(x)$, so the solution to $f(x) = 50$ is $x = 40$.

b. The point (88, 80) is on the graph of $y = f(x)$, so the solution to $f(x) = 80$ is $x = 88$.

c. The point $(-40, 0)$ is on the graph of $y = f(x)$, so the solution to $f(x) = 0$ is $x = -40$.

d. The y-coordinates of the graph of $y = f(x)$ are above 50 when the x-coordinates are larger than 40. Thus, the solution to $f(x) > 50$ is $\{x | x > 40\}$ or $(40, \infty)$.

e. The y-coordinates of the graph of $y = f(x)$ are below 80 when the x-coordinates are smaller than 88. Thus, the solution to $f(x) \leq 80$ is $\{x | x \leq 88\}$ or $(-\infty, 88]$.

f. The y-coordinates of the graph of $y = f(x)$ are between 0 and 80 when the x-coordinates are between -40 and 88. Thus, the solution to $0 < f(x) < 80$ is $\{x | -40 < x < 88\}$ or $(-40, 88)$.

38. a. The point (5, 20) is on the graph of $y = g(x)$, so the solution to $g(x) = 20$ is $x = 5$.

b. The point $(-15, 60)$ is on the graph of $y = g(x)$, so the solution to $g(x) = 60$ is $x = -15$.

c. The point (15, 0) is on the graph of $y = g(x)$, so the solution to $g(x) = 0$ is $x = 15$.

d. The y-coordinates of the graph of $y = g(x)$ are above 20 when the x-coordinates are smaller than 5. Thus, the solution to $g(x) > 20$ is $\{x | x < 5\}$ or $(-\infty, 5)$.

e. The y-coordinates of the graph of $y = f(x)$ are below 60 when the x-coordinates are larger than -15. Thus, the solution to $g(x) \leq 60$ is $\{x | x \geq -15\}$ or $[-15, \infty)$.

f. The y-coordinates of the graph of $y = f(x)$ are between 0 and 60 when the x-coordinates are between -15 and 15. Thus, the solution to $0 < f(x) < 60$ is $\{x | -15 < x < 15\}$ or $(-15, 15)$.

39. a. $f(x) = g(x)$ when their graphs intersect. Thus, $x = -4$.

b. $f(x) \leq g(x)$ when the graph of f is above the graph of g. Thus, the solution is $\{x | x < -4\}$ or $(-\infty, -4)$.

40. a. $f(x) = g(x)$ when their graphs intersect. Thus, $x = 2$.

b. $f(x) \leq g(x)$ when the graph of f is below or intersects the graph of g. Thus, the solution is $\{x | x \leq 2\}$ or $(-\infty, 2]$.

41. a. $f(x) = g(x)$ when their graphs intersect. Thus, $x = -6$.

b. $g(x) \leq f(x) < h(x)$ when the graph of f is above or intersects the graph of g and below the graph of h. Thus, the solution is $\{x | -6 \leq x < 5\}$ or $[-6, 5)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

42. a. $f(x) = g(x)$ when their graphs intersect. Thus, $x = 7$.

b. $g(x) \le f(x) < h(x)$ when the graph of f is above or intersects the graph of g and below the graph of h. Thus, the solution is $\{x | -4 \le x < 7\}$ or $[-4, 7)$.

43. $C(x) = 2.5x + 85$

a. $C(40) = 2.5(40) + 85 = \$185$.

b. Solve $C(x) = 2.5x + 85 = 245$

$$2.5x + 85 = 245$$
$$2.5x = 100$$
$$x = \frac{160}{2.5} = 64 \text{ miles}$$

c. Solve $C(x) = 0.35x + 45 \le 150$

$$2.5x + 85 \le 150$$
$$2.5x \le 105$$
$$x \le \frac{65}{2.5} = 26 \text{ miles}$$

d. The number of mile towed cannot be negative, so the implied domain for C is $\{x | x \ge 0\}$ or $[0, \infty)$.

e. The cost of being towed increases \$2.50 for each mile, or there is a charge of \$2.50 per mile towed in addition to a fixed charge of \$85.

f. It costs \$85 for towing 0 miles, or there is a fixed charge of \$85 for towing in addition to a charge that depends on mileage.

44. $C(x) = 0.07x + 24.99$

a. $C(50) = 0.07(50) + 24.99 = \$28.49$.

b. Solve $C(x) = 0.07x + 24.99 = 31.85$

$$0.07x + 24.99 = 31.85$$
$$0.07x = 6.86$$
$$x = \frac{6.86}{0.07} = 98 \text{ minutes}$$

c. Solve $C(x) = 0.07x + 24.99 \le 36$

$$0.07x + 24.99 \le 36$$
$$0.07x \le 11.01$$
$$x \le \frac{11.01}{0.07} \approx 157 \text{ minutes}$$

d. The number of minutes cannot be negative, so $x \ge 0$. If there are 30 days in the month, then the number of minutes can be at most $30 \cdot 24 \cdot 60 = 43,200$. Thus, the implied domain for C is $\{x | 0 \le x \le 43,200\}$ or $[0, 43200]$.

e. The monthly cost of the plan increases \$0.07 for each minute used, or there is a charge of \$0.07 per minute to use the phone in addition to a fixed charge of \$24.99.

f. It costs \$24.99 per month for the plan if 0 minutes are used, or there is a fixed charge of \$24.99 per month for the plan in addition to a charge that depends on the number of minutes used.

45. $S(p) = -600 + 50p;\ D(p) = 1200 - 25p$

a. Solve $S(p) = D(p)$.

$$-600 + 50p = 1200 - 25p$$
$$75p = 1800$$
$$p = \frac{1800}{75} = 24$$
$$S(24) = -600 + 50(24) = 600$$

Thus, the equilibrium price is \$24, and the equilibrium quantity is 600 T-shirts.

b. Solve $D(p) > S(p)$.

$$1200 - 25p > -600 + 50p$$
$$1800 > 75p$$
$$\frac{1800}{75} > p$$
$$24 > p$$

The demand will exceed supply when the price is less than \$24 (but still greater than \$0). That is, $\$0 < p < \$24$.

c. The price will eventually be increased.

46. $S(p) = -2000 + 3000p;\ D(p) = 10000 - 1000p$

a. Solve $S(p) = D(p)$.

$$-2000 + 3000p = 10000 - 1000p$$
$$4000p = 12000$$
$$p = \frac{12000}{4000} = 3$$
$$S(3) = -2000 + 3000(3) = 7000$$

Thus, the equilibrium price is \$3, and the equilibrium quantity is 7000 hot dogs.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

b. Solve $D(p) < S(p)$.

$$10000 - 1000p < -2000 + 3000p$$
$$12000 < 4000p$$
$$\frac{12000}{4000} < p$$
$$3 < p$$

The demand will be less than the supply when the price is greater than \$3.

c. The price will eventually be decreased.

47. a. We are told that the tax function T is for adjusted gross incomes x between \$9,325 and \$37,950, inclusive. Thus, the domain is $\{x \mid 9{,}325 \le x \le 37{,}950\}$ or $[9325,\ 37950]$.

b. $T(20{,}000) = 0.15(20{,}000 - 9325) + 932.50$
$= 2533.75$

If a single filer's adjusted gross income is \$20,000, then his or her tax bill will be \$2533.75.

c. The independent variable is adjusted gross income, x. The dependent variable is the tax bill, T.


d. Evaluate T at $x = 9325,\ 20000,$ and 37950.

$T(9325) = 0.15(9325 - 9325) + 932.50$
$= 932.50$

$T(20{,}000) = 0.15(20{,}000 - 9325) + 932.50$
$= 2533.75$

$T(37{,}950) = 0.15(37950 - 9325) + 932.50$
$= 5226.25$

Thus, the points $(9325, 932.50)$, $(20000, 2533.75)$, and $(37950, 5226.25)$ are on the graph.



e. We must solve $T(x) = 3673.75$.

$0.15(x - 9325) + 932.50 = 3673.75$
$0.15x - 1398.75 + 932.50 = 3673.75$
$0.15x - 466.25 = 3673.75$
$0.15x = 4140$
$x = 27600$

A single filer with an adjusted gross income of \$27,600 will have a tax bill of \$3673.75.

f. For each additional dollar of taxable income between \$9325 and \$37,950, the tax bill of a single person in 2013 increased by \$0.15.

48. a. The independent variable is payroll, p. The payroll tax only applies if the payroll exceeds \$189 million. Thus, the domain of T is $\{p \mid p > 189\}$ or $(189,\ \infty)$.

b. $T(243.8) = 0.5(243.8 - 189) = 27.4$

The luxury tax for the New York Yankees was \$27.4 million.


c. Evaluate T at $p = 189$, 243.8, and 300 million.

$T(189) = 0.5(189 - 189) = 0$ million

$T(243.8) = 0.5(243.8 - 189)$
$= 27.4$ million

$T(300) = 0.5(300 - 189) = 55.5$ million

Thus, the points $(189 \text{ million},\ 0 \text{ million})$, $(243.8 \text{ million},\ 27.4 \text{ million})$, and $(300 \text{ million},\ 55.5 \text{ million})$ are on the graph.



d. We must solve $T(p) = 31.8$.

$0.5(p - 189) = 31.8$
$0.5p - 94.5 = 31.8$
$0.5p = 126.3$
$p = 252.6$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

If the luxury tax is \$31.8 million, then the payroll of the team is \$252.6 million.

e. For each additional million dollars of payroll in excess of \$189 million in 2016, the luxury tax of a team increased by \$0.5 million.

49. $R(x) = 8x;\ \ C(x) = 4.5x + 17{,}500$

a. Solve $R(x) = C(x)$.

$8x = 4.5x + 17{,}500$

$3.5x = 17{,}500$

$x = 5000$

The break-even point occurs when the company sells 5000 units.

b. Solve $R(x) > C(x)$

$8x > 4.5x + 17{,}500$

$3.5x > 17{,}500$

$x > 5000$

The company makes a profit if it sells more than 5000 units.

50. $R(x) = 12x;\ \ C(x) = 10x + 15{,}000$

a. Solve $R(x) = C(x)$

$12x = 10x + 15{,}000$

$2x = 15{,}000$

$x = 7500$

The break-even point occurs when the company sells 7500 units.

b. Solve $R(x) > C(x)$

$12x > 10x + 15{,}000$

$2x > 15{,}000$

$x > 7500$

The company makes a profit if it sells more than 7500 units.

51. a. Consider the data points (x, y), where x = the age in years of the computer and y = the value in dollars of the computer. So we have the points $(0, 3000)$ and $(3, 0)$. The slope formula yields:


$$m = \frac{\Delta y}{\Delta x} = \frac{0 - 3000}{3 - 0} = \frac{-3000}{3} = -1000$$

The y-intercept is $(0, 3000)$, so $b = 3000$.

Therefore, the linear function is $V(x) = mx + b = -1000x + 3000$.

b. The age of the computer cannot be negative, and the book value of the computer will be \$0 after 3 years. Thus, the implied domain for V is $\{x \mid 0 \le x \le 3\}$ or $[0, 3]$.

c. The graph of $V(x) = -1000x + 3000$



d. $V(2) = -1000(2) + 3000 = 1000$

The computer's book value after 2 years will be \$1000.

e. Solve $V(x) = 2000$

$-1000x + 3000 = 2000$

$-1000x = -1000$

$x = 1$

The computer will have a book value of \$2000 after 1 year.

52. a. Consider the data points (x, y), where x = the age in years of the machine and y = the value in dollars of the machine. So we have the points $(0, 120000)$ and $(10, 0)$. The slope formula yields:


$$m = \frac{\Delta y}{\Delta x} = \frac{0 - 120000}{10 - 0} = \frac{-120000}{10} = -12000$$

The y-intercept is $(0, 120000)$, so $b = 120{,}000$.

Therefore, the linear function is $V(x) = mx + b = -12{,}000x + 120{,}000$.

b. The age of the machine cannot be negative, and the book value of the machine will be \$0 after 10 years. Thus, the implied domain for V is $\{x \mid 0 \le x \le 10\}$ or $[0, 10]$.

c. The graph of $V(x) = -12{,}000x + 120{,}000$



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

d. $V(4) = -12000(4) + 120000 = 72000$

The machine's value after 4 years is given by \$72,000.


e. Solve $V(x) = 72000$.

$$-12000x + 120000 = 72000$$
$$-12000x = -48000$$
$$x = 4$$

The machine will be worth \$72,000 after 4 years.

53. a. Let x = the number of bicycles manufactured. We can use the cost function $C(x) = mx + b$, with $m = 90$ and $b = 1800$. Therefore $C(x) = 90x + 1800$

b. The graph of $C(x) = 90x + 1800$



c. The cost of manufacturing 14 bicycles is given by $C(14) = 90(14) + 1800 = \$3060$.

d. Solve $C(x) = 90x + 1800 = 3780$

$$90x + 1800 = 3780$$
$$90x = 1980$$
$$x = 22$$

So 22 bicycles could be manufactured for \$3780.


54. a. The new daily fixed cost is

$$1800 + \frac{100}{20} = \$1805$$

b. Let x = the number of bicycles manufactured. We can use the cost function $C(x) = mx + b$, with $m = 90$ and $b = 1805$.

Therefore $C(x) = 90x + 1805$

c. The graph of $C(x) = 90x + 1805$



d. The cost of manufacturing 14 bicycles is given by $C(14) = 90(14) + 1805 = \$3065$.

e. Solve $C(x) = 90x + 1805 = 3780$

$$90x + 1805 = 3780$$
$$90x = 1975$$
$$x \approx 21.94$$

So approximately 21 bicycles could be manufactured for \$3780.

55. a. Let x = number of miles driven, and let C = cost in dollars. Total cost = (cost per mile)(number of miles) + fixed cost

$C(x) = 0.89x + 39.95$

b. $C(110) = (0.89)(110) + 39.95 = \$137.85$

$C(230) = (0.89)(230) + 39.95 = \$244.65$

56. a. Let x = number of megabytes used, and let C = cost in dollars. Total cost = (cost per megabyte)(number of megabytes over 200) + fixed cost:

$$C(x) = 0.25(x - 200) + 40$$
$$= 0.25x - 50 + 40$$
$$= 0.25x - 10, x > 200$$


b. $C(265) = (0.25)(265) - 10 = \$56.25$

$C(300) = (0.25)(300) - 10 = \$65$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

57. a.



b.

m	n	Avg. rate of change $= \dfrac{\Delta n}{\Delta m}$
4	30	
16	120	$\dfrac{120-30}{16-4}=\dfrac{90}{12}=\dfrac{15}{2}$
64	480	$\dfrac{480-120}{64-16}=\dfrac{360}{48}=\dfrac{15}{2}$
128	960	$\dfrac{960-480}{128-64}=\dfrac{480}{64}=\dfrac{15}{2}$

Since each input (memory) corresponds to a single output (recording time), we know that recording time is a function of memory. Also, because the average rate of change is constant at 7.5 minutes per gigabyte, the function is linear.

c. From part (b), we know $slope = 7.5$. Using $(m_1, n_1) = (4, 30)$, we get the equation:

$$t - t_1 = s(m - m_1)$$
$$t - 30 = 7.5(m - 4)$$
$$t - 30 = 7.5m - 30$$
$$t = 7.5m$$


Using function notation, we have $t(m) = 7.5m$.

d. The memory cannot be negative, so $m \geq 0$. Likewise, the time cannot be negative, so, $t(m) \geq 0$.

$$7.5m \geq 0$$
$$m \geq 0$$


Thus, the implied domain for n(m) is $\{m \mid m \geq 0\}$ or $[0, \infty)$.

e.



f. If memory increases by 1 GB, then the number of songs increases by 218.75.

58. a.



b.

s	h	Avg. rate of change $= \dfrac{\Delta h}{\Delta s}$
20	0	
15	3	$\dfrac{3-0}{15-20}=\dfrac{3}{-5}=-0.6$
10	6	$\dfrac{6-3}{10-15}=\dfrac{3}{-5}=-0.6$
5	9	$\dfrac{9-6}{5-10}=\dfrac{3}{-5}=-0.6$

Since each input (soda) corresponds to a single output (hot dogs), we know that number of hot dogs purchased is a function of number of sodas purchased. Also, because the average rate of change is constant at -0.6 hot dogs per soda, the function is linear.

c. From part (b), we know $m = -0.6$. Using $(s_1, h_1) = (20, 0)$, we get the equation:

$$h - h_1 = m(s - s_1)$$
$$h - 0 = -0.6(s - 20)$$
$$h = -0.6s + 12$$

Using function notation, we have $h(s) = -0.6s + 12$.

Copyright © 2019 Pearson Education, Inc.


Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

d. The number of sodas cannot be negative, so $s \ge 0$. Likewise, the number of hot dogs cannot be negative, so, $h(s) \ge 0$.

$$-0.6s + 12 \ge 0$$
$$-0.6s \ge -12$$
$$s \le 20$$

Thus, the implied domain for h(s) is $\{s \mid 0 \le s \le 20\}$ or $[0, 20]$.

e.



f. If the number of hot dogs purchased increases by \$1, then the number of sodas purchased decreases by 0.6.

g. s-intercept: If 0 hot dogs are purchased, then 20 sodas can be purchased.
h-intercept: If 0 sodas are purchased, then 12 hot dogs may be purchased.

59. The graph shown has a positive slope and a positive y-intercept. Therefore, the function from (d) and (e) might have the graph shown.

60. The graph shown has a negative slope and a positive y-intercept. Therefore, the function from (b) and (e) might have the graph shown.

61. A linear function $f(x) = mx + b$ will be odd provided $f(-x) = -f(x)$.

That is, provided $m(-x) + b = -(mx + b)$.

$$-mx + b = -mx - b$$
$$b = -b$$
$$2b = 0$$
$$b = 0$$

So a linear function $f(x) = mx + b$ will be odd provided $b = 0$.

A linear function $f(x) = mx + b$ will be even provided $f(-x) = f(x)$.

That is, provided $m(-x) + b = mx + b$.


$$-mx + b = mx + b$$
$$-mxb = mx$$
$$0 = 2mx$$
$$m = 0$$

So, yes, a linear function $f(x) = mx + b$ cab be even provided $m = 0$.

62. If you solve the linear function $f(x) = mx + b$ for 0 you are actually finding the x-intercept. Therefore using x-intercept of the graph of $f(x) = mx + b$ would be same x-value as solving $mx + b > 0$ for x. Then the appropriate interval could be determined

63.
$$x^2 - 4x + y^2 + 10y - 7 = 0$$
$$(x^2 - 4x + 4) + (y^2 + 10y + 25) = 7 + 4 + 25$$
$$(x - 2)^2 + (y + 5)^2 = 6^2$$

Center: (2, -5); Radius = 6



64.
$$f(x) = \frac{2x + B}{x - 3}$$
$$f(5) = 8 = \frac{2(5) + B}{5 - 3}$$
$$8 = \frac{10 + B}{2}$$
$$16 = 10 + B$$
$$B = 6$$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


65. $\dfrac{f(3)-f(1)}{3-1}$

$= \dfrac{12-(-2)}{2}$

$= \dfrac{14}{2}$


$= 7$

66.



Section 2.2

1.



No, the relation is not a function because an input, 1, corresponds to two different outputs, 5 and 12.

2. Let $(x_1, y_1) = (1, 4)$ and $(x_2, y_2) = (3, 8)$.

$m = \dfrac{y_2 - y_1}{x_2 - x_1} = \dfrac{8-4}{3-1} = \dfrac{4}{2} = 2$

$y - y_1 = m(x - x_1)$

$y - 4 = 2(x - 1)$

$y - 4 = 2x - 2$

$y = 2x + 2$

3. scatter diagram

4. decrease; 0.008

5. Linear relation, $m > 0$

6. Nonlinear relation


7. Linear relation, $m < 0$

8. Linear relation, $m > 0$

9. Nonlinear relation

10. Nonlinear relation

11. a.



b. Answers will vary. We select (4, 6) and (8, 14). The slope of the line containing these points is:

$m = \dfrac{14-6}{8-4} = \dfrac{8}{4} = 2$

The equation of the line is:


$y - y_1 = m(x - x_1)$

$y - 6 = 2(x - 4)$

$y - 6 = 2x - 8$

$y = 2x - 2$


c.




d. Using the LINear REGression program, the line of best fit is:

$y = 2.0357x - 2.3571$

e.



12. a.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

b. Answers will vary. We select (5, 2) and (11, 9). The slope of the line containing these points is: $m = \frac{9-2}{11-5} = \frac{7}{6}$

The equation of the line is:


$y - y_1 = m(x - x_1)$

$y - 2 = \frac{7}{6}(x - 5)$

$y - 2 = \frac{7}{6}x - \frac{35}{6}$

$y = \frac{7}{6}x - \frac{23}{6}$


c.




d. Using the LINear REGression program, the line of best fit is:

$y = 1.1286x - 3.8619$

e.



13. a.



b. Answers will vary. We select (–2,–4) and (2, 5). The slope of the line containing these points is: $m = \frac{5-(-4)}{2-(-2)} = \frac{9}{4}$.

The equation of the line is:


$y - y_1 = m(x - x_1)$

$y - (-4) = \frac{9}{4}(x - (-2))$

$y + 4 = \frac{9}{4}x + \frac{9}{2}$


$y = \frac{9}{4}x + \frac{1}{2}$

c.




e. Using the LINear REGression program, the line of best fit is:

$y = 2.2x + 1.2$



14. a.



b. Answers will vary. We select (–2, 7) and (2, 0). The slope of the line containing these points is: $m = \frac{0-7}{2-(-2)} = \frac{-7}{4} = -\frac{7}{4}$.

The equation of the line is:


$y - y_1 = m(x - x_1)$

$y - 7 = -\frac{7}{4}(x - (-2))$

$y - 7 = -\frac{7}{4}x - \frac{7}{2}$

$y = -\frac{7}{4}x + \frac{7}{2}$

c.




d. Using the LINear REGression program, the line of best fit is:

$y = -1.8x + 3.6$


Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

e.



15. a.




b. Answers will vary. We select (–20,100) and (–10,140). The slope of the line containing these points is:

$$m = \frac{140-100}{-10-(-20)} = \frac{40}{10} = 4$$

The equation of the line is:

$$y - y_1 = m(x - x_1)$$
$$y - 100 = 4(x - (-20))$$
$$y - 100 = 4x + 80$$
$$y = 4x + 180$$


c.




d. Using the LINear REGression program, the line of best fit is:

$$y = 3.8613x + 180.2920$$

e.



16. a.




b. Selection of points will vary. We select (–30, 10) and (–14, 18). The slope of the line containing these points is:

$$m = \frac{18-10}{-14-(-30)} = \frac{8}{16} = \frac{1}{2}$$

The equation of the line is:

$$y - y_1 = m(x - x_1)$$
$$y - 10 = \frac{1}{2}(x - (-30))$$
$$y - 10 = \frac{1}{2}x + 15$$
$$y = \frac{1}{2}x + 25$$


c.




d. Using the LINear REGression program, the line of best fit is:

$$y = 0.4421x + 23.4559$$

e.



17. a.



b. Linear.


c. Answers will vary. We will use the points $(39.52,\ 210)$ and $(66.45,\ 280)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$$m = \frac{280-210}{66.45-39.52} = \frac{70}{26.93} \approx 2.5993316$$
$$y - 210 = 2.5993316(x - 39.52)$$
$$y - 210 = 2.5993316x - 102.7255848$$
$$y = 2.599x + 107.274$$

d.




e. $x = 62.3: \; y = 2.599(62.3) + 107.274 \approx 269$

We predict that a candy bar weighing 62.3 grams will contain 269 calories.

f. If the weight of a candy bar is increased by one gram, then the number of calories will increase by 2.599.

18. a.




b. Linear with positive slope.

c. Answers will vary. We will use the points $(200,\ 2.5)$ and $(500,\ 5.8)$.

$$m = \frac{5.8-2.5}{500-200} = \frac{3.3}{300} = 0.011$$
$$L - L_1 = m(w - w_1)$$
$$L - 2.5 = 0.011(w - 200)$$
$$L - 2.5 = 0.011w - 2.2$$
$$L = 0.011w + 0.3$$

d.




e. $L(450) = 0.011(450) + 0.3 = 5.25$

We predict that the approximately length of a 450 yard wide tornado is 5.25 miles.

f. For each 1-yard increase in the width of a tornado, the length of the tornado increases by 0.011 mile, on average.

19. a. The independent variable is the number of hours spent playing video games and cumulative grade-point average is the dependent variable because we are using number of hours playing video games to predict (or explain) cumulative grade-point average.

b.



c. Using the LINear REGression program, the line of best fit is: $G(h) = -0.0942h + 3.2763$

d. If the number of hours playing video games in a week increases by 1 hour, the cumulative grade-point average decreases 0.09, on average.

e. $G(8) = -0.0942(8) + 3.2763 = 2.52$

We predict a grade-point average of approximately 2.52 for a student who plays 8 hours of video games each week.


f.
$$2.40 = -0.0942(h) + 3.2763$$
$$2.40 - 3.2763 = -0.0942h$$
$$-0.8763 = -0.0942h$$
$$9.3 = h$$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

A student who has a grade-point average of 2.40 will have played approximately 9.3 hours of video games.

20. a.



b. Using the LINear REGression program, the line of best fit is:
$w(p) = -1.1857p + 1231.8279$

c. For each 10-millibar increase in the atmospheric pressure, the wind speed of the tropical system decreases by 1.1857 knots, on average.

d. $w(990) = -1.1857(990) + 1231.8279 \approx 58$ knots


e. To find the pressure, we solve the following equation:

$$85 = -1.1857p + 1231.8279$$
$$-1146.8279 = -1.1857p$$
$$967 \approx p$$

A hurricane with a wind speed of 85 knots would have a pressure of approximately 967 millibars.

21. a. This relation does not represent a function since the values of the input variable s are repeated.

b.



c. Using the LINear REGression program, the line of best fit is: $d = 3.3641s + 51.8233$

d. For each 1-mph increase in the speed off bat, the homerun distance increases by 3.3641 feet, on average.

e. $d(s) = 3.3641s + 51.8233$


f. Since the speed off bat must be greater than 0 the domain is $\{s \mid s > 0\}$.

g. $d(103) = 3.3641(103) + 51.8233 \approx 398$ ft

A hurricane with a wind speed of 85 knots would have a pressure of approximately 967 millibars.

22. a. The relation is a function because none of the invariables are repeated.

b.



c. Using the LINear REGression program, the line of best fit is: $m = -0.2277n + 15.9370$.

d. If Internet ad spending increases by 1%, magazine ad spending goes down by about 0.2277%, on average.

e. $m(n) = -0.2277n + 15.9370$

f. Domain: $\{n \mid 0 < n \le 70.0\}$

Note that the m-intercept is roughly 15.9 and that the percent of Internet sales cannot be negative.


g. $D(28) = -0.2277(26.0) + 15.9370 \approx 10.0$

Percent of magazine sales is about 10.0%.

Copyright © 2019 Pearson Education, Inc.


Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

23.



The data do not follow a linear pattern so it would not make sense to find the line of best fit.

24. Using the ordered pairs $(1, 5)$ and $(3, 8)$, the line of best fit is $y = 1.5x + 3.5$.



The correlation coefficient is $r = 1$. This is reasonable because two points determine a line.

25. A correlation coefficient of 0 implies that the data do not have a linear relationship.

26. The y-intercept would be the calories of a candy bar with weight 0 which would not be meaningful in this problem.

27. $G(0) = -0.0942(0) + 3.2763 = 3.2763$. The approximate grade-point average of a student who plays 0 hours of video games per week would be 3.28.

28. $m = \dfrac{-3-5}{3-(-1)} = \dfrac{-8}{4} = -2$

$y - y_1 = m(x - x_1)$

$y - 5 = -2(x + 1)$

$y - 5 = -2x - 2$

$y = -2x + 3$ or

$2x + y = 3$

29. The domain would be all real numbers except those that make the denominator zero.

$x^2 - 25 = 0$

$x^2 = 25 \rightarrow x = \pm 5$

So the domain is: $\{x \mid x \neq 5, -5\}$

30. $f(x) = 5x - 8$ and $g(x) = x^2 - 3x + 4$

$(g - f)(x) = (x^2 - 3x + 4) - (5x - 8)$

$= x^2 - 3x + 4 - 5x + 8$

$= x^2 - 8x + 12$

31. Since y is shifted to the left 3 units we would use $y = (x + 3)^2$. Since y is also shifted down 4 units,we would use $y = (x + 3)^2 - 4$.

Section 2.3

1. a. $x^2 - 5x - 6 = (x - 6)(x + 1)$

b. $2x^2 - x - 3 = (2x - 3)(x + 1)$

2. $\sqrt{8^2 - 4 \cdot 2 \cdot 3} = \sqrt{64 - 24}$

$= \sqrt{40} = \sqrt{4 \cdot 10} = 2\sqrt{10}$

3. $(x - 3)(3x + 5) = 0$

$x - 3 = 0$ or $3x + 5 = 0$

$x = 3$ $\qquad 3x = -5$

$x = -\dfrac{5}{3}$

The solution set is $\left\{-\dfrac{5}{3}, 3\right\}$.

4. add; $\left(\dfrac{1}{2} \cdot 6\right)^2 = 9$

5. If $f(4) = 10$, then the point (4, 10) is on the graph of *f*.

6. $f(-3) = (-3)^2 + 4(-3) + 3$

$= 9 - 12 + 3 = 0$

-3 is a zero of $f(x)$.

7. repeated; multiplicity 2

8. discriminant; negative

9. A quadratic functions can have either 0, 1 or 2 real zeros.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

10. $x = \dfrac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

11. False; the equation will have only two real solution but not necessarily negatives of one another.

12. b

13.
$$f(x) = 0$$
$$x^2 - 9x = 0$$
$$x(x-9) = 0$$
$$x = 0 \quad \text{or} \quad x - 9 = 0$$
$$x = 9$$

The zeros of $f(x) = x^2 - 9x$ are 0 and 9. The x-intercepts of the graph of f are 0 and 9.

14.
$$f(x) = 0$$
$$x^2 + 4x = 0$$
$$x(x+4) = 0$$
$$x = 0 \quad \text{or} \quad x + 4 = 0$$
$$x = -4$$

The zeros of $f(x) = x^2 + 4x$ are -4 and 0. The x-intercepts of the graph of f are -4 and 0.

15.
$$g(x) = 0$$
$$x^2 - 25 = 0$$
$$(x+5)(x-5) = 0$$
$$x + 5 = 0 \quad \text{or} \quad x - 5 = 0$$
$$x = -5 \qquad x = 5$$

The zeros of $g(x) = x^2 - 25$ are -5 and 5. The x-intercepts of the graph of g are -5 and 5.

16.
$$G(x) = 0$$
$$x^2 - 9 = 0$$
$$(x+3)(x-3) = 0$$
$$x + 3 = 0 \quad \text{or} \quad x - 3 = 0$$
$$x = -3 \qquad x = 3$$

The zeros of $G(x) = x^2 - 9$ are -3 and 3. The x-intercepts of the graph of G are -3 and 3.

17.
$$F(x) = 0$$
$$x^2 + x - 6 = 0$$
$$(x+3)(x-2) = 0$$
$$x + 3 = 0 \quad \text{or} \quad x - 2 = 0$$
$$x = -3 \qquad x = 2$$

The zeros of $F(x) = x^2 + x - 6$ are -3 and 2.
The x-intercepts of the graph of F are -3 and 2.

18.
$$H(x) = 0$$
$$x^2 + 7x + 6 = 0$$
$$(x+6)(x+1) = 0$$
$$x + 6 = 0 \quad \text{or} \quad x + 1 = 0$$
$$x = -6 \qquad x = -1$$

The zeros of $H(x) = x^2 + 7x + 6$ are -6 and -1.
The x-intercepts of the graph of H are -6 and -1.

19.
$$g(x) = 0$$
$$2x^2 - 5x - 3 = 0$$
$$(2x+1)(x-3) = 0$$
$$2x + 1 = 0 \quad \text{or} \quad x - 3 = 0$$
$$x = -\frac{1}{2} \qquad x = 3$$

The zeros of $g(x) = 2x^2 - 5x - 3$ are $-\dfrac{1}{2}$ and 3.
The x-intercepts of the graph of g are $-\dfrac{1}{2}$ and 3.

20.
$$f(x) = 0$$
$$3x^2 + 5x + 2 = 0$$
$$(3x+2)(x+1) = 0$$
$$3x + 2 = 0 \quad \text{or} \quad x + 1 = 0$$
$$x = -\frac{2}{3} \qquad x = -1$$

The zeros of $f(x) = 3x^2 + 5x + 2$ are -1 and $-\dfrac{2}{3}$. The x-intercepts of the graph of f are -1 and $-\dfrac{2}{3}$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

21. $P(x)=0$

$3x^2-48=0$

$3(x^2-16)=0$

$3(x+4)(x-4)=0$

$t+4=0$ or $t-4=0$

$t=-4$ $\quad$ $t=4$

The zeros of $P(x)=3x^2-48$ are -4 and 4.

The x-intercepts of the graph of P are -4 and 4.

22. $H(x)=0$

$2x^2-50=0$

$2(x^2-25)=0$

$2(x+5)(x-5)=0$

$y+5=0$ or $y-5=0$

$y=-5$ $\quad$ $y=5$

The zeros of $H(x)=2x^2-50$ are -5 and 5.

The x-intercepts of the graph of H are -5 and 5.

23. $g(x)=0$

$x(x+8)+12=0$

$x^2+8x+12=0$

$(x+6)(x+2)=0$

$x=-6$ or $x=-2$

The zeros of $g(x)=x(x+8)+12$ are -6 and -2.

The x-intercepts of the graph of g are -6 and -2.

24. $f(x)=0$

$x(x-4)-12=0$

$x^2-4x-12=0$

$(x-6)(x+2)=0$

$x=-2$ or $x=6$

The zeros of $f(x)=x(x-4)-12$ are -2 and 6.

The x-intercepts of the graph of f are -2 and 6.

25. $G(x)=0$

$4x^2+9-12x=0$

$4x^2-12x+9=0$

$(2x-3)(2x-3)=0$

$2x-3=0$ or $2x-3=0$

$x=\frac{3}{2}$ $\quad$ $x=\frac{3}{2}$

The only zero of $G(x)=4x^2+9-12x$ is $\frac{3}{2}$.

The only x-intercept of the graph of G is $\frac{3}{2}$.

26. $F(x)=0$

$25x^2+16-40x=0$

$25x^2-40x+16=0$

$(5x-4)(5x-4)=0$

$5x-4=0$ or $5x-4=0$

$x=\frac{4}{5}$ $\quad$ $x=\frac{4}{5}$

The only zero of $F(x)=25x^2+16-40x$ is $\frac{4}{5}$.

The only x-intercept of the graph of F is $\frac{4}{5}$.

27. $f(x)=0$

$x^2-8=0$

$x^2=8$

$x=\pm\sqrt{8}=\pm 2\sqrt{2}$

The zeros of $f(x)=x^2-8$ are $-2\sqrt{2}$ and $2\sqrt{2}$.

The x-intercepts of the graph of f are $-2\sqrt{2}$ and $2\sqrt{2}$.

28. $g(x)=0$

$x^2-18=0$

$x^2=18$

$x=\pm\sqrt{18}=\pm 3\sqrt{3}$

The zeros of $g(x)=x^2-18$ are $-3\sqrt{3}$ and $3\sqrt{3}$. The x-intercepts of the graph of g are $-3\sqrt{3}$ and $3\sqrt{3}$.

29. $g(x)=0$

$(x-1)^2-4=0$

$(x-1)^2=4$

$x-1=\pm\sqrt{4}$

$x-1=\pm 2$

$x-1=2$ or $x-1=-2$

$x=3$ $\quad$ $x=-1$

The zeros of $g(x)=(x-1)^2-4$ are -1 and 3.

The x-intercepts of the graph of g are -1 and 3.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

30. $G(x)=0$

$(x+2)^2-1=0$

$(x+2)^2=1$

$x+2=\pm\sqrt{1}$

$x+2=\pm 1$

$x+2=1$ or $x+2=-1$

$x=-1$ $\quad$ $x=-3$

The zeros of $G(x)=(x+2)^2-1$ are -3 and -1.

The x-intercepts of the graph of G are -3 and -1.

31. $F(x)=0$

$(2x+3)^2-32=0$

$(2x+3)^2=32$

$2x+3=\pm\sqrt{32}$

$2x+3=\pm 4\sqrt{2}$

$2x=-3\pm 4\sqrt{2}$

$x=\dfrac{-3\pm 4\sqrt{2}}{2}$

The zeros of $F(x)=(2x+3)^2-32$ are $\dfrac{-3+4\sqrt{2}}{2}$ and $\dfrac{-3-4\sqrt{2}}{2}$. The x-intercepts of the graph of F are $\dfrac{-3+4\sqrt{2}}{2}$ and $\dfrac{-3-4\sqrt{2}}{2}$.

32. $F(x)=0$

$(3x-2)^2-75=0$

$(3x-2)^2=75$

$3x-2=\pm\sqrt{75}$

$3x-2=\pm 5\sqrt{3}$

$3x=2\pm 5\sqrt{3}$

$x=\dfrac{2\pm 5\sqrt{3}}{3}$

The zeros of $G(x)=(3x-2)^2-75$ are $\dfrac{2+5\sqrt{3}}{3}$ and $\dfrac{2-5\sqrt{3}}{3}$. The x-intercepts of the graph of G are $\dfrac{2-5\sqrt{3}}{3}$ and $\dfrac{2+5\sqrt{3}}{3}$.

33. $f(x)=0$

$x^2+4x-8=0$

$x^2+4x=8$

$x^2+4x+4=8+4$

$(x+2)^2=12$

$x+2=\pm\sqrt{12}$

$x+2=\pm 2\sqrt{3}$

$x=-2\pm 2\sqrt{3}$

$x=-2+2\sqrt{3}$ or $x=-2-2\sqrt{3}$

The zeros of $f(x)=x^2+4x-8$ are $-2+2\sqrt{3}$ and $-2-2\sqrt{3}$. The x-intercepts of the graph of f are $-2+2\sqrt{3}$ and $-2-2\sqrt{3}$.

34. $f(x)=0$

$x^2-6x-9=0$

$x^2-6x+9=9+9$

$(x-3)^2=18$

$x-3=\pm\sqrt{18}$

$x=3\pm 3\sqrt{2}$

The zeros of $f(x)=x^2-6x-9$ are $3-3\sqrt{2}$ and $3+3\sqrt{2}$. The x-intercepts of the graph of f are $3-3\sqrt{2}$ and $3+3\sqrt{2}$.

35. $g(x)=0$

$x^2-\dfrac{1}{2}x-\dfrac{3}{16}=0$

$x^2-\dfrac{1}{2}x=\dfrac{3}{16}$

$x^2-\dfrac{1}{2}x+\dfrac{1}{16}=\dfrac{3}{16}+\dfrac{1}{16}$

$\left(x-\dfrac{1}{4}\right)^2=\dfrac{1}{4}$

$x-\dfrac{1}{4}=\pm\sqrt{\dfrac{1}{4}}=\pm\dfrac{1}{2}$

$x=\dfrac{1}{4}\pm\dfrac{1}{2}$

$x=\dfrac{3}{4}$ or $x=-\dfrac{1}{4}$

The zeros of $g(x)=x^2-\dfrac{1}{2}x-\dfrac{3}{16}$ are $-\dfrac{1}{4}$ and $\dfrac{3}{4}$.

The x-intercepts of the graph of g are $-\dfrac{1}{4}$ and $\dfrac{3}{4}$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

36. $g(x)=0$

$$x^2+\frac{2}{3}x-\frac{1}{3}=0$$
$$x^2+\frac{2}{3}x=\frac{1}{3}$$
$$x^2+\frac{2}{3}x+\frac{1}{9}=\frac{1}{3}+\frac{1}{9}$$
$$\left(x+\frac{1}{3}\right)^2=\frac{4}{9}$$
$$x+\frac{1}{3}=\pm\sqrt{\frac{4}{9}}=\pm\frac{2}{3}$$
$$x=-\frac{1}{3}\pm\frac{2}{3}$$
$$x=\frac{1}{3} \text{ or } x=-1$$

The zeros of $g(x)=x^2+\frac{2}{3}x-\frac{1}{3}$ are -1 and $\frac{1}{3}$.

The x-intercepts of the graph of g are -1 and $\frac{1}{3}$.

37. $F(x)=0$

$$3x^2+x-\frac{1}{2}=0$$
$$x^2+\frac{1}{3}x-\frac{1}{6}=0$$
$$x^2+\frac{1}{3}x=\frac{1}{6}$$
$$x^2+\frac{1}{3}x+\frac{1}{36}=\frac{1}{6}+\frac{1}{36}$$
$$\left(x+\frac{1}{6}\right)^2=\frac{7}{36}$$
$$x+\frac{1}{6}=\pm\sqrt{\frac{7}{36}}=\pm\frac{\sqrt{7}}{6}$$
$$x=\frac{-1\pm\sqrt{7}}{6}$$

The zeros of $F(x)=3x^2+x-\frac{1}{2}$ are $\frac{-1-\sqrt{7}}{6}$ and $\frac{-1+\sqrt{7}}{6}$. The x-intercepts of the graph of F are $\frac{-1-\sqrt{7}}{6}$ and $\frac{-1+\sqrt{7}}{6}$.

38. $G(x)=0$

$$2x^2-3x-1=0$$
$$x^2-\frac{3}{2}x-\frac{1}{2}=0$$
$$x^2-\frac{3}{2}x=\frac{1}{2}$$
$$x^2-\frac{3}{2}x+\frac{9}{16}=\frac{1}{2}+\frac{9}{16}$$
$$\left(x-\frac{3}{4}\right)^2=\frac{17}{16}$$
$$x-\frac{3}{4}=\pm\sqrt{\frac{17}{16}}=\pm\frac{\sqrt{17}}{4}$$
$$x=\frac{3\pm\sqrt{17}}{4}$$

The zeros of $G(x)=2x^2-3x-1$ are $\frac{3-\sqrt{17}}{4}$ and $\frac{3+\sqrt{17}}{4}$. The x-intercepts of the graph of G are $\frac{3-\sqrt{17}}{4}$ and $\frac{3+\sqrt{17}}{4}$.

39. $f(x)=0$

$$x^2-4x+2=0$$
$$a=1,\quad b=-4,\quad c=2$$
$$x=\frac{-(-4)\pm\sqrt{(-4)^2-4(1)(2)}}{2(1)}=\frac{4\pm\sqrt{16-8}}{2}$$
$$=\frac{4\pm\sqrt{8}}{2}=\frac{4\pm2\sqrt{2}}{2}=2\pm\sqrt{2}$$

The zeros of $f(x)=x^2-4x+2$ are $2-\sqrt{2}$ and $2+\sqrt{2}$. The x-intercepts of the graph of f are $2-\sqrt{2}$ and $2+\sqrt{2}$.

40. $f(x)=0$

$$x^2+4x+2=0$$
$$a=1,\quad b=4,\quad c=2$$
$$x=\frac{-4\pm\sqrt{4^2-4(1)(2)}}{2(1)}=\frac{-4\pm\sqrt{16-8}}{2}$$
$$=\frac{-4\pm\sqrt{8}}{2}=\frac{-4\pm2\sqrt{2}}{2}=-2\pm\sqrt{2}$$

The zeros of $f(x)=x^2+4x+2$ are $-2-\sqrt{2}$ and $-2+\sqrt{2}$. The x-intercepts of the graph of f are $-2-\sqrt{2}$ and $-2+\sqrt{2}$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

41. $g(x)=0$

$x^2-4x-1=0$

$a=1,\quad b=-4,\quad c=-1$

$$x=\frac{-(-4)\pm\sqrt{(-4)^2-4(1)(-1)}}{2(1)}=\frac{4\pm\sqrt{16+4}}{2}$$

$$=\frac{4\pm\sqrt{20}}{2}=\frac{4\pm2\sqrt{5}}{2}=2\pm\sqrt{5}$$

The zeros of $g(x)=x^2-4x-1$ are $2-\sqrt{5}$ and $2+\sqrt{5}$. The x-intercepts of the graph of g are $2-\sqrt{5}$ and $2+\sqrt{5}$.

42. $g(x)=0$

$x^2+6x+1=0$

$a=1,\quad b=6,\quad c=1$

$$x=\frac{-6\pm\sqrt{6^2-4(1)(1)}}{2(1)}=\frac{-6\pm\sqrt{36-4}}{2}$$

$$=\frac{-6\pm\sqrt{32}}{2}=\frac{-6\pm4\sqrt{2}}{2}=-3\pm2\sqrt{2}$$

The zeros of $g(x)=x^2+6x+1$ are $-3-2\sqrt{2}$ and $-3+2\sqrt{2}$. The x-intercepts of the graph of g are $-3-2\sqrt{2}$ and $-3+2\sqrt{2}$.

43. $F(x)=0$

$2x^2-5x+3=0$

$a=2,\quad b=-5,\quad c=3$

$$x=\frac{-(-5)\pm\sqrt{(-5)^2-4(2)(3)}}{2(2)}=\frac{5\pm\sqrt{25-24}}{4}$$

$$=\frac{5\pm1}{4}=\frac{3}{2}\text{ or }1$$

The zeros of $F(x)=2x^2-5x+3$ are 1 and $\frac{3}{2}$.

The x-intercepts of the graph of F are 1 and $\frac{3}{2}$.

44. $g(x)=0$

$2x^2+5x+3=0$

$a=2,\quad b=5,\quad c=3$

$$x=\frac{-5\pm\sqrt{5^2-4(2)(3)}}{2(2)}=\frac{-5\pm\sqrt{25-24}}{4}$$

$$=\frac{-5\pm1}{4}=-1\text{ or }-\frac{3}{2}$$

The zeros of $g(x)=2x^2+5x+3$ are $-\frac{3}{2}$ and -1.

The x-intercepts of the graph of g are $-\frac{3}{2}$ and -1.

45. $P(x)=0$

$4x^2-x+2=0$

$a=4,\quad b=-1,\quad c=2$

$$x=\frac{-(-1)\pm\sqrt{(-1)^2-4(4)(2)}}{2(4)}=\frac{1\pm\sqrt{1-32}}{8}$$

$$=\frac{1\pm\sqrt{-31}}{8}=\text{not real}$$

The function $P(x)=4x^2-x+2$ has no real zeros, and the graph of P has no x-intercepts.

46. $H(x)=0$

$4x^2+x+1=0$

$a=4,\quad b=1,\quad c=1$

$$t=\frac{-1\pm\sqrt{1^2-4(4)(1)}}{2(4)}=\frac{-1\pm\sqrt{1-16}}{8}$$

$$=\frac{-1\pm\sqrt{-15}}{8}=\text{not real}$$

The function $H(x)=4x^2+x+1$ has no real zeros, and the graph of H has no x-intercepts.

47. $f(x)=0$

$4x^2-1+2x=0$

$4x^2+2x-1=0$

$a=4,\quad b=2,\quad c=-1$

$$x=\frac{-2\pm\sqrt{2^2-4(4)(-1)}}{2(4)}=\frac{-2\pm\sqrt{4+16}}{8}$$

$$=\frac{-2\pm\sqrt{20}}{8}=\frac{-2\pm2\sqrt{5}}{8}=\frac{-1\pm\sqrt{5}}{4}$$

The zeros of $f(x)=4x^2-1+2x$ are $\frac{-1-\sqrt{5}}{4}$ and $\frac{-1+\sqrt{5}}{4}$. The x-intercepts of the graph of f are $\frac{-1-\sqrt{5}}{4}$ and $\frac{-1+\sqrt{5}}{4}$.

48. $f(x)=0$

$2x^2-1+2x=0$

$2x^2+2x-1=0$

$a=2,\quad b=2,\quad c=-1$

$$x=\frac{-2\pm\sqrt{2^2-4(2)(-1)}}{2(2)}=\frac{-2\pm\sqrt{4+8}}{4}$$

$$=\frac{-2\pm\sqrt{12}}{4}=\frac{-2\pm2\sqrt{3}}{4}=\frac{-1\pm\sqrt{3}}{2}$$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

The zeros of $f(x)=2x^2-1+2x$ are $\frac{-1-\sqrt{3}}{2}$ and $\frac{-1+\sqrt{3}}{2}$. The x-intercepts of the graph of f are $\frac{-1-\sqrt{3}}{2}$ and $\frac{-1+\sqrt{3}}{2}$.

49. $G(x)=0$

$2x(x+2)-3=0$

$2x^2+4x-3=0$

$a=2, \quad b=4, \quad c=-3$

$$x=\frac{-(4)\pm\sqrt{(4)^2-4(2)(-3)}}{2(2)}=\frac{-4\pm\sqrt{16+24}}{4}$$

$$=\frac{-4\pm\sqrt{40}}{4}=\frac{-4\pm2\sqrt{10}}{4}=\frac{-2\pm\sqrt{10}}{2}$$

The zeros of $G(x)=2x(x+2)-3$ are $\frac{-2+\sqrt{10}}{2}$ and $\frac{-2-\sqrt{10}}{2}$. The x-intercepts of the graph of G are $\frac{-2+\sqrt{10}}{2}$ and $\frac{-2-\sqrt{10}}{2}$.

50. $F(x)=0$

$3x(x+2)-1=0 \Rightarrow 3x^2+6x-1=0$

$a=3, \quad b=6, \quad c=-1$

$$x=\frac{-(6)\pm\sqrt{(6)^2-4(3)(-1)}}{2(3)}=\frac{-6\pm\sqrt{36+12}}{6}$$

$$=\frac{-6\pm\sqrt{48}}{6}=\frac{-6\pm4\sqrt{3}}{6}=\frac{-3\pm2\sqrt{3}}{3}$$

The zeros of $F(x)=3x(x+2)-2$ are $\frac{-3+2\sqrt{3}}{3}$ and $\frac{-3-2\sqrt{3}}{3}$. The x-intercepts of the graph of G are $\frac{-3+2\sqrt{3}}{3}$ and $\frac{-3-2\sqrt{3}}{3}$.

51. $p(x)=0$

$9x^2-6x+1=0$

$a=9, \quad b=-6, \quad c=1$

$$x=\frac{-(-6)\pm\sqrt{(-6)^2-4(9)(1)}}{2(9)}=\frac{6\pm\sqrt{36-36}}{18}$$

$$=\frac{6\pm0}{18}=\frac{1}{3}$$

The only real zero of $p(x)=9x^2-6x+1$ is $\frac{1}{3}$.

The only x-intercept of the graph of g is $\frac{1}{3}$.

52. $q(x)=0$

$4x^2+20x+25=0$

$a=4, \quad b=20, \quad c=25$

$$x=\frac{-20\pm\sqrt{(20)^2-4(4)(25)}}{2(4)}=\frac{-20\pm\sqrt{400-400}}{8}$$

$$=\frac{-20\pm0}{8}=-\frac{20}{8}=-\frac{5}{2}$$

The only real zero of $q(x)=4x^2+20x+25$ is $-\frac{5}{2}$. The only x-intercept of the graph of F is $-\frac{5}{2}$.

53. $f(x)=g(x)$

$x^2+6x+3=3$

$x^2+6x=0 \Rightarrow x(x+6)=0$

$x=0 \quad \text{or} \quad x+6=0$

$x=-6$

The x-coordinates of the points of intersection are -6 and 0. The y-coordinates are $g(-6)=3$ and $g(0)=3$. The graphs of the f and g intersect at the points $(-6,3)$ and $(0,3)$.

54. $f(x)=g(x)$

$x^2-4x+3=3$

$x^2-4x=0$

$x(x-4)=0$

$x=0 \quad \text{or} \quad x-4=0$

$x=4$

The x-coordinates of the points of intersection are 0 and 4. The y-coordinates are $g(0)=3$ and $g(4)=3$. The graphs of the f and g intersect at the points $(0,3)$ and $(4,3)$.

55. $f(x)=g(x)$

$-2x^2+1=3x+2$

$0=2x^2+3x+1$

$0=(2x+1)(x+1)$

$2x+1=0 \quad \text{or} \quad x+1=0$

$x=-\frac{1}{2} \qquad x=-1$

The x-coordinates of the points of intersection are -1 and $-\frac{1}{2}$. The y-coordinates are $g(-1)=3(-1)+2=-3+2=-1$ and

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$g\left(-\frac{1}{2}\right)=3\left(-\frac{1}{2}\right)+2=-\frac{3}{2}+2=\frac{1}{2}$.

The graphs of the f and g intersect at the points $(-1,-1)$ and $\left(-\frac{1}{2},\frac{1}{2}\right)$.

56.
$$f(x)=g(x)$$
$$3x^2-7=10x+1$$
$$3x^2-10x-8=0$$
$$(3x+2)(x-4)=0$$
$$3x+2=0 \quad \text{or} \quad x-4=0$$
$$x=-\frac{2}{3} \qquad x=4$$

The x-coordinates of the points of intersection are $-\frac{2}{3}$ and 4. The y-coordinates are $g\left(-\frac{2}{3}\right)=10\left(-\frac{2}{3}\right)+1=-\frac{20}{3}+1=-\frac{17}{3}$ and $g(4)=10(4)+1=40+1=41$.

The graphs of the f and g intersect at the points $\left(-\frac{2}{3},-\frac{17}{3}\right)$ and $(4,41)$.

57.
$$f(x)=g(x)$$
$$x^2-x+1=2x^2-3x-14$$
$$0=x^2-2x-15$$
$$0=(x+3)(x-5)$$
$$x+3=0 \quad \text{or} \quad x-5=0$$
$$x=-3 \qquad x=5$$

The x-coordinates of the points of intersection are -3 and 5. The y-coordinates are $f(-3)=(-3)^2-(-3)+1=9+3+1=13$ and $f(5)=5^2-5+1=25-5+1=21$.

The graphs of the f and g intersect at the points $(-3,13)$ and $(5,21)$.

58.
$$f(x)=g(x)$$
$$x^2+5x-3=2x^2+7x-27$$
$$0=x^2+2x-24$$
$$0=(x+6)(x-4)$$
$$x+6=0 \quad \text{or} \quad x-4=0$$
$$x=-6 \qquad x=4$$

The x-coordinates of the points of intersection are -6 and 4. The y-coordinates are $f(-6)=(-6)^2+5(-6)-3=36-30-3=3$ and $f(4)=4^2+5(4)-3=16+20-3=33$.

The graphs of the f and g intersect at the points $(-6,3)$ and $(4,33)$.

59.
$$P(x)=0$$
$$x^4-6x^2-16=0$$
$$(x^2+2)(x^2-8)=0$$
$$x^2+2=0 \quad \text{or} \quad x^2-8=0$$
$$x^2=-2 \qquad x^2=8$$
$$x=\pm\sqrt{-2} \qquad x=\pm\sqrt{8}$$
$$=\text{not real} \qquad =\pm 2\sqrt{2}$$

The zeros of $P(x)=x^4-6x^2-16$ are $-2\sqrt{2}$ and $2\sqrt{2}$. The x-intercepts of the graph of P are $-2\sqrt{2}$ and $2\sqrt{2}$.

60.
$$H(x)=0$$
$$x^4-3x^2-4=0$$
$$(x^2+1)(x^2-4)=0$$
$$x^2+1=0 \quad \text{or} \quad x^2-4=0$$
$$x^2=-1 \qquad x^2=4$$
$$x=\pm\sqrt{-1} \qquad x=\pm\sqrt{4}$$
$$=\text{not real} \qquad =\pm 2$$

The zeros of $H(x)=x^4-3x^2-4$ are -2 and 2. The x-intercepts of the graph of H are -2 and 2.

61.
$$f(x)=0$$
$$x^4-5x^2+4=0$$
$$(x^2-4)(x^2-1)=0$$
$$x^2-4=0 \text{ or } x^2-1=0$$
$$x=\pm 2 \text{ or } x=\pm 1$$

The zeros of $f(x)=x^4-5x^2+4$ are -2, -1, 1, and 2. The x-intercepts of the graph of f are -2, -1, 1, and 2.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

62. $f(x)=0$

$x^4-10x^2+24=0$

$(x^2-4)(x^2-6)=0$

$x^2-4=0$ or $x^2-6=0$

$x^2=4$ $\quad$ $x^2=6$

$x=\pm 2$ $\quad$ $x=\pm\sqrt{6}$

The zeros of $f(x)=x^4-10x^2+24$ are $-\sqrt{6}$, $\sqrt{6}$, 2 and -2. The x-intercepts of the graph of f are $-\sqrt{6}$, $\sqrt{6}$, 2 and -2.

63. $G(x)=0$

$3x^4-2x^2-1=0$

$(3x^2+1)(x^2-1)=0$

$3x^2+1=0$ or $x^2-1=0$

$x^2=-\frac{1}{3}$ $\quad$ $x^2=1$

$x=\pm\sqrt{-\frac{1}{3}}$ $\quad$ $x=\pm\sqrt{1}$

$x=$ not real $\quad$ $=\pm 1$

The zeros of $G(x)=3x^4-2x^2-1$ are -1 and 1. The x-intercepts of the graph of G are -1 and 1.

64. $F(x)=0$

$2x^4-5x^2-12=0$

$(2x^2+3)(x^2-4)=0$

$2x^2+3=0$ or $x^2-4=0$

$x^2=-\frac{3}{2}$ $\quad$ $x^2=4$

$x=\pm\sqrt{-\frac{3}{2}}$ $\quad$ $x=\pm\sqrt{4}$

$=$ not real $\quad$ $=\pm 2$

The zeros of $F(x)=2x^4-5x^2-12$ are -2 and 2. The x-intercepts of the graph of F are -2 and 2.

65. $g(x)=0$

$x^6+7x^3-8=0$

$(x^3+8)(x^3-1)=0$

$x^3+8=0$ or $x^3-1=0$

$x^3=-8$ $\quad$ $x^3=1$

$x=-2$ $\quad$ $x=1$

The zeros of $g(x)=x^6+7x^3-8$ are -2 and 1. The x-intercepts of the graph of g are -2 and 1.

66. $g(x)=0$

$x^6-7x^3-8=0$

$(x^3-8)(x^3+1)=0$

$x^3-8=0$ or $x^3+1=0$

$x^3=8$ $\quad$ $x^3=-1$

$x=2$ $\quad$ $x=-1$

The zeros of $g(x)=x^6-7x^3-8$ are -1 and 2. The x-intercepts of the graph of g are -1 and 2.

67. $G(x)=0$

$(x+2)^2+7(x+2)+12=0$

Let $u=x+2\rightarrow u^2=(x+2)^2$

$u^2+7u+12=0$

$(u+3)(u+4)=0$

$u+3=0$ or $u+4=0$

$u=-3$ $\quad$ $u=-4$

$x+2=-3$ $\quad$ $x+2=-4$

$x=-5$ $\quad$ $x=-6$

The zeros of $G(x)=(x+2)^2+7(x+2)+12$ are -6 and -5. The x-intercepts of the graph of G are -6 and -5.

68. $f(x)=0$

$(2x+5)^2-(2x+5)-6=0$

Let $u=2x+5\rightarrow u^2=(2x+5)^2$

$u^2-u-6=0$

$(u-3)(u+2)=0$

$u-3=0$ or $u+2=0$

$u=3$ $\quad$ $u=-2$

$2x+5=3$ $\quad$ $2x+5=-2$

$x=-1$ $\quad$ $x=-\frac{7}{2}$

The zeros of $f(x)=(2x+5)^2-(2x+5)-6$ are $-\frac{7}{2}$ and -1. The x-intercepts of the graph of f are $-\frac{7}{2}$ and -1.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

69. $f(x)=0$

$(3x+4)^2-6(3x+4)+9=0$

Let $u=3x+4 \rightarrow u^2=(3x+4)^2$

$u^2-6u+9=0$

$(u-3)^2=0$

$u-3=0$

$u=3$

$3x+4=3$

$x=-\frac{1}{3}$

The only zero of $f(x)=(3x+4)^2-6(3x+4)+9$ is $-\frac{1}{3}$. The x-intercept of the graph of f is $-\frac{1}{3}$.

70. $H(x)=0$

$(2-x)^2+(2-x)-20=0$

Let $u=2-x \rightarrow u^2=(2-x)^2$

$u^2+u-20=0$

$(u+5)(u-4)=0$

$u+5=0$ or $u-4=0$

$u=-5$ $\quad$ $u=4$

$2-x=-5$ $\quad$ $2-x=4$

$x=7$ $\quad$ $x=-2$

The zeros of $H(x)=(2-x)^2+(2-x)-20$ are -2 and 7. The x-intercepts of the graph of H are -2 and 7.

71. $P(x)=0$

$2(x+1)^2-5(x+1)-3=0$

Let $u=x+1 \rightarrow u^2=(x+1)^2$

$2u^2-5u-3=0$

$(2u+1)(u-3)=0$

$2u+1=0$ or $u-3=0$

$u=-\frac{1}{2}$ $\quad$ $u=3$

$x+1=-\frac{1}{2}$ $\quad$ $x+1=3$

$x=-\frac{3}{2}$ $\quad$ $x=2$

The zeros of $P(x)=2(x+1)^2-5(x+1)-3$ are $-\frac{3}{2}$ and 2. The x-intercepts of the graph of P are $-\frac{3}{2}$ and 2.

72. $H(x)=0$

$3(1-x)^2+5(1-x)+2=0$

Let $u=1-x \rightarrow u^2=(1-x)^2$

$3u^2+5u+2=0$

$(3u+2)(u+1)=0$

$3u+2=0$ or $u+1=0$

$u=-\frac{2}{3}$ $\quad$ $u=-1$

$1-x=-\frac{2}{3}$ $\quad$ $1-x=-1$

$x=\frac{5}{3}$ $\quad$ $x=2$

The zeros of $H(x)=3(1-x)^2+5(1-x)+2$ are $\frac{5}{3}$ and 2. The x-intercepts of the graph of H are $\frac{5}{3}$ and 2.

73. $G(x)=0$

$x-4\sqrt{x}=0$

Let $u=\sqrt{x} \rightarrow u^2=x$

$u^2-4u=0$

$u(u-4)=0$

$u=0$ or $u-4=0$

$u=4$

$\sqrt{x}=0$ $\quad$ $\sqrt{x}=4$

$x=0^2=0$ $\quad$ $x=4^2=16$

Check:

$G(0)=0-4\sqrt{0}=0$

$G(16)=16-4\sqrt{16}=16-16=0$

The zeros of $G(x)=x-4\sqrt{x}$ are 0 and 16. The x-intercepts of the graph of G are 0 and 16.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

74. $f(x)=0$

$x+8\sqrt{x}=0$

Let $u=\sqrt{x}\rightarrow u^2=x$

$u^2+8u=0$

$u(u+8)=0$

$u=0$ or $u+8=0$

$u=-8$

$\sqrt{x}=0$ $\sqrt{x}=-8$

$x=0^2=0$ $x=$ not real

Check: $f(0)=0+8\sqrt{0}=0$

The only zero of $f(x)=x+8\sqrt{x}$ is 0. The only x-intercept of the graph of f is 0.

75. $g(x)=0$

$x+\sqrt{x}-20=0$

Let $u=\sqrt{x}\rightarrow u^2=x$

$u^2+u-20=0$

$(u+5)(u-4)=0$

$u+5=0$ or $u-4=0$

$u=-5$ $u=4$

$\sqrt{x}=-5$ $\sqrt{x}=4$

$x=$ not real $x=4^2=16$

Check: $g(16)=16+\sqrt{16}-20=16+4-20=0$

The only zero of $g(x)=x+\sqrt{x}-20$ is 16. The only x-intercept of the graph of g is 16.

76. $f(x)=0$

$x+\sqrt{x}-2=0$

Let $u=\sqrt{x}\rightarrow u^2=x$

$u^2+u-2=0$

$(u-1)(u+2)=0$

$u-1=0$ or $u+2=0$

$u=1$ $u=-2$

$\sqrt{x}=1$ $\sqrt{x}=-2$

$x=1^2=1$ $x=$ not real

Check: $f(1)=1+\sqrt{1}-2=1+1-2=0$

The only zero of $f(x)=x+\sqrt{x}-2$ is 1. The only x-intercept of the graph of f is 1.

77. $f(x)=0$

$x^2-50=0$

$x^2=50\Rightarrow x=\pm\sqrt{50}=\pm5\sqrt{2}$

The zeros of $f(x)=x^2-50$ are $-5\sqrt{2}$ and $5\sqrt{2}$. The x-intercepts of the graph of f are $-5\sqrt{2}$ and $5\sqrt{2}$.

78. $f(x)=0$

$x^2-20=0$

$x^2=20\Rightarrow x=\pm\sqrt{20}=\pm2\sqrt{5}$

The zeros of $f(x)=x^2-6$ are $-2\sqrt{5}$ and $2\sqrt{5}$. The x-intercepts of the graph of f are $-2\sqrt{5}$ and $2\sqrt{5}$.

79. $g(x)=0$

$16x^2-8x+1=0$

$(4x-1)^2=0$

$4x-1=0\Rightarrow x=\frac{1}{4}$

The only real zero of $g(x)=16x^2-8x+1$ is $\frac{1}{4}$.

The only x-intercept of the graph of g is $\frac{1}{4}$.

80. $F(x)=0$

$4x^2-12x+9=0$

$(2x-3)^2=0$

$2x-3=0\Rightarrow x=\frac{3}{2}$

The only real zero of $F(x)=4x^2-12x+9$ is $\frac{3}{2}$.

The only x-intercept of the graph of F is $\frac{3}{2}$.

81. $G(x)=0$

$10x^2-19x-15=0$

$(5x+3)(2x-5)=0$

$5x+3=0$ or $2x-5=0$

$x=-\frac{3}{5}$ $x=\frac{5}{2}$

The zeros of $G(x)=10x^2-19x-15$ are $-\frac{3}{5}$ and

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$\frac{5}{2}$. The x-intercepts of the graph of G are $-\frac{3}{5}$ and $\frac{5}{2}$.

82.
$$f(x)=0$$
$$6x^2+7x-20=0$$
$$(3x-4)(2x+5)=0$$
$$3x-4=0 \quad \text{or} \quad 2x+5=0$$
$$x=\frac{4}{3} \qquad x=-\frac{5}{2}$$

The zeros of $f(x)=6x^2+7x-20$ are $-\frac{5}{2}$ and $\frac{4}{3}$.

The x-intercepts of the graph of f are $-\frac{5}{2}$ and $\frac{4}{3}$.

83.
$$P(x)=0$$
$$6x^2-x-2=0$$
$$(3x-2)(2x+1)=0$$
$$3x-2=0 \quad \text{or} \quad 2x+1=0$$
$$x=\frac{2}{3} \qquad x=-\frac{1}{2}$$

The zeros of $P(x)=6x^2-x-2$ are $-\frac{1}{2}$ and $\frac{2}{3}$.

The x-intercepts of the graph of P are $-\frac{1}{2}$ and $\frac{2}{3}$.

84.
$$H(x)=0$$
$$6x^2+x-2=0$$
$$(3x+2)(2x-1)=0$$
$$3x+2=0 \quad \text{or} \quad 2x-1=0$$
$$x=-\frac{2}{3} \qquad x=\frac{1}{2}$$

The zeros of $H(x)=6x^2+x-2$ are $-\frac{2}{3}$ and $\frac{1}{2}$.

The x-intercepts of the graph of H are $-\frac{2}{3}$ and $\frac{1}{2}$.

85.
$$G(x)=0$$
$$x^2+\sqrt{2}x-\frac{1}{2}=0$$
$$2\left(x^2+\sqrt{2}x-\frac{1}{2}\right)=(0)(2)$$
$$2x^2+2\sqrt{2}x-1=0$$
$$a=2, \quad b=2\sqrt{2}, \quad c=-1$$
$$x=\frac{-(2\sqrt{2})\pm\sqrt{(2\sqrt{2})^2-4(2)(-1)}}{2(2)}$$
$$=\frac{-2\sqrt{2}\pm\sqrt{8+8}}{4}=\frac{-2\sqrt{2}\pm\sqrt{16}}{4}$$
$$=\frac{-2\sqrt{2}\pm 4}{4}=\frac{-\sqrt{2}\pm 2}{2}$$

The zeros of $G(x)=x^2+\sqrt{2}x-\frac{1}{2}$ are $\frac{-\sqrt{2}-2}{2}$ and $\frac{-\sqrt{2}+2}{2}$. The x-intercepts of the graph of G are $\frac{-\sqrt{2}-2}{2}$ and $\frac{-\sqrt{2}+2}{2}$.

86.
$$F(x)=0$$
$$\frac{1}{2}x^2-\sqrt{2}x-1=0$$
$$2\left(\frac{1}{2}x^2-\sqrt{2}x-1\right)=(0)(2)$$
$$x^2-2\sqrt{2}x-2=0$$
$$a=1, \quad b=-2\sqrt{2}, \quad c=-2$$
$$x=\frac{-(-2\sqrt{2})\pm\sqrt{(-2\sqrt{2})^2-4(1)(-2)}}{2(1)}$$
$$=\frac{2\sqrt{2}\pm\sqrt{16}}{2}=\frac{2\sqrt{2}\pm 4}{2}=\frac{\sqrt{2}\pm 2}{1}$$

The zeros of $F(x)=\frac{1}{2}x^2-\sqrt{2}x-1$ are $\sqrt{2}-2$ and $\sqrt{2}+2$. The x-intercepts of the graph of F are $\sqrt{2}-2$ and $\sqrt{2}+2$.

87.
$$f(x)=0$$
$$x^2+x-4=0$$
$$a=1, \quad b=1, \quad c=-4$$
$$x=\frac{-(1)\pm\sqrt{(1)^2-4(1)(-4)}}{2(1)}$$
$$=\frac{-1\pm\sqrt{1+16}}{2}=\frac{-1\pm\sqrt{17}}{2}$$

The zeros of $f(x)=x^2+x-4$ are $\frac{-1-\sqrt{17}}{2}$ and $\frac{-1+\sqrt{17}}{2}$. The x-intercepts of the graph of f are $\frac{-1-\sqrt{17}}{2}$ and $\frac{-1+\sqrt{17}}{2}$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

88. $g(x)=0$

$x^2+x-1=0$


$a=1,\quad b=1,\quad c=-1$

$$x=\frac{-(1)\pm\sqrt{(1)^2-4(1)(-1)}}{2(1)}=\frac{-1\pm\sqrt{5}}{2}$$

The zeros of $g(x)=x^2+x-1$ are $\frac{-1-\sqrt{5}}{2}$ and $\frac{-1+\sqrt{5}}{2}$. The x-intercepts of the graph of g are $\frac{-1-\sqrt{5}}{2}$ and $\frac{-1+\sqrt{5}}{2}$.

89. **a.** $g(x)=(x-1)^2-4$

Using the graph of $y=x^2$, horizontally shift to the right 1 unit, and then vertically shift downward 4 units.



b. $g(x)=0$

$(x-1)^2-4=0$


$x^2-2x+1-4=0$

$x^2-2x-3=0$

$(x+1)(x-3)=0\Rightarrow x=-1$ or $x=3$

90. **a.** $F(x)=(x+3)^2-9$

Using the graph of $y=x^2$, horizontally shift to the left 3 units, and then vertically shift downward 9 units.



b. $F(x)=0$

$(x+3)^2-9=0$


$x^2+6x+9-9=0$

$x^2+6x=0$

$x(x+6)=0\Rightarrow x=0$ or $x=-6$

91. **a.** $f(x)=2(x+4)^2-8$

Using the graph of $y=x^2$, horizontally shift to the left 4 units, vertically stretch by a factor of 2, and then vertically shift downward 8 units.



b. $f(x)=0$

$2(x+4)^2-8=0$

$2(x^2+8x+16)-8=0$


$2x^2+16x+32-8=0$

$2x^2+16x+24=0$

$2(x+2)(x+6)=0\Rightarrow x=-2$ or $x=-6$

92. **a.** $h(x)=3(x-2)^2-12$

Using the graph of $y=x^2$, horizontally shift to the right 2 units, vertically stretch by a factor of 3, and then vertically shift downward 12 units.




Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

b.
$$h(x) = 0$$
$$3(x-2)^2 - 12 = 0$$
$$3(x^2 - 4x + 4) - 12 = 0$$
$$3x^2 - 12x + 12 - 12 = 0$$
$$3x^2 - 12x = 0$$
$$3x(x-4) = 0 \Rightarrow x = 0 \text{ or } x = 4$$

93. a. $H(x) = -3(x-3)^2 + 6$

Using the graph of $y = x^2$, horizontally shift to the right 3 units, vertically stretch by a factor of 3, reflect about the x-axis, and then vertically shift upward 6 units.




b.
$$H(x) = 0$$
$$-3(x-3)^2 + 6 = 0$$
$$-3(x^2 - 6x + 9) + 6 = 0$$
$$-3x^2 + 18x - 27 + 6 = 0$$
$$-3x^2 + 18x - 21 = 0$$
$$-3(x^2 - 6x + 7) = 0$$
$$a = 1, \quad b = -6, \quad c = 7$$

$$x = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(1)(7)}}{2(1)} = \frac{6 \pm \sqrt{36 - 28}}{2}$$
$$= \frac{6 \pm \sqrt{8}}{2} = \frac{6 \pm 2\sqrt{2}}{2} = 3 \pm \sqrt{2}$$

94. a. $f(x) = -2(x+1)^2 + 12$

Using the graph of $y = x^2$, horizontally shift to the left 1 unit, vertically stretch by a factor of 2, reflect about the x-axis, and then vertically shift upward 12 units.



b.
$$f(x) = 0$$
$$-2(x+1)^2 + 12 = 0$$
$$-2(x^2 + 2x + 1) + 12 = 0$$
$$-2x^2 - 4x - 2 + 12 = 0$$
$$-2x^2 - 4x + 10 = 0$$
$$-2(x^2 + 2x - 5) = 0$$
$$a = 1, \quad b = 2, \quad c = -5$$

$$x = \frac{-(2) \pm \sqrt{(2)^2 - 4(1)(-5)}}{2(1)} = \frac{-2 \pm \sqrt{4 + 20}}{2}$$
$$= \frac{-2 \pm \sqrt{24}}{2} = \frac{-2 \pm 2\sqrt{6}}{2} = -1 \pm \sqrt{6}$$

95.
$$f(x) = g(x)$$
$$5x(x-1) = -7x^2 + 2$$
$$5x^2 - 5x = -7x^2 + 2$$
$$12x^2 - 5x - 2 = 0$$
$$(3x-2)(4x+1) = 0 \Rightarrow x = \frac{2}{3} \text{ or } x = -\frac{1}{4}$$
$$f\left(\frac{2}{3}\right) = 5\left(\frac{2}{3}\right)\left[\left(\frac{2}{3}\right) - 1\right]$$
$$= \left(\frac{10}{3}\right)\left(-\frac{1}{3}\right) = -\frac{10}{9}$$
$$f\left(-\frac{1}{4}\right) = 5\left(-\frac{1}{4}\right)\left[\left(-\frac{1}{4}\right) - 1\right]$$
$$= \left(-\frac{5}{4}\right)\left(-\frac{5}{4}\right) = \frac{25}{16}$$

The points of intersection are:
$\left(\frac{2}{3}, -\frac{10}{9}\right)$ and $\left(-\frac{1}{4}, \frac{25}{16}\right)$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

96.
$$f(x) = g(x)$$
$$10x(x+2) = -3x+5$$
$$10x^2 + 20x = -3x+5$$
$$10x^2 + 23x - 5 = 0$$
$$(2x+5)(5x-1) = 0 \Rightarrow x = -\frac{5}{2} \text{ or } x = \frac{1}{5}$$
$$f\left(-\frac{5}{2}\right) = 10\left(-\frac{5}{2}\right)\left[\left(-\frac{5}{2}\right)+2\right]$$
$$= (-25)\left(-\frac{1}{2}\right) = \frac{25}{2}$$
$$f\left(\frac{1}{5}\right) = 10\left(\frac{1}{5}\right)\left[\left(\frac{1}{5}\right)+2\right]$$
$$= (2)\left(\frac{11}{5}\right) = \frac{22}{5}$$

The points of intersection are:
$\left(-\frac{5}{2}, \frac{25}{2}\right)$ and $\left(\frac{1}{5}, \frac{22}{5}\right)$

97.
$$f(x) = g(x)$$
$$3(x^2-4) = 3x^2+2x+4$$
$$3x^2 - 12 = 3x^2+2x+4$$
$$-12 = 2x+4$$
$$-16 = 2x \Rightarrow x = -8$$
$$f(-8) = 3\left[(-8)^2 - 4\right]$$
$$= 3[64-4] = 180$$

The point of intersection is: $(-8, 180)$

98.
$$f(x) = g(x)$$
$$4(x^2+1) = 4x^2 - 3x - 8$$
$$4x^2 + 4 = 4x^2 - 3x - 8$$
$$4 = -3x - 8$$
$$12 = -3x \Rightarrow x = -4$$
$$f(-4) = 4\left[(-4)^2 + 1\right]$$
$$= 4[16+1] = 68$$

The point of intersection is: $(-4, 68)$

99.
$$f(x) = g(x)$$
$$\frac{3x}{x+2} - \frac{5}{x+1} = \frac{-5}{x^2+3x+2}$$
$$\frac{3x}{x+2} - \frac{5}{x+1} = \frac{-5}{(x+2)(x+1)}$$
$$3x(x+1) - 5(x+2) = -5$$
$$3x^2 + 3x - 5x - 10 = -5$$
$$3x^2 - 2x - 5 = 0$$
$$(3x-5)(x+1) = 0$$
$$x = \frac{5}{3} \text{ or } \cancel{x = -1}$$
$$f\left(\frac{5}{3}\right) = \frac{3\left(\frac{5}{3}\right)}{\left(\frac{5}{3}\right)+2} - \frac{5}{\left(\frac{5}{3}\right)+1}$$
$$= \frac{(5)}{\left(\frac{11}{3}\right)} - \frac{5}{\left(\frac{8}{3}\right)}$$
$$= \frac{15}{11} - \frac{15}{8}$$
$$= -\frac{45}{88}$$

The point of intersection is: $\left(\frac{5}{3}, -\frac{45}{88}\right)$

100.
$$f(x) = g(x)$$
$$\frac{2x}{x-3} - \frac{3}{x+1} = \frac{2x+18}{x^2-2x-3}$$
$$\frac{2x}{x-3} - \frac{3}{x+1} = \frac{2x+18}{(x-3)(x+1)}$$
$$2x(x+1) - 3(x-3) = 2x+18$$
$$2x^2 + 2x - 3x + 9 = 2x+18$$
$$2x^2 - 3x - 9 = 0$$
$$(2x+3)(x-3) = 0$$
$$x = -\frac{3}{2} \text{ or } \cancel{x = 3}$$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$$f\left(-\frac{3}{2}\right)=\frac{2\left(-\frac{3}{2}\right)}{\left(-\frac{3}{2}\right)-3}-\frac{3}{\left(-\frac{3}{2}\right)+1}$$
$$=\frac{(-3)}{\left(-\frac{9}{2}\right)}-\frac{3}{\left(-\frac{1}{2}\right)}$$
$$=\frac{6}{9}+6=\frac{2}{3}+6=\frac{20}{3}$$

The point of intersection is: $\left(-\frac{3}{2},\frac{20}{3}\right)$

101. a. $(f+g)(x)=$
$$=x^2+5x-14+x^2+3x-4$$
$$=2x^2+8x-18$$
$$2x^2+8x-18=0$$
$$x^2+4x-9=0$$
$$x=\frac{-(4)\pm\sqrt{(4)^2-4(1)(-9)}}{2(1)}=\frac{-4\pm\sqrt{16+36}}{2}$$
$$=\frac{-4\pm\sqrt{52}}{2}=\frac{-4\pm2\sqrt{13}}{2}=-2\pm\sqrt{13}$$

b. $(f-g)(x)=$
$$=\left(x^2+5x-14\right)-\left(x^2+3x-4\right)$$
$$=x^2+5x-14-x^2-3x+4$$
$$=2x-10$$
$$2x-10=0\Rightarrow x=5$$

c. $(f\cdot g)(x)=$
$$=\left(x^2+5x-14\right)\left(x^2+3x-4\right)$$
$$=(x+7)(x-2)(x+4)(x-1)$$

$$(f\cdot g)(x)=0$$
$$0=(x+7)(x-2)(x+4)(x-1)$$
$$\Rightarrow x=-7 \text{ or } x=2 \text{ or } x=-4 \text{ or } x=1$$

102. a. $(f+g)(x)=$
$$=x^2-3x-18+x^2+2x-3$$
$$=2x^2-x-21$$
$$2x^2-x-21=0$$
$$(2x-7)(x+3)=0\Rightarrow x=\frac{7}{2} \text{ or } x=-3$$

b. $(f-g)(x)=$
$$=\left(x^2-3x-18\right)-\left(x^2+2x-3\right)$$
$$=x^2-3x-18-x^2-2x+3$$
$$=-5x-15$$
$$-5x-15=0\Rightarrow x=-3$$

c. $(f\cdot g)(x)=$
$$=\left(x^2-3x-18\right)\left(x^2+2x-3\right)$$
$$=(x+3)(x-6)(x+3)(x-1)$$
$$(f\cdot g)(x)=0$$
$$0=(x+3)(x-6)(x+3)(x-1)$$
$$\Rightarrow x=-3 \text{ or } x=6 \text{ or } x=1$$

103.
$$A(x)=143$$
$$x(x+2)=143$$
$$x^2+2x-143=0$$
$$(x+13)(x-11)=0$$
$$\cancel{x=-13} \text{ or } x=11$$

Discard the negative solution since width cannot be negative. The width of the rectangular window is 11 feet and the length is 13 feet.

104.
$$A(x)=306$$
$$x(x+1)=306$$
$$x^2+x-306=0$$
$$(x+18)(x-17)=0$$
$$\cancel{x=-18} \text{ or } x=17$$

Discard the negative solution since width cannot be negative. The width of the rectangular window is 17 cm and the length is 18 cm.

105.
$$V(x)=4$$
$$(x-2)^2=4$$
$$x-2=\pm\sqrt{4}$$
$$x-2=\pm2$$
$$x=2\pm2$$
$$x=4 \text{ or } x=0$$

Discard $x=0$ since that is not a feasible length for the original sheet. Therefore, the original sheet should measure 4 feet on each side.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

106. $V(x) = 4$

$(x-2)^2 = 16$

$x-2 = \pm\sqrt{16}$

$x-2 = \pm 4$

$x = 2 \pm 4$

$x = 6$ or ~~$x = -2$~~

Discard $x = -2$ since width cannot be negative. Therefore, the original sheet should measure 6 feet on each side.

107. **a.** When the ball strikes the ground, the distance from the ground will be 0. Therefore, we solve

$s = 0$

$96 + 80t - 16t^2 = 0$

$-16t^2 + 80t + 96 = 0$

$t^2 - 5t - 6 = 0$

$(t-6)(t+1) = 0$

$t = 6$ or ~~$t = -1$~~

Discard the negative solution since the time of flight must be positive. The ball will strike the ground after 6 seconds.

b. When the ball passes the top of the building, it will be 96 feet from the ground. Therefore, we solve

$s = 96$

$96 + 80t - 16t^2 = 96$

$-16t^2 + 80t = 0$

$t^2 - 5t = 0$

$t(t-5) = 0$

$t = 0$ or $t = 5$

The ball is at the top of the building at time $t = 0$ seconds when it is thrown. It will pass the top of the building on the way down after 5 seconds.

108. **a.** To find when the object will be 15 meters above the ground, we solve

$s = 15$

$-4.9t^2 + 20t = 15$

$-4.9t^2 + 20t - 15 = 0$

$a = -4.9,\ b = 20,\ c = -15$

$$t = \frac{-20 \pm \sqrt{20^2 - 4(-4.9)(-15)}}{2(-4.9)}$$

$$= \frac{-20 \pm \sqrt{106}}{-9.8}$$

$$= \frac{20 \pm \sqrt{106}}{9.8}$$

$t \approx 0.99$ or $t \approx 3.09$

The object will be 15 meters above the ground after about 0.99 seconds (on the way up) and about 3.09 seconds (on the way down).

b. The object will strike the ground when the distance from the ground is 0. Thus, we solve

$s = 0$

$-4.9t^2 + 20t = 0$

$t(-4.9t + 20) = 0$

$t = 0$ or $-4.9t + 20 = 0$

$-4.9t = -20$

$t \approx 4.08$

The object will strike the ground after about 4.08 seconds.

c. $s = 100$

$-4.9t^2 + 20t = 100$

$-4.9t^2 + 20t - 100 = 0$

$a = -4.9,\ b = 20,\ c = -100$

$$t = \frac{-20 \pm \sqrt{20^2 - 4(-4.9)(-100)}}{2(-4.9)}$$

$$= \frac{-20 \pm \sqrt{-1560}}{-9.8}$$

There is no real solution. The object never reaches a height of 100 meters.

109. For the sum to be 210, we solve

$S(n) = 210$

$\frac{1}{2}n(n+1) = 210$

$n(n+1) = 420$

$n^2 + n - 420 = 0$

$(n-20)(n+21) = 0$

$n - 20 = 0$ or $n + 21 = 0$

$n = 20$ ~~$n = -21$~~

Discard the negative solution since the number of consecutive integers must be positive. For a sum of 210, we must add the 20 consecutive integers, starting at 1.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

110. To determine the number of sides when a polygon has 65 diagonals, we solve

$$D(n) = 65$$
$$\frac{1}{2}n(n-3) = 65$$
$$n(n-3) = 130$$
$$n^2 - 3n - 130 = 0$$
$$(n+10)(n-13) = 0$$
$$n+10 = 0 \quad \text{or} \quad n-13 = 0$$
$$\cancel{n = -10} \qquad n = 13$$

Discard the negative solution since the number of sides must be positive. A polygon with 65 diagonals will have 13 sides.

To determine the number of sides if a polygon has 80 diagonals, we solve

$$D(n) = 80$$
$$\frac{1}{2}n(n-3) = 80$$
$$n(n-3) = 160$$
$$n^2 - 3n - 160 = 0$$
$$a = 1,\ b = -3,\ c = -160$$
$$t = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(-160)}}{2(1)}$$
$$= \frac{3 \pm \sqrt{649}}{2}$$

Since the solutions are not integers, a polygon with 80 diagonals is not possible.

111. The roots of a quadratic equation are

$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \text{ and } x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a},$$

so the sum of the roots is

$$x_1 + x_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} + \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$
$$= \frac{-b - \sqrt{b^2 - 4ac} - b + \sqrt{b^2 - 4ac}}{2a}$$
$$= \frac{-2b}{2a} = -\frac{b}{a}$$

112. The roots of a quadratic equation are

$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \text{ and } x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a},$$

so the product of the roots is

$$x_1 \cdot x_2 = \left(\frac{-b - \sqrt{b^2 - 4ac}}{2a}\right)\left(\frac{-b + \sqrt{b^2 - 4ac}}{2a}\right)$$
$$= \frac{(-b)^2 - \left(\sqrt{b^2 - 4ac}\right)^2}{(2a)^2} = \frac{b^2 - (b^2 - 4ac)}{4a^2}$$
$$= \frac{b^2 - b^2 + 4ac}{4a^2} = \frac{4ac}{4a^2} = \frac{c}{a}$$

113. In order to have one repeated real zero, we need the discriminant to be 0.

$$b^2 - 4ac = 0$$
$$1^2 - 4(k)(k) = 0$$
$$1 - 4k^2 = 0$$
$$4k^2 = 1$$
$$k^2 = \frac{1}{4}$$
$$k = \pm\sqrt{\frac{1}{4}}$$
$$k = \frac{1}{2} \quad \text{or} \quad k = -\frac{1}{2}$$

114. In order to have one repeated real zero, we need the discriminant to be 0.

$$b^2 - 4ac = 0$$
$$(-k)^2 - 4(1)(4) = 0$$
$$k^2 - 16 = 0$$
$$(k-4)(k+4) = 0$$
$$k = 4 \quad \text{or} \quad k = -4$$

115. For $f(x) = ax^2 + bx + c = 0$:

$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \text{ and } x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

For $f(x) = ax^2 - bx + c = 0$:

$$x_1^* = \frac{-(-b) - \sqrt{(-b)^2 - 4ac}}{2a}$$
$$= \frac{b - \sqrt{b^2 - 4ac}}{2a} = -\left(\frac{-b + \sqrt{b^2 - 4ac}}{2a}\right) = -x_2$$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

and

$$x_2^* = \frac{-(-b)+\sqrt{(-b)^2-4ac}}{2a}$$
$$= \frac{b+\sqrt{b^2-4ac}}{2a} = -\left(\frac{-b-\sqrt{b^2-4ac}}{2a}\right) = -x_1$$

116. For $f(x) = ax^2+bx+c = 0$:

$$x_1 = \frac{-b-\sqrt{b^2-4ac}}{2a} \text{ and } x_2 = \frac{-b+\sqrt{b^2-4ac}}{2a}$$

For $f(x) = cx^2+bx+a = 0$:

$$x_1^* = \frac{-b-\sqrt{b^2-4(c)(a)}}{2c} = \frac{-b-\sqrt{b^2-4ac}}{2c}$$
$$= \frac{-b-\sqrt{b^2-4ac}}{2c}\cdot\frac{-b+\sqrt{b^2-4ac}}{-b+\sqrt{b^2-4ac}}$$
$$= \frac{b^2-\left(b^2-4ac\right)}{2c\left(-b+\sqrt{b^2-4ac}\right)} = \frac{4ac}{2c\left(-b+\sqrt{b^2-4ac}\right)}$$
$$= \frac{2a}{-b+\sqrt{b^2-4ac}} = \frac{1}{x_2}$$

and

$$x_2^* = \frac{-b+\sqrt{b^2-4(c)(a)}}{2c} = \frac{-b+\sqrt{b^2-4ac}}{2c}$$
$$= \frac{-b+\sqrt{b^2-4ac}}{2c}\cdot\frac{-b-\sqrt{b^2-4ac}}{-b-\sqrt{b^2-4ac}}$$
$$= \frac{b^2-\left(b^2-4ac\right)}{2c\left(-b-\sqrt{b^2-4ac}\right)} = \frac{4ac}{2c\left(-b-\sqrt{b^2-4ac}\right)}$$
$$= \frac{2a}{-b-\sqrt{b^2-4ac}} = \frac{1}{x_1}$$

117. **a.** $x^2 = 9$ and $x = 3$ are not equivalent because they do not have the same solution set. In the first equation we can also have $x = -3$.

b. $x = \sqrt{9}$ and $x = 3$ are equivalent because $\sqrt{9} = 3$.

c. $(x-1)(x-2) = (x-1)^2$ and $x-2 = x-1$ are not equivalent because they do not have the same solution set. The first equation has the solution set $\{1\}$ while the second equation has no solutions.

118. Answers may vary. Methods discussed in this section include factoring, the square root method, completing the square, and the quadratic formula.

119. Answers will vary. Knowing the discriminant allows us to know how many real solutions the equation will have.

120. Answers will vary. One possibility:

Two distinct: $f(x) = x^2-3x-18$


One repeated: $f(x) = x^2-14x+49$

No real: $f(x) = x^2+x+4$

121. Answers will vary.

122. Two quadratic functions can intersect 0, 1, or 2 times.

123. The graph is shifted vertically by 4 units and is reflected about the x-axis.



124. Domain: $\{-3,-1,1,3\}$ Range: $\{2,4\}$

125. $\bar{x} = \frac{-10+2}{2} = \frac{-8}{2} = -4$

$\bar{y} = \frac{4+(-1)}{2} = \frac{3}{2}$

So the midpoint is: $\left(-4,\frac{3}{2}\right)$.

126. If the graph is symmetric with respect to the y-axis then x and –x are on the graph. Thus if $(-1,4)$ is on the graph, then so is $(1,4)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Section 2.4

1. $y = x^2 - 9$

To find the y-intercept, let $x = 0$:

$y = 0^2 - 9 = -9$.

To find the x-intercept(s), let $y = 0$:

$x^2 - 9 = 0$

$x^2 = 9$

$x = \pm\sqrt{9} = \pm 3$

The intercepts are $(0, -9), (-3, 0),$ and $(3, 0)$.

2. $2x^2 + 7x - 4 = 0$

$(2x - 1)(x + 4) = 0$

$2x - 1 = 0$ or $x + 4 = 0$

$2x = 1$ or $x = -4$

$x = \frac{1}{2}$ or $x = -4$

The solution set is $\left\{-4, \frac{1}{2}\right\}$..

3. $\left(\frac{1}{2} \cdot (-5)\right)^2 = \frac{25}{4}$

4. right; 4

5. parabola

6. axis (or axis of symmetry)

7. $-\frac{b}{2a}$

8. True; $a = 2 > 0$.

9. True; $-\frac{b}{2a} = -\frac{4}{2(-1)} = 2$

10. True

11. a

12. d

13. C

14. E

15. F

16. A

17. G


18. B

19. H

20. D


21. $f(x) = \frac{1}{4}x^2$

Using the graph of $y = x^2$, compress vertically by a factor of $\frac{1}{4}$.




22. $f(x) = 2x^2 + 4$

Using the graph of $y = x^2$, stretch vertically by a factor of 2, then shift up 4 units.



23. $f(x) = (x + 2)^2 - 2$

Using the graph of $y = x^2$, shift left 2 units, then shift down 2 units.




Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

24. $f(x) = (x-3)^2 - 10$

Using the graph of $y = x^2$, shift right 3 units, then shift down 10 units.




25. $f(x) = x^2 + 4x + 1$

$= (x^2 + 4x + 4) + 1 - 4$

$= (x+2)^2 - 3$

Using the graph of $y = x^2$, shift left 2 units, then shift down 3 units.




26. $f(x) = x^2 - 6x - 1$

$= (x^2 - 6x + 9) - 1 - 9$

$= (x-3)^2 - 10$

Using the graph of $y = x^2$, shift right 3 units, then shift down 10 units.




27. $f(x) = 2x^2 - 4x + 1$

$= 2\left(x^2 - 2x\right) + 1$

$= 2(x^2 - 2x + 1) + 1 - 2$

$= 2(x-1)^2 - 1$

Using the graph of $y = x^2$, shift right 1 unit, stretch vertically by a factor of 2, then shift down 1 unit.




28. $f(x) = 3x^2 + 6x$

$= 3\left(x^2 + 2x\right)$

$= 3(x^2 + 2x + 1) - 3$

$= 3(x+1)^2 - 3$

Using the graph of $y = x^2$, shift left 1 unit, stretch vertically by a factor of 3, then shift down 3 units.



29. $f(x) = -x^2 - 2x$

$= -\left(x^2 + 2x\right)$


$= -(x^2 + 2x + 1) + 1$

$= -(x+1)^2 + 1$

Using the graph of $y = x^2$, shift left 1 unit, reflect across the x-axis, then shift up 1 unit.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual




30. $f(x) = -2x^2 + 6x + 2$

$= -2\left(x^2 - 3x\right) + 2$

$= -2\left(x^2 - 3x + \frac{9}{4}\right) + 2 + \frac{9}{2}$

$= -2\left(x - \frac{3}{2}\right)^2 + \frac{13}{2}$

Using the graph of $y = x^2$, shift right $\frac{3}{2}$ units, reflect about the x-axis, stretch vertically by a factor of 2, then shift up $\frac{13}{2}$ units.




31. $f(x) = \frac{1}{2}x^2 + x - 1$

$= \frac{1}{2}\left(x^2 + 2x\right) - 1$

$= \frac{1}{2}\left(x^2 + 2x + 1\right) - 1 - \frac{1}{2}$

$= \frac{1}{2}(x+1)^2 - \frac{3}{2}$

Using the graph of $y = x^2$, shift left 1 unit, compress vertically by a factor of $\frac{1}{2}$, then shift down $\frac{3}{2}$ units.




32. $f(x) = \frac{2}{3}x^2 + \frac{4}{3}x - 1$

$= \frac{2}{3}\left(x^2 + 2x\right) - 1$

$= \frac{2}{3}\left(x^2 + 2x + 1\right) - 1 - \frac{2}{3}$

$= \frac{2}{3}(x+1)^2 - \frac{5}{3}$

Using the graph of $y = x^2$, shift left 1 unit, compress vertically by a factor of $\frac{2}{3}$, then shift down $\frac{5}{3}$ unit.



33. a. For $f(x) = x^2 + 2x$, $a = 1$, $b = 2$, $c = 0$.
Since $a = 1 > 0$, the graph opens up.
The x-coordinate of the vertex is
$x = \frac{-b}{2a} = \frac{-(2)}{2(1)} = \frac{-2}{2} = -1$.
The y-coordinate of the vertex is
$f\left(\frac{-b}{2a}\right) = f(-1) = (-1)^2 + 2(-1) = 1 - 2 = -1.$
Thus, the vertex is $(-1, -1)$.
The axis of symmetry is the line $x = -1$.
The discriminant is
$b^2 - 4ac = (2)^2 - 4(1)(0) = 4 > 0$, so the graph has two x-intercepts.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

The x-intercepts are found by solving:


$x^2 + 2x = 0$

$x(x+2) = 0$

$x = 0$ or $x = -2$

The x-intercepts are -2 and 0.

The y-intercept is $f(0) = 0$.



b. The domain is $(-\infty, \infty)$.

The range is $[-1, \infty)$.

c. Decreasing on $(-\infty, -1]$.

Increasing on $[-1, \infty)$.

34. a. For $f(x) = x^2 - 4x$, $a = 1$, $b = -4$, $c = 0$.

Since $a = 1 > 0$, the graph opens up.

The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-4)}{2(1)} = \frac{4}{2} = 2.$$

The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(2) = (2)^2 - 4(2) = 4 - 8 = -4.$$

Thus, the vertex is $(2, -4)$.

The axis of symmetry is the line $x = 2$.

The discriminant is:

$b^2 - 4ac = (-4)^2 - 4(1)(0) = 16 > 0$, so the graph has two x-intercepts.

The x-intercepts are found by solving:


$x^2 - 4x = 0$

$x(x-4) = 0$

$x = 0$ or $x = 4$.

The x-intercepts are 0 and 4.

The y-intercept is $f(0) = 0$.



b. The domain is $(-\infty, \infty)$.

The range is $[-4, \infty)$.

c. Decreasing on $(-\infty, 2]$.

Increasing on $[2, \infty)$.

35. a. For $f(x) = -x^2 - 6x$, $a = -1$, $b = -6$, $c = 0$. Since $a = -1 < 0$, the graph opens down. The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-6)}{2(-1)} = \frac{6}{-2} = -3.$$

The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(-3) = -(-3)^2 - 6(-3)$$

$$= -9 + 18 = 9.$$

Thus, the vertex is $(-3, 9)$.

The axis of symmetry is the line $x = -3$.

The discriminant is:

$b^2 - 4ac = (-6)^2 - 4(-1)(0) = 36 > 0$,

so the graph has two x-intercepts.

The x-intercepts are found by solving:


$-x^2 - 6x = 0$

$-x(x+6) = 0$

$x = 0$ or $x = -6$.

The x-intercepts are -6 and 0.

The y-intercepts are $f(0) = 0$.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, 9]$.

c. Increasing on $(-\infty, -3]$.
Decreasing on $[-3, \infty)$.


36. a. For $f(x) = -x^2 + 4x$, $a = -1$, $b = 4$, $c = 0$.
Since $a = -1 < 0$, the graph opens down.
The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-4}{2(-1)} = \frac{-4}{-2} = 2.$$

The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(2)$$
$$= -(2)^2 + 4(2)$$
$$= 4.$$

Thus, the vertex is $(2, 4)$.
The axis of symmetry is the line $x = 2$.
The discriminant is:
$b^2 - 4ac = 4^2 - 4(-1)(0) = 16 > 0$,
so the graph has two x-intercepts.
The x-intercepts are found by solving:
$-x^2 + 4x = 0$
$-x(x - 4) = 0$
$x = 0$ or $x = 4$.
The x-intercepts are 0 and 4.
The y-intercept is $f(0) = 0$.



b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, 4]$.

c. Increasing on $(-\infty, 2]$.
Decreasing on $[2, \infty)$.


37. a. For $f(x) = x^2 + 2x - 8$, $a = 1$, $b = 2$, $c = -8$.
Since $a = 1 > 0$, the graph opens up.
The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-2}{2(1)} = \frac{-2}{2} = -1.$$

The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(-1) = (-1)^2 + 2(-1) - 8$$
$$= 1 - 2 - 8 = -9.$$

Thus, the vertex is $(-1, -9)$.
The axis of symmetry is the line $x = -1$.
The discriminant is:
$b^2 - 4ac = 2^2 - 4(1)(-8) = 4 + 32 = 36 > 0$,
so the graph has two x-intercepts.
The x-intercepts are found by solving:
$x^2 + 2x - 8 = 0$
$(x + 4)(x - 2) = 0$
$x = -4$ or $x = 2$.
The x-intercepts are -4 and 2.
The y-intercept is $f(0) = -8$.



b. The domain is $(-\infty, \infty)$.
The range is $[-9, \infty)$.

c. Decreasing on $(-\infty, -1]$.
Increasing on $[-1, \infty)$.

38. a. For $f(x) = x^2 - 2x - 3$, $a = 1$, $b = -2$, $c = -3$.
Since $a = 1 > 0$, the graph opens up.
The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-2)}{2(1)} = \frac{2}{2} = 1.$$

The y-coordinate of the vertex is


$$f\left(\frac{-b}{2a}\right) = f(1) = 1^2 - 2(1) - 3 = -4.$$

Thus, the vertex is $(1, -4)$.
The axis of symmetry is the line $x = 1$.
The discriminant is:
$b^2 - 4ac = (-2)^2 - 4(1)(-3) = 4 + 12 = 16 > 0$,

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

so the graph has two x-intercepts.
The x-intercepts are found by solving:
$x^2 - 2x - 3 = 0$
$(x+1)(x-3) = 0$
$x = -1$ or $x = 3$.
The x-intercepts are -1 and 3.
The y-intercept is $f(0) = -3$.



b. The domain is $(-\infty, \infty)$. The range is $[-4, \infty)$.

c. Decreasing on $(-\infty, 1]$. Increasing on $[1, \infty)$.

39. a. For $f(x) = x^2 + 2x + 1$, $a = 1$, $b = 2$, $c = 1$.
Since $a = 1 > 0$, the graph opens up.
The x-coordinate of the vertex is


$$x = \frac{-b}{2a} = \frac{-2}{2(1)} = \frac{-2}{2} = -1.$$

The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(-1)$$

$$= (-1)^2 + 2(-1) + 1 = 1 - 2 + 1 = 0.$$

Thus, the vertex is $(-1, 0)$.
The axis of symmetry is the line $x = -1$.
The discriminant is:
$b^2 - 4ac = 2^2 - 4(1)(1) = 4 - 4 = 0$,
so the graph has one x-intercept.
The x-intercept is found by solving:
$x^2 + 2x + 1 = 0$
$(x+1)^2 = 0$
$x = -1$.
The x-intercept is -1.
The y-intercept is $f(0) = 1$.



b. The domain is $(-\infty, \infty)$.
The range is $[0, \infty)$.

c. Decreasing on $(-\infty, -1]$.
Increasing on $[-1, \infty)$.

40. a. For $f(x) = x^2 + 6x + 9$, $a = 1$, $b = 6$, $c = 9$.
Since $a = 1 > 0$, the graph opens up.
The x-coordinate of the vertex is


$$x = \frac{-b}{2a} = \frac{-6}{2(1)} = \frac{-6}{2} = -3.$$

The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(-3)$$

$$= (-3)^2 + 6(-3) + 9 = 9 - 18 + 9 = 0.$$

Thus, the vertex is $(-3, 0)$.
The axis of symmetry is the line $x = -3$.
The discriminant is:
$b^2 - 4ac = 6^2 - 4(1)(9) = 36 - 36 = 0$,
so the graph has one x-intercept.
The x-intercept is found by solving:
$x^2 + 6x + 9 = 0$
$(x+3)^2 = 0$
$x = -3$.
The x-intercept is -3.
The y-intercept is $f(0) = 9$.




b. The domain is $(-\infty, \infty)$. The range is $[0, \infty)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

c. Decreasing on $(-\infty, -3]$.
Increasing on $[-3, \infty)$.


41. a. For $f(x) = 2x^2 - x + 2$, $a = 2$, $b = -1$, $c = 2$.
Since $a = 2 > 0$, the graph opens up.
The x-coordinate of the vertex is
$$x = \frac{-b}{2a} = \frac{-(-1)}{2(2)} = \frac{1}{4}.$$
The y-coordinate of the vertex is
$$f\left(\frac{-b}{2a}\right) = f\left(\frac{1}{4}\right) = 2\left(\frac{1}{4}\right)^2 - \frac{1}{4} + 2$$
$$= \frac{1}{8} - \frac{1}{4} + 2 = \frac{15}{8}.$$
Thus, the vertex is $\left(\frac{1}{4}, \frac{15}{8}\right)$.
The axis of symmetry is the line $x = \frac{1}{4}$.
The discriminant is:
$b^2 - 4ac = (-1)^2 - 4(2)(2) = 1 - 16 = -15$,
so the graph has no x-intercepts.
The y-intercept is $f(0) = 2$.



b. The domain is $(-\infty, \infty)$. The range is $\left[\frac{15}{8}, \infty\right)$.

c. Decreasing on $\left(-\infty, \frac{1}{4}\right]$. Increasing on $\left[\frac{1}{4}, \infty\right)$.

42. a. For $f(x) = 4x^2 - 2x + 1$, $a = 4$, $b = -2$, $c = 1$.
Since $a = 4 > 0$, the graph opens up.
The x-coordinate of the vertex is
$$x = \frac{-b}{2a} = \frac{-(-2)}{2(4)} = \frac{2}{8} = \frac{1}{4}.$$
The y-coordinate of the vertex is
$$f\left(\frac{-b}{2a}\right) = f\left(\frac{1}{4}\right) = 4\left(\frac{1}{4}\right)^2 - 2\left(\frac{1}{4}\right) + 1$$
$$= \frac{1}{4} - \frac{1}{2} + 1 = \frac{3}{4}.$$
Thus, the vertex is $\left(\frac{1}{4}, \frac{3}{4}\right)$.
The axis of symmetry is the line $x = \frac{1}{4}$.
The discriminant is:
$b^2 - 4ac = (-2)^2 - 4(4)(1) = 4 - 16 = -12$,
so the graph has no x-intercepts.
The y-intercept is $f(0) = 1$.



b. The domain is $(-\infty, \infty)$.
The range is $\left[\frac{3}{4}, \infty\right)$.


c. Decreasing on $\left(-\infty, \frac{1}{4}\right]$.
Increasing on $\left[\frac{1}{4}, \infty\right)$.

43. a. For $f(x) = -2x^2 + 2x - 3$, $a = -2$, $b = 2$, $c = -3$. Since $a = -2 < 0$, the graph opens down.
The x-coordinate of the vertex is
$$x = \frac{-b}{2a} = \frac{-(2)}{2(-2)} = \frac{-2}{-4} = \frac{1}{2}.$$
The y-coordinate of the vertex is
$$f\left(\frac{-b}{2a}\right) = f\left(\frac{1}{2}\right) = -2\left(\frac{1}{2}\right)^2 + 2\left(\frac{1}{2}\right) - 3$$
$$= -\frac{1}{2} + 1 - 3 = -\frac{5}{2}.$$
Thus, the vertex is $\left(\frac{1}{2}, -\frac{5}{2}\right)$.
The axis of symmetry is the line $x = \frac{1}{2}$.
The discriminant is:
$b^2 - 4ac = 2^2 - 4(-2)(-3) = 4 - 24 = -20$,
so the graph has no x-intercepts.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

The y-intercept is $f(0) = -3$.



b. The domain is $(-\infty, \infty)$.

The range is $\left(-\infty, -\frac{5}{2}\right]$.

c. Increasing on $\left(-\infty, \frac{1}{2}\right]$.

Decreasing on $\left[\frac{1}{2}, \infty\right)$.

44. **a.** For $f(x) = -3x^2 + 3x - 2$, $a = -3$, $b = 3$, $c = -2$. Since $a = -3 < 0$, the graph opens down.
The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-3}{2(-3)} = \frac{-3}{-6} = \frac{1}{2}.$$


The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f\left(\frac{1}{2}\right) = -3\left(\frac{1}{2}\right)^2 + 3\left(\frac{1}{2}\right) - 2$$
$$= -\frac{3}{4} + \frac{3}{2} - 2 = -\frac{5}{4}.$$

Thus, the vertex is $\left(\frac{1}{2}, -\frac{5}{4}\right)$.

The axis of symmetry is the line $x = \frac{1}{2}$.

The discriminant is:
$b^2 - 4ac = 3^2 - 4(-3)(-2) = 9 - 24 = -15$,
so the graph has no x-intercepts.
The y-intercept is $f(0) = -2$.



b. The domain is $(-\infty, \infty)$.

The range is $\left(-\infty, -\frac{5}{4}\right]$.

c. Increasing on $\left(-\infty, \frac{1}{2}\right]$.

Decreasing on $\left[\frac{1}{2}, \infty\right)$.

45. **a.** For $f(x) = 3x^2 + 6x + 2$, $a = 3$, $b = 6$, $c = 2$. Since $a = 3 > 0$, the graph opens up.
The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-6}{2(3)} = \frac{-6}{6} = -1.$$

The y-coordinate of the vertex is


$$f\left(\frac{-b}{2a}\right) = f(-1) = 3(-1)^2 + 6(-1) + 2$$
$$= 3 - 6 + 2 = -1.$$

Thus, the vertex is $(-1, -1)$.
The axis of symmetry is the line $x = -1$.
The discriminant is:
$b^2 - 4ac = 6^2 - 4(3)(2) = 36 - 24 = 12$,
so the graph has two x-intercepts.
The x-intercepts are found by solving:

$$3x^2 + 6x + 2 = 0$$
$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$
$$= \frac{-6 \pm \sqrt{12}}{6} = \frac{-6 \pm 2\sqrt{3}}{6} = \frac{-3 \pm \sqrt{3}}{3}$$

The x-intercepts are $-1 - \frac{\sqrt{3}}{3}$ and $-1 + \frac{\sqrt{3}}{3}$.

The y-intercept is $f(0) = 2$.



b. The domain is $(-\infty, \infty)$.

The range is $[-1, \infty)$.

c. Decreasing on $(-\infty, -1]$.

Increasing on $[-1, \infty)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

46. a. For $f(x) = 2x^2 + 5x + 3$, $a = 2$, $b = 5$, $c = 3$. Since $a = 2 > 0$, the graph opens up.
The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-5}{2(2)} = -\frac{5}{4}.$$

The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f\left(-\frac{5}{4}\right)$$
$$= 2\left(-\frac{5}{4}\right)^2 + 5\left(-\frac{5}{4}\right) + 3$$
$$= \frac{25}{8} - \frac{25}{4} + 3$$
$$= -\frac{1}{8}.$$

Thus, the vertex is $\left(-\frac{5}{4}, -\frac{1}{8}\right)$.


The axis of symmetry is the line $x = -\frac{5}{4}$.

The discriminant is:
$b^2 - 4ac = 5^2 - 4(2)(3) = 25 - 24 = 1$,
so the graph has two x-intercepts.
The x-intercepts are found by solving:

$$2x^2 + 5x + 3 = 0$$
$$(2x + 3)(x + 1) = 0$$
$$x = -\frac{3}{2} \text{ or } x = -1.$$

The x-intercepts are $-\frac{3}{2}$ and -1.

The y-intercept is $f(0) = 3$.



b. The domain is $(-\infty, \infty)$.

The range is $\left[-\frac{1}{8}, \infty\right)$.

c. Decreasing on $\left(-\infty, -\frac{5}{4}\right]$.

Increasing on $\left[-\frac{5}{4}, \infty\right)$.

47. a. For $f(x) = -4x^2 - 6x + 2$, $a = -4$, $b = -6$, $c = 2$. Since $a = -4 < 0$, the graph opens down.
The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-6)}{2(-4)} = \frac{6}{-8} = -\frac{3}{4}.$$

The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f\left(-\frac{3}{4}\right) = -4\left(-\frac{3}{4}\right)^2 - 6\left(-\frac{3}{4}\right) + 2$$
$$= -\frac{9}{4} + \frac{9}{2} + 2 = \frac{17}{4}.$$

Thus, the vertex is $\left(-\frac{3}{4}, \frac{17}{4}\right)$.


The axis of symmetry is the line $x = -\frac{3}{4}$.

The discriminant is:
$b^2 - 4ac = (-6)^2 - 4(-4)(2) = 36 + 32 = 68$,
so the graph has two x-intercepts.
The x-intercepts are found by solving:

$$-4x^2 - 6x + 2 = 0$$
$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-6) \pm \sqrt{68}}{2(-4)}$$
$$= \frac{6 \pm \sqrt{68}}{-8} = \frac{6 \pm 2\sqrt{17}}{-8} = \frac{3 \pm \sqrt{17}}{-4}$$

The x-intercepts are $\frac{-3 + \sqrt{17}}{4}$ and $\frac{-3 - \sqrt{17}}{4}$.

The y-intercept is $f(0) = 2$.



b. The domain is $(-\infty, \infty)$.

The range is $\left(-\infty, \frac{17}{4}\right]$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

c. Decreasing on $\left[-\frac{3}{4}, \infty\right)$.

Increasing on $\left(-\infty, -\frac{3}{4}\right]$.

48. a. For $f(x) = 3x^2 - 8x + 2,\ a = 3,\ b = -8,\ c = 2$.
Since $a = 3 > 0$, the graph opens up.
The x-coordinate of the vertex is
$x = \frac{-b}{2a} = \frac{-(-8)}{2(3)} = \frac{8}{6} = \frac{4}{3}$.
The y-coordinate of the vertex is


$$f\left(\frac{-b}{2a}\right) = f\left(\frac{4}{3}\right) = 3\left(\frac{4}{3}\right)^2 - 8\left(\frac{4}{3}\right) + 2$$
$$= \frac{16}{3} - \frac{32}{3} + 2 = -\frac{10}{3}.$$

Thus, the vertex is $\left(\frac{4}{3}, -\frac{10}{3}\right)$.

The axis of symmetry is the line $x = \frac{4}{3}$.
The discriminant is:
$b^2 - 4ac = (-8)^2 - 4(3)(2) = 64 - 24 = 40$,
so the graph has two x-intercepts.
The x-intercepts are found by solving:
$3x^2 - 8x + 2 = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-8) \pm \sqrt{40}}{2(3)}$$
$$= \frac{8 \pm \sqrt{40}}{6} = \frac{8 \pm 2\sqrt{10}}{6} = \frac{4 \pm \sqrt{10}}{3}$$

The x-intercepts are $\frac{4+\sqrt{10}}{3}$ and $\frac{4-\sqrt{10}}{3}$.
The y-intercept is $f(0) = 2$.



b. The domain is $(-\infty, \infty)$.

The range is $\left[-\frac{10}{3}, \infty\right)$.

c. Decreasing on $\left(-\infty, \frac{4}{3}\right]$.

Increasing on $\left[\frac{4}{3}, \infty\right)$.

49. Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(-1,-2)$ so we have $h = -1$ and $k = -2$. The graph also passes through the point $(x, y) = (0,-1)$. Substituting these values for *x, y, h,* and *k*, we can solve for *a*:

$$-1 = a(0-(-1))^2 + (-2)$$
$$-1 = a(1)^2 - 2$$
$$-1 = a - 2$$
$$1 = a$$

The quadratic function is
$f(x) = (x+1)^2 - 2 = x^2 + 2x - 1$.

50. Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(2,1)$ so we have $h = 2$ and $k = 1$. The graph also passes through the point $(x, y) = (0,5)$. Substituting these values for *x, y, h,* and *k*, we can solve for *a*:

$$5 = a(0-2)^2 + 1$$
$$5 = a(-2)^2 + 1$$
$$5 = 4a + 1$$
$$4 = 4a$$
$$1 = a$$

The quadratic function is
$f(x) = (x-2)^2 + 1 = x^2 - 4x + 5$.

51. Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(-3,5)$ so we have $h = -3$ and $k = 5$. The graph also passes through the point $(x, y) = (0,-4)$. Substituting these values for *x, y, h,* and *k*, we can solve for *a*:

$$-4 = a(0-(-3))^2 + 5$$
$$-4 = a(3)^2 + 5$$
$$-4 = 9a + 5$$
$$-9 = 9a$$
$$-1 = a$$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

The quadratic function is
$f(x) = -(x+3)^2 + 5 = -x^2 - 6x - 4$.

52. Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(2,3)$ so we have $h = 2$ and $k = 3$. The graph also passes through the point $(x, y) = (0, -1)$. Substituting these values for *x, y, h,* and *k*, we can solve for *a*:

$-1 = a(0-2)^2 + 3$

$-1 = a(-2)^2 + 3$

$-1 = 4a + 3$

$-4 = 4a$

$-1 = a$

The quadratic function is
$f(x) = -(x-2)^2 + 3 = -x^2 + 4x - 1$.

53. Consider the form $y = ax^2 + bx + c$. Substituting the three points from the graph into the general form we have the following three equations.

$5 = a(-1)^2 + b(-1) + c \Rightarrow 5 = a - b + c$

and

$5 = a(3)^2 + b(3) + c \Rightarrow 5 = 9a + 3b + c$

and

$-1 = a(0)^2 + b(0) + c \Rightarrow -1 = c$

Since $-1 = c$, we have the following equations:
$5 = a - b - 1, \quad 5 = 9a + 3b - 1, \quad -1 = c$
Solving the first two simultaneously we have

$$\left.\begin{array}{l} 5 = a - b - 1 \\ 5 = 9a + 3b - 1 \end{array}\right)$$

$$\left.\begin{array}{l} 6 = a - b \\ 6 = 9a + 3b \end{array}\right) \rightarrow \quad a = 2, b = -4$$

The quadratic function is $f(x) = 2x^2 - 4x - 1$.

54. Consider the form $y = ax^2 + bx + c$. Substituting the three points from the graph into the general form we have the following three equations.

$-2 = a(-4)^2 + b(-4) + c \Rightarrow -2 = 16a - 4b + c$

and

$4 = a(-1)^2 + b(-1) + c \Rightarrow 4 = a - b + c$

and

$-2 = a(0)^2 + b(0) + c \Rightarrow -2 = c$

Since $-2 = c$, we have the following equations:
$-2 = 16a - 4b - 2, \quad 4 = a - b - 2, \quad -2 = c$
Solving the first two simultaneously we have

$$\left.\begin{array}{l} -2 = 16a - 4b - 2 \\ 4 = a - b - 2 \end{array}\right)$$

$$\left.\begin{array}{l} 0 = 16a - 4b \\ 6 = a - b \end{array}\right) \rightarrow \quad a = -2, b = -8$$

The quadratic function is $f(x) = -2x^2 - 8x - 2$.

55. For $f(x) = 2x^2 + 12x, \quad a = 2, \quad b = 12, \quad c = 0$.
Since $a = 2 > 0,$ the graph opens up, so the vertex is a minimum point. The minimum occurs at $x = \dfrac{-b}{2a} = \dfrac{-12}{2(2)} = \dfrac{-12}{4} = -3.$
The minimum value is
$f(-3) = 2(-3)^2 + 12(-3) = 18 - 36 = -18$.

56. For $f(x) = -2x^2 + 12x, \quad a = -2, \quad b = 12, \quad c = 0,$.
Since $a = -2 < 0,$ the graph opens down, so the vertex is a maximum point. The maximum occurs at $x = \dfrac{-b}{2a} = \dfrac{-12}{2(-2)} = \dfrac{-12}{-4} = 3.$
The maximum value is
$f(3) = -2(3)^2 + 12(3) = -18 + 36 = 18$.

57. For $f(x) = 2x^2 + 12x - 3, \; a = 2, \; b = 12, \quad c = -3.$
Since $a = 2 > 0,$ the graph opens up, so the vertex is a minimum point. The minimum occurs at $x = \dfrac{-b}{2a} = \dfrac{-12}{2(2)} = \dfrac{-12}{4} = -3.$ The minimum value is
$f(-3) = 2(-3)^2 + 12(-3) - 3 = 18 - 36 - 3 = -21$.

58. For $f(x) = 4x^2 - 8x + 3, \quad a = 4, \quad b = -8, \quad c = 3.$
Since $a = 4 > 0,$ the graph opens up, so the vertex is a minimum point. The minimum occurs at $x = \dfrac{-b}{2a} = \dfrac{-(-8)}{2(4)} = \dfrac{8}{8} = 1.$ The minimum value is
$f(1) = 4(1)^2 - 8(1) + 3 = 4 - 8 + 3 = -1$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

59. For $f(x) = -x^2 + 10x - 4$, $a = -1$, $b = 10$, $c = -4$. Since $a = -1 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at $x = \dfrac{-b}{2a} = \dfrac{-10}{2(-1)} = \dfrac{-10}{-2} = 5$. The maximum value is

$f(5) = -(5)^2 + 10(5) - 4 = -25 + 50 - 4 = 21$.

60. For $f(x) = -2x^2 + 8x + 3$, $a = -2$, $b = 8$, $c = 3$. Since $a = -2 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at $x = \dfrac{-b}{2a} = \dfrac{-8}{2(-2)} = \dfrac{-8}{-4} = 2$. The maximum value is

$f(2) = -2(2)^2 + 8(2) + 3 = -8 + 16 + 3 = 11$.

61. For $f(x) = -3x^2 + 12x + 1$, $a = -3$, $b = 12$, $c = 1$. Since $a = -3 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at $x = \dfrac{-b}{2a} = \dfrac{-12}{2(-3)} = \dfrac{-12}{-6} = 2$. The maximum value is $f(2) = -3(2)^2 + 12(2) + 1 = -12 + 24 + 1 = 13$.


62. For $f(x) = 4x^2 - 4x$, $a = 4$, $b = -4$, $c = 0$. Since $a = 4 > 0$, the graph opens up, so the vertex is a minimum point. The minimum occurs at $x = \dfrac{-b}{2a} = \dfrac{-(-4)}{2(4)} = \dfrac{4}{8} = \dfrac{1}{2}$. The minimum value is

$$f\left(\frac{1}{2}\right) = 4\left(\frac{1}{2}\right)^2 - 4\left(\frac{1}{2}\right) = 1 - 2 = -1.$$

63. **a.** For $f(x) = x^2 - 2x - 15$, $a = 1$, $b = -2$, $c = -15$. Since $a = 1 > 0$, the graph opens up.
The x-coordinate of the vertex is
$x = \dfrac{-b}{2a} = \dfrac{-(-2)}{2(1)} = \dfrac{2}{2} = 1$.
The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(1) = (1)^2 - 2(1) - 15 = 1 - 2 - 15 = -16.$$

Thus, the vertex is $(1, -16)$.
The discriminant is:
$b^2 - 4ac = (-2)^2 - 4(1)(-15) = 4 + 60 = 64 > 0$,
so the graph has two x-intercepts.
The x-intercepts are found by solving:
$x^2 - 2x - 15 = 0$
$(x + 3)(x - 5) = 0$
$x = -3$ or $x = 5$
The x-intercepts are -3 and 5.
The y-intercept is $f(0) = -15$.



b. The domain is $(-\infty, \infty)$.
The range is $[-16, \infty)$.

c. Decreasing on $(-\infty, 1]$.
Increasing on $[1, \infty)$.


64. **a.** For $f(x) = x^2 - 2x - 8$, $a = 1$, $b = -2$, $c = -8$. Since $a = 1 > 0$, the graph opens up.
The x-coordinate of the vertex is
$x = \dfrac{-b}{2a} = \dfrac{-(-2)}{2(1)} = \dfrac{2}{2} = 1$.
The y-coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(1) = (1)^2 - 2(1) - 8 = 1 - 2 - 8 = -9.$$

Thus, the vertex is $(1, -9)$.
The discriminant is:
$b^2 - 4ac = (-2)^2 - 4(1)(-8) = 4 + 32 = 36 > 0$,
so the graph has two x-intercepts.
The x-intercepts are found by solving:
$x^2 - 2x - 8 = 0$
$(x + 2)(x - 4) = 0$
$x = -2$ or $x = 4$
The x-intercepts are -2 and 4.
The y-intercept is $f(0) = -8$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual



b. The domain is $(-\infty, \infty)$. The range is $[-9, \infty)$.

c. Decreasing on $(-\infty, 1]$. Increasing on $[1, \infty)$.


65. a. $F(x) = 2x - 5$ is a linear function.

The x-intercept is found by solving:

$$2x - 5 = 0$$
$$2x = 5$$
$$x = \frac{5}{2}$$

The x-intercept is $\frac{5}{2}$.

The y-intercept is $F(0) = -5$.



b. The domain is $(-\infty, \infty)$.

The range is $(-\infty, \infty)$.

c. Increasing on $(-\infty, \infty)$.


66. a. $f(x) = \frac{3}{2}x - 2$ is a linear function.

The x-intercept is found by solving:

$$\frac{3}{2}x - 2 = 0$$
$$\frac{3}{2}x = 2$$
$$x = 2 \cdot \frac{2}{3} = \frac{4}{3}$$

The x-intercept is $\frac{4}{3}$.

The y-intercept is $f(0) = -2$.




b. The domain is $(-\infty, \infty)$.

The range is $(-\infty, \infty)$.

c. Increasing on $(-\infty, \infty)$.

67. a. $g(x) = -2(x-3)^2 + 2$

Using the graph of $y = x^2$, shift right 3 units, reflect about the x-axis, stretch vertically by a factor of 2, then shift up 2 units.



b. The domain is $(-\infty, \infty)$.


The range is $(-\infty, 2]$.

c. Increasing on $(-\infty, 3]$.

Decreasing on $[3, \infty)$.

68. a. $h(x) = -3(x+1)^2 + 4$

Using the graph of $y = x^2$, shift left 1 unit, reflect about the x-axis, stretch vertically by a factor of 3, then shift up 4 units.




Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, 4]$.

c. Increasing on $(-\infty, -1]$.
Decreasing on $[-1, \infty)$.


69. a. For $f(x) = 2x^2 + x + 1$, $a = 2$, $b = 1$, $c = 1$.
Since $a = 2 > 0$, the graph opens up.
The x-coordinate of the vertex is
$$x = \frac{-b}{2a} = \frac{-1}{2(2)} = \frac{-1}{4} = -\frac{1}{4}.$$
The y-coordinate of the vertex is
$$f\left(\frac{-b}{2a}\right) = f\left(-\frac{1}{4}\right) = 2\left(-\frac{1}{4}\right)^2 + \left(-\frac{1}{4}\right) + 1$$
$$= \frac{1}{8} - \frac{1}{4} + 1 = \frac{7}{8}.$$
Thus, the vertex is $\left(-\frac{1}{4}, \frac{7}{8}\right)$.
The discriminant is:
$b^2 - 4ac = 1^2 - 4(2)(1) = 1 - 8 = -7$,
so the graph has no x-intercepts.
The y-intercept is $f(0) = 1$.



b. The domain is $(-\infty, \infty)$.
The range is $\left[\frac{7}{8}, \infty\right)$.

c. Decreasing on $\left(-\infty, -\frac{1}{4}\right]$.
Increasing on $\left[-\frac{1}{4}, \infty\right)$.

70. a. For $G(x) = 3x^2 + 2x + 5$, $a = 3$, $b = 2$, $c = 5$.
Since $a = 3 > 0$, the graph opens up.
The x-coordinate of the vertex is
$$x = \frac{-b}{2a} = \frac{-2}{2(3)} = \frac{-2}{6} = -\frac{1}{3}.$$
The y-coordinate of the vertex is
$$G\left(\frac{-b}{2a}\right) = G\left(-\frac{1}{3}\right) = 3\left(-\frac{1}{3}\right)^2 + 2\left(-\frac{1}{3}\right) + 5$$
$$= \frac{1}{3} - \frac{2}{3} + 5 = \frac{14}{3}.$$
Thus, the vertex is $\left(-\frac{1}{3}, \frac{14}{3}\right)$.
The discriminant is:
$b^2 - 4ac = 2^2 - 4(3)(5) = 4 - 60 = -56$,
so the graph has no x-intercepts.
The y-intercept is $G(0) = 5$.




b. The domain is $(-\infty, \infty)$.
The range is $\left[\frac{14}{3}, \infty\right)$.

c. Decreasing on $\left(-\infty, -\frac{1}{3}\right]$.
Increasing on $\left[-\frac{1}{3}, \infty\right)$.

71. a. $h(x) = -\frac{2}{5}x + 4$ is a linear function.
The x-intercept is found by solving:
$$-\frac{2}{5}x + 4 = 0$$
$$-\frac{2}{5}x = -4$$
$$x = -4\left(-\frac{5}{2}\right) = 10$$
The x-intercept is 10.
The y-intercept is $h(0) = 4$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual




b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, \infty)$.

c. Decreasing on $(-\infty, \infty)$.

72. a. $f(x) = -3x + 2$ is a linear function.
The x-intercept is found by solving:

$$-3x + 2 = 0$$
$$-3x = -2$$
$$x = \frac{-2}{-3} = \frac{2}{3}$$

The x-intercept is $\frac{2}{3}$.
The y-intercept is $f(0) = 2$.



b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, \infty)$.

c. Decreasing on $(-\infty, \infty)$.

73. a. For $H(x) = -4x^2 - 4x - 1$, $a = -4$, $b = -4$, $c = -1$. Since $a = -4 < 0$, the graph opens down. The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-4)}{2(-4)} = \frac{4}{-8} = -\frac{1}{2}.$$

The y-coordinate of the vertex is


$$H\left(\frac{-b}{2a}\right) = H\left(-\frac{1}{2}\right) = -4\left(-\frac{1}{2}\right)^2 - 4\left(-\frac{1}{2}\right) - 1$$
$$= -1 + 2 - 1 = 0$$

Thus, the vertex is $\left(-\frac{1}{2}, 0\right)$.

The discriminant is:
$b^2 - 4ac = (-4)^2 - 4(-4)(-1) = 16 - 16 = 0$,
so the graph has one x-intercept.
The x-intercept is found by solving:

$$-4x^2 - 4x - 1 = 0$$
$$4x^2 + 4x + 1 = 0$$
$$(2x + 1)^2 = 0$$
$$2x + 1 = 0$$
$$x = -\frac{1}{2}$$

The x-intercept is $-\frac{1}{2}$.
The y-intercept is $H(0) = -1$.



b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, 0]$.

c. Increasing on $\left(-\infty, -\frac{1}{2}\right]$.
Decreasing on $\left[-\frac{1}{2}, \infty\right)$.

74. a. For $F(x) = -4x^2 + 20x - 25$, $a = -4$, $b = 20$, $c = -25$. Since $a = -4 < 0$, the graph opens down. The x-coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-20}{2(-4)} = \frac{-20}{-8} = \frac{5}{2}.$$

The y-coordinate of the vertex is

$$F\left(\frac{-b}{2a}\right) = F\left(\frac{5}{2}\right) = -4\left(\frac{5}{2}\right)^2 + 20\left(\frac{5}{2}\right) - 25$$
$$= -25 + 50 - 25 = 0$$

Thus, the vertex is $\left(\frac{5}{2}, 0\right)$.

The discriminant is:

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


$$b^2 - 4ac = (20)^2 - 4(-4)(-25)$$
$$= 400 - 400 = 0,$$

so the graph has one x-intercept.
The x-intercept is found by solving:

$$-4x^2 + 20x - 25 = 0$$
$$4x^2 - 20x + 25 = 0$$
$$(2x-5)^2 = 0$$
$$2x - 5 = 0$$
$$x = \frac{5}{2}$$

The x-intercept is $\frac{5}{2}$.

The y-intercept is $F(0) = -25$.



b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, 0]$.

c. Increasing on $\left(-\infty, \frac{5}{2}\right]$.
Decreasing on $\left[\frac{5}{2}, \infty\right)$.

75. Use the form $f(x) = a(x-h)^2 + k$.
The vertex is $(0,2)$, so $h = 0$ and $k = 2$.
$f(x) = a(x-0)^2 + 2 = ax^2 + 2$.
Since the graph passes through $(1, 8)$, $f(1) = 8$.

$$f(x) = ax^2 + 2$$
$$8 = a(1)^2 + 2$$
$$8 = a + 2$$
$$6 = a$$


$f(x) = 6x^2 + 2$.
$a = 6, b = 0, c = 2$

76. Use the form $f(x) = a(x-h)^2 + k$.
The vertex is $(1, 4)$, so $h = 1$ and $k = 4$.
$f(x) = a(x-1)^2 + 4$.
Since the graph passes through $(-1, -8)$,
$f(-1) = -8$.

$$-8 = a(-1-1)^2 + 4$$
$$-8 = a(-2)^2 + 4$$
$$-8 = 4a + 4$$
$$-12 = 4a$$
$$-3 = a$$
$$f(x) = -3(x-1)^2 + 4$$
$$= -3(x^2 - 2x + 1) + 4$$
$$= -3x^2 + 6x - 3 + 4$$
$$= -3x^2 + 6x + 1$$

$a = -3, b = 6, c = 1$

77. **a** and **d.**



b.
$$f(x) = g(x)$$
$$2x - 1 = x^2 - 4$$
$$0 = x^2 - 2x - 3$$
$$0 = (x+1)(x-3)$$
$$x + 1 = 0 \quad \text{or} \quad x - 3 = 0$$
$$x = -1 \qquad x = 3$$

The solution set is $\{-1, 3\}$.


c.
$$f(-1) = 2(-1) - 1 = -2 - 1 = -3$$
$$g(-1) = (-1)^2 - 4 = 1 - 4 = -3$$
$$f(3) = 2(3) - 1 = 6 - 1 = 5$$
$$g(3) = (3)^2 - 4 = 9 - 4 = 5$$

Thus, the graphs of f and g intersect at the points $(-1, -3)$ and $(3, 5)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

78. **a** and **d.**



b. $f(x) = g(x)$

$-2x - 1 = x^2 - 9$

$0 = x^2 + 2x - 8$

$0 = (x + 4)(x - 2)$

$x + 4 = 0$ or $x - 2 = 0$

$x = -4$ $\quad x = 2$

The solution set is $\{-4, 2\}$.

c. $f(-4) = -2(-4) - 1 = 8 - 1 = 7$


$g(-4) = (-4)^2 - 9 = 16 - 9 = 7$

$f(2) = -2(2) - 1 = -4 - 1 = -5$

$g(2) = (2)^2 - 9 = 4 - 9 = -5$

Thus, the graphs of f and g intersect at the points $(-4,\ 7)$ and $(2, -5)$.

79. **a** and **d.**



b. $f(x) = g(x)$

$-x^2 + 4 = -2x + 1$

$0 = x^2 - 2x - 3$

$0 = (x + 1)(x - 3)$

$x + 1 = 0$ or $x - 3 = 0$

$x = -1$ $\quad x = 3$

The solution set is $\{-1, 3\}$.

c. $f(1) = -(-1)^2 + 4 = -1 + 4 = 3$


$g(1) = -2(-1) + 1 = 2 + 1 = 3$

$f(3) = -(3)^2 + 4 = -9 + 4 = -5$

$g(3) = -2(3) + 1 = -6 + 1 = -5$

Thus, the graphs of f and g intersect at the points $(-1,\ 3)$ and $(3, -5)$.

80. **a** and **d.**



b. $f(x) = g(x)$

$-x^2 + 9 = 2x + 1$

$0 = x^2 + 2x - 8$

$0 = (x + 4)(x - 2)$

$x + 4 = 0$ or $x - 2 = 0$

$x = -4$ $\quad x = 2$

The solution set is $\{-4, 2\}$.

c. $f(-4) = -(-4)^2 + 9 = -16 + 9 = -7$

$g(-4) = 2(-4) + 1 = -8 + 1 = -7$

$f(2) = -(2)^2 + 9 = -4 + 9 = 5$


$g(2) = 2(2) + 1 = 4 + 1 = 5$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Thus, the graphs of f and g intersect at the points $(-4,-7)$ and $(2,\ 5)$.

81. **a** and **d.**



b. $f(x)=g(x)$

$$-x^2+5x=x^2+3x-4$$
$$0=2x^2-2x-4$$
$$0=x^2-x-2$$
$$0=(x+1)(x-2)$$
$$x+1=0 \quad \text{or} \quad x-2=0$$
$$x=-1 \qquad\qquad x=2$$

The solution set is $\{-1, 2\}$.

c. $f(-1)=-(-1)^2+5(-1)=-1-5=-6$


$g(-1)=(-1)^2+3(-1)-4=1-3-4=-6$

$f(2)=-(2)^2+5(2)=-4+10=6$

$g(2)=2^2+3(2)-4=4+6-4=6$

Thus, the graphs of f and g intersect at the points $(-1,-6)$ and $(2,\ 6)$.

82. **a** and **d.**



b. $f(x)=g(x)$

$$-x^2+7x-6=x^2+x-6$$
$$0=2x^2-6x$$
$$0=2x(x-3)$$
$$2x=0 \quad \text{or} \quad x-3=0$$
$$x=0 \qquad\qquad x=3$$

The solution set is $\{0, 3\}$.

c. $f(0)=-(0)^2+7(0)-6=-6$

$g(0)=0^2+0-6=-6$

$f(3)=-(3)^2+7(3)-6=-9+21-6=6$

$g(3)=3^2+3-6=9+3-6=6$

Thus, the graphs of f and g intersect at the points $(0,-6)$ and $(3,\ 6)$.

83. **a.** For $a=1$:

$$f(x)=a(x-r_1)(x-r_2)$$
$$=1(x-(-3))(x-1)$$
$$=(x+3)(x-1)=x^2+2x-3$$

For $a=2$:

$$f(x)=2(x-(-3))(x-1)$$
$$=2(x+3)(x-1)$$
$$=2(x^2+2x-3)=2x^2+4x-6$$

For $a=-2$:

$$f(x)=-2(x-(-3))(x-1)$$
$$=-2(x+3)(x-1)$$
$$=-2(x^2+2x-3)=-2x^2-4x+6$$

For $a=5$:

$$f(x)=5(x-(-3))(x-1)$$
$$=5(x+3)(x-1)$$
$$=5(x^2+2x-3)=5x^2+10x-15$$

b. The x-intercepts are not affected by the value of a. The y-intercept is multiplied by the value of a.

c. The axis of symmetry is unaffected by the value of a. For this problem, the axis of symmetry is $x=-1$ for all values of a.

d. The x-coordinate of the vertex is not affected by the value of a. The y-coordinate of the vertex is multiplied by the value of a.

e. The x-coordinate of the vertex is the mean of the x-intercepts.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

84. a. For $a = 1$:

$$f(x) = 1(x-(-5))(x-3)$$
$$= (x+5)(x-3) = x^2 + 2x - 15$$

For $a = 2$:

$$f(x) = 2(x-(-5))(x-3)$$
$$= 2(x+5)(x-3)$$
$$= 2(x^2 + 2x - 15) = 2x^2 + 4x - 30$$

For $a = -2$:

$$f(x) = -2(x-(-5))(x-3)$$
$$= -2(x+5)(x-3)$$
$$= -2(x^2 + 2x - 15) = -2x^2 - 4x + 30$$

For $a = 5$:

$$f(x) = 5(x-(-5))(x-3)$$
$$= 5(x+5)(x-3)$$
$$= 5(x^2 + 2x - 15) = 5x^2 + 10x - 75$$

b. The x-intercepts are not affected by the value of a. The y-intercept is multiplied by the value of a.

c. The axis of symmetry is unaffected by the value of a. For this problem, the axis of symmetry is $x = -1$ for all values of a.

d. The x-coordinate of the vertex is not affected by the value of a. The y-coordinate of the vertex is multiplied by the value of a.

e. The x-coordinate of the vertex is the mean of the x-intercepts.

85. a. $x = -\dfrac{b}{2a} = -\dfrac{4}{2(1)} = -2$

$y = f(-2) = (-2)^2 + 4(-2) - 21 = -25$

The vertex is $(-2, -25)$.

b. $f(x) = 0$

$x^2 + 4x - 21 = 0$

$(x+7)(x-3) = 0$

$x + 7 = 0$ or $x - 3 = 0$

$x = -7$ $\quad$ $x = 3$

The x-intercepts of f are $(-7, 0)$ and $(3, 0)$.

c. $f(x) = -21$

$x^2 + 4x - 21 = -21$

$x^2 + 4x = 0$

$x(x+4) = 0$


$x = 0$ or $x + 4 = 0$

$x = -4$

The solutions $f(x) = -21$ are -4 and 0.

Thus, the points $(-4, -21)$ and $(0, -21)$ are on the graph of f.

d.



86. a. $x = -\dfrac{b}{2a} = -\dfrac{2}{2(1)} = -1$

$y = f(-1) = (-1)^2 + 2(-1) - 8 = -9$

The vertex is $(-1, -9)$.

b. $f(x) = 0$

$x^2 + 2x - 8 = 0$

$(x+4)(x-2) = 0$

$x + 4 = 0$ or $x - 2 = 0$

$x = -4$ $\quad$ $x = 2$

The x-intercepts of f are $(-4, 0)$ and $(2, 0)$.

c. $f(x) = -8$

$x^2 + 2x - 8 = -8$

$x^2 + 2x = 0$

$x(x+2) = 0$

$x = 0$ or $x + 2 = 0$

$x = -2$


The solutions $f(x) = -8$ are -2 and 0. Thus,

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

the points $(-2,-8)$ and $(0,-8)$ are on the graph of f.

d.



87. Let (x, y) represent a point on the line $y = x$. Then the distance from (x, y) to the point (3, 1) is $d = \sqrt{(x-3)^2 + (y-1)^2}$. Since $y = x$, we can replace the y variable with x so that we have the distance expressed as a function of x:

$$\begin{aligned} d(x) &= \sqrt{(x-3)^2 + (x-1)^2} \\ &= \sqrt{x^2 - 6x + 9 + x^2 - 2x + 1} \\ &= \sqrt{2x^2 - 8x + 10} \end{aligned}$$

Squaring both sides of this function, we obtain $[d(x)]^2 = 2x^2 - 8x + 10$.

Now, the expression on the right is quadratic. Since $a = 2 > 0$, it has a minimum. Finding the x-coordinate of the minimum point of $[d(x)]^2$ will also give us the x-coordinate of the minimum of $d(x)$: $x = \dfrac{-b}{2a} = \dfrac{-(-8)}{2(2)} = \dfrac{8}{4} = 2$. So, 2 is the x-coordinate of the point on the line $y = x$ that is closest to the point (3, 1). Since $y = x$, the y-coordinate is also 2. Thus, the point is (2, 2) is the point on the line $y = x$ that is closest to (3, 1).

88. Let (x, y) represent a point on the line $y = x + 1$. Then the distance from (x, y) to the point (4, 1) is $d = \sqrt{(x-4)^2 + (y-1)^2}$. Replacing the y variable with $x + 1$, we find the distance expressed as a function of x:

$$\begin{aligned} d(x) &= \sqrt{(x-4)^2 + ((x+1)-1)^2} \\ &= \sqrt{x^2 - 8x + 16 + x^2} \\ &= \sqrt{2x^2 - 8x + 16} \end{aligned}$$

Squaring both sides of this function, we obtain $[d(x)]^2 = 2x^2 - 8x + 16$.

Now, the expression on the right is quadratic. Since $a = 2 > 0$, it has a minimum. Finding the x-coordinate of the minimum point of $[d(x)]^2$ will also give us the x-coordinate of the minimum of $d(x)$: $x = \dfrac{-b}{2a} = \dfrac{-(-8)}{2(2)} = \dfrac{8}{4} = 2$. So, 2 is the x-coordinate of the point on the line $y = x + 1$ that is closest to the point (4, 1). The y-coordinate is $y = 2 + 1 = 3$. Thus, the point is (2, 3) is the point on the line $y = x + 1$ that is closest to (4, 1).

89. $R(p) = -4p^2 + 4000p$, $a = -4, b = 4000, c = 0$.

Since $a = -4 < 0$ the graph is a parabola that opens down, so the vertex is a maximum point. The maximum occurs at $p = \dfrac{-b}{2a} = \dfrac{-4000}{2(-4)} = 500$.

Thus, the unit price should be \$500 for maximum revenue. The maximum revenue is

$$\begin{aligned} R(500) &= -4(500)^2 + 4000(500) \\ &= -1000000 + 2000000 \\ &= \$1,000,000 \end{aligned}$$

90. $R(p) = -\dfrac{1}{2}p^2 + 1900p$, $a = -\dfrac{1}{2}, b = 1900, c = 0$.

Since $a = -\dfrac{1}{2} < 0$, the graph is a parabola that opens down, so the vertex is a maximum point. The maximum occurs at $p = \dfrac{-b}{2a} = \dfrac{-1900}{2(-1/2)} = \dfrac{-1900}{-1} = 1900$. Thus, the unit price should be \$1900 for maximum revenue. The maximum revenue is

$$\begin{aligned} R(1900) &= -\frac{1}{2}(1900)^2 + 1900(1900) \\ &= -1805000 + 3610000 \\ &= \$1,805,000 \end{aligned}$$

91. a. $C(x) = x^2 - 140x + 7400$, $a = 1, b = -140, c = 7400$. Since $a = 1 > 0$, the graph opens up, so the vertex is a minimum point. The minimum marginal cost occurs at $x = \dfrac{-b}{2a} = \dfrac{-(-140)}{2(1)} = \dfrac{140}{2} = 70$, 70,000 digital music players produced.

b. The minimum marginal cost is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) = f(70) &= (70)^2 - 140(70) + 7400 \\ &= 4900 - 9800 + 7400 \\ &= \$2500 \end{aligned}$$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

92. a. $C(x) = 5x^2 - 200x + 4000$,
$a = 5, b = -200, c = 4000$. Since $a = 5 > 0$, the graph opens up, so the vertex is a minimum point. The minimum marginal cost occurs at $x = \frac{-b}{2a} = \frac{-(-200)}{2(5)} = \frac{200}{10} = 20$,
20,000 thousand smartphones manufactured.

b. The minimum marginal cost is
$$f\left(\frac{-b}{2a}\right) = f(20) = 5(20)^2 - 200(20) + 4000$$
$$= 2000 - 4000 + 4000$$
$$= \$2000$$

93. a. $R(x) = 75x - 0.2x^2$
$a = -0.2, b = 75, c = 0$
The maximum revenue occurs when
$$x = \frac{-b}{2a} = \frac{-75}{2(-0.2)} = \frac{-75}{-0.4} = 187.5$$
The maximum revenue occurs when $x = 187$ or $x = 188$ watches.
The maximum revenue is:
$$R(187) = 75(187) - 0.2(187)^2 = \$7031.20$$
$$R(188) = 75(188) - 0.2(188)^2 = \$7031.20$$

b. $P(x) = R(x) - C(x)$
$$= 75x - 0.2x^2 - (32x + 1750)$$
$$= -0.2x^2 + 43x - 1750$$

c. $P(x) = -0.2x^2 + 43x - 1750$
$a = -0.2, b = 43, c = -1750$
$$x = \frac{-b}{2a} = \frac{-43}{2(-0.2)} = \frac{-43}{-0.4} = 107.5$$
The maximum profit occurs when $x = 107$ or $x = 108$ watches.
The maximum profit is:
$$P(107) = -0.2(107)^2 + 43(107) - 1750$$
$$= \$561.20$$
$$P(108) = -0.2(108)^2 + 43(108) - 1750$$
$$= \$561.20$$

d. Answers will vary.

94. a. $R(x) = 9.5x - 0.04x^2$
$a = -0.04, b = 9.5, c = 0$
The maximum revenue occurs when
$$x = \frac{-b}{2a} = \frac{-9.5}{2(-0.04)} = \frac{-9.5}{-0.08}$$
$$= 118.75 \approx 119 \text{ boxes of candy}$$
The maximum revenue is:
$$R(119) = 9.5(119) - 0.04(119)^2 = \$564.06$$

b. $P(x) = R(x) - C(x)$
$$= 9.5x - 0.04x^2 - (1.25x + 250)$$
$$= -0.04x^2 + 8.25x - 250$$

c. $P(x) = -0.04x^2 + 8.25x - 250$
$a = -0.04, b = 8.25, c = -250$
The maximum profit occurs when
$$x = \frac{-b}{2a} = \frac{-8.25}{2(-0.04)} = \frac{-8.25}{-0.08}$$
$$= 103.125 \approx 103 \text{ boxes of candy}$$
The maximum profit is:
$$P(103) = -0.04(103)^2 + 8.25(103) - 250$$
$$= \$175.39$$

d. Answers will vary.

95. a. $d(v) = 1.1v + 0.06v^2$
$$d(45) = 1.1(45) + 0.06(45)^2$$
$$= 49.5 + 121.5 = 171 \text{ ft.}$$

b. $200 = 1.1v + 0.06v^2$
$$0 = -200 + 1.1v + 0.06v^2$$
$$x = \frac{-(1.1) \pm \sqrt{(1.1)^2 - 4(0.06)(-200)}}{2(0.06)}$$
$$= \frac{-1.1 \pm \sqrt{49.21}}{0.12}$$
$$\approx \frac{-1.1 \pm 7.015}{0.12}$$
$v \approx 49$ or $v \approx -68$
Disregard the negative value since we are talking about speed. So the maximum speed you can be traveling would be approximately 49 mph.

c. The 1.1v term might represent the reaction time.

96. a. $a = \frac{-b}{2a} = \frac{-19.09}{2(-0.34)} = \frac{-19.09}{-0.68} = 28.1$ years old

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


b. $B(28.1) = -0.34(28.1)^2 + 19.09(28.1) - 203.98$
≈ 63.98 births per 1000 unmarried women

c. $B(40) = -0.34(40)^2 + 19.09(40) - 203.98$
$= 15.62$ births/1000 unmarried women over 40

97. If x is even, then ax^2 and bx are even. When two even numbers are added to an odd number the result is odd. Thus, $f(x)$ is odd. If x is odd, then ax^2 and bx are odd. The sum of three odd numbers is an odd number. Thus, $f(x)$ is odd.


98. Answers will vary.

99. $y = x^2 + 2x - 3$; $y = x^2 + 2x + 1$; $y = x^2 + 2x$



Each member of this family will be a parabola with the following characteristics:
(i) opens upwards since a > 0;
(ii) vertex occurs at $x = -\frac{b}{2a} = -\frac{2}{2(1)} = -1$;
(iii) There is at least one x-intercept since $b^2 - 4ac \geq 0$.

100. $y = x^2 - 4x + 1$; $y = x^2 + 1$; $y = x^2 + 4x + 1$



Each member of this family will be a parabola with the following characteristics:
(i) opens upwards since $a > 0$
(ii) y-intercept occurs at (0, 1).

101. The graph of the quadratic function $f(x) = ax^2 + bx + c$ will not have any x-intercepts whenever $b^2 - 4ac < 0$.

102. By completing the square on the quadratic function $f(x) = ax^2 + bx + c$ we obtain the equation $y = a\left(x + \frac{b}{2a}\right)^2 + c - \frac{b^2}{4a}$. We can then draw the graph by applying transformations to the graph of the basic parabola $y = x^2$, which opens up. When $a > 0$, the basic parabola will either be stretched or compressed vertically. When $a < 0$, the basic parabola will either be stretched or compressed vertically as well as reflected across the x-axis. Therefore, when $a > 0$, the graph of $f(x) = ax^2 + bx + c$ will open up, and when $a < 0$, the graph of $f(x) = ax^2 + bx + c$ will open down.

103. No. We know that the graph of a quadratic function $f(x) = ax^2 + bx + c$ is a parabola with vertex $\left(-\frac{b}{2a}, f\left(-\frac{b}{2a}\right)\right)$. If $a > 0$, then the vertex is a minimum point, so the range is $\left[f\left(-\frac{b}{2a}\right), \infty\right)$. If $a < 0$, then the vertex is a maximum point, so the range is $\left(-\infty, f\left(-\frac{b}{2a}\right)\right]$. Therefore, it is impossible for the range to be $(-\infty, \infty)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

104. Two quadratic functions can intersect 0, 1, or 2 times.

105. $x^2 + 4y^2 = 16$

To check for symmetry with respect to the x-axis, replace y with –y and see if the equations are equivalent.

$x^2 + 4(-y)^2 = 16$

$x^2 + 4y^2 = 16$

So the graph is symmetric with respect to the x-axis.

To check for symmetry with respect to the y-axis, replace x with –x and see if the equations are equivalent.

$(-x)^2 + 4y^2 = 16$

$x^2 + 4y^2 = 16$

So the graph is symmetric with respect to the y-axis.

To check for symmetry with respect to the origin, replace x with –x and y with –y and see if the equations are equivalent.

$(-x)^2 + 4(-y)^2 = 16$

$x^2 + 4y^2 = 16$

So the graph is symmetric with respect to the origin.

106. The radicand must be non-negative, so

$8 - 2x \ge 0$

$-2x \ge -8$

$x \le 4$

So the solution set is: $(-\infty,\ 4]$ or $\{x \mid x \le 4\}$.

107. $x^2 + y^2 - 10x + 4y + 20 = 0$

$x^2 - 10x + y^2 + 4y = -20$

$(x^2 - 10x + 25) + (y^2 + 4y + 4) = -20 + 25 + 4$

$(x-5)^2 + (y+2)^2 = 3^2$

Center: (5, -2); Radius = 3

108. To reflect a graph about the y-axis, we change $f(x)$ to $f(-x)$ so to reflect $y = \sqrt{x}$ about the y-axis we change it to $y = \sqrt{-x}$.

Section 2.5


1. $-3x - 2 < 7$

$-3x < 9$

$x > -3$

The solution set is $\{x \mid x > -3\}$ or $(-3, \infty)$.

2. $(-2, 7]$ represents the numbers between -2 and 7, including 7 but not including -2. Using inequality notation, this is written as $-2 < x \le 7$.



3. a. $f(x) > 0$ when the graph of f is above the x-axis. Thus, $\{x \mid x < -2 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -2) \cup (2, \infty)$.

b. $f(x) \le 0$ when the graph of f is below or intersects the x-axis. Thus, $\{x \mid -2 \le x \le 2\}$ or, using interval notation, $[-2,\ 2]$.

4. a. $g(x) < 0$ when the graph of g is below the x-axis. Thus, $\{x \mid x < -1 \text{ or } x > 4\}$ or, using interval notation, $(-\infty, -1) \cup (4, \infty)$.

b. $g(x) \ge 0$ when the graph of f is above or intersects the x-axis. Thus, $\{x \mid -1 \le x \le 4\}$ or, using interval notation, $[-1,\ 4]$.

5. a. $g(x) \ge f(x)$ when the graph of g is above or intersects the graph of f. Thus $\{x \mid -2 \le x \le 1\}$ or, using interval notation, $[-2,\ 1]$.

b. $f(x) > g(x)$ when the graph of f is above the graph of g. Thus, $\{x \mid x < -2 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, -2) \cup (1, \infty)$.

6. a. $f(x) < g(x)$ when the graph of f is below the graph of g. Thus, $\{x \mid x < -3 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, -3) \cup (1, \infty)$.

b. $f(x) \ge g(x)$ when the graph of f is above or intersects the graph of g. Thus,

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$\{x|-3 \le x \le 1\}$ or, using interval notation, $[-3,\ 1]$.

7. $x^2 - 3x - 10 < 0$

We graph the function $f(x) = x^2 - 3x - 10$. The intercepts are


y-intercept: $f(0) = -10$

x-intercepts: $x^2 - 3x - 10 = 0$

$(x-5)(x+2) = 0$

$x = 5,\ x = -2$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-3)}{2(1)} = \dfrac{3}{2}$. Since $f\left(\dfrac{3}{2}\right) = -\dfrac{49}{4}$, the vertex is $\left(\dfrac{3}{2}, -\dfrac{49}{4}\right)$.



The graph is below the x-axis for $-2 < x < 5$. Since the inequality is strict, the solution set is $\{x|-2 < x < 5\}$ or, using interval notation, $(-2,\ 5)$.

8. $x^2 + 3x - 10 > 0$

We graph the function $f(x) = x^2 + 3x - 10$. The intercepts are


y-intercept: $f(0) = -10$

x-intercepts: $x^2 + 3x - 10 = 0$

$(x+5)(x-2) = 0$

$x = -5,\ x = 2$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(3)}{2(1)} = -\dfrac{3}{2}$. Since $f\left(-\dfrac{3}{2}\right) = -\dfrac{49}{4}$, the vertex is $\left(-\dfrac{3}{2}, -\dfrac{49}{4}\right)$.



The graph is above the x-axis when $x < -5$ or $x > 2$. Since the inequality is strict, the solution set is $\{x|x < -5 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -5) \cup (2,\ \infty)$.

9. $x^2 - 4x > 0$

We graph the function $f(x) = x^2 - 4x$. The intercepts are


y-intercept: $f(0) = 0$

x-intercepts: $x^2 - 4x = 0$

$x(x-4) = 0$

$x = 0,\ x = 4$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-4)}{2(1)} = \dfrac{4}{2} = 2$. Since $f(2) = -4$, the vertex is $(2, -4)$.



The graph is above the x-axis when $x < 0$ or $x > 4$. Since the inequality is strict, the solution set is $\{x|x < 0 \text{ or } x > 4\}$ or, using interval notation, $(-\infty,\ 0) \cup (4,\ \infty)$.

10. $x^2 + 8x > 0$

We graph the function $f(x) = x^2 + 8x$. The intercepts are

y-intercept: $f(0) = 0$

x-intercepts: $x^2 + 8x = 0$

$x(x+8) = 0$


$x = 0,\ x = -8$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

The vertex is at $x = \frac{-b}{2a} = \frac{-(8)}{2(1)} = \frac{-8}{2} = -4$.

Since $f(-4) = -16$, the vertex is $(-4, -16)$.



The graph is above the x-axis when $x < -8$ or $x > 0$. Since the inequality is strict, the solution set is $\{x \mid x < -8 \text{ or } x > 0\}$ or, using interval notation, $(-\infty, -8) \cup (0, \infty)$.

11. $x^2 - 9 < 0$

We graph the function $f(x) = x^2 - 9$. The intercepts are


y-intercept: $f(0) = -9$

x-intercepts: $x^2 - 9 = 0$

$(x+3)(x-3) = 0$

$x = -3, x = 3$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(1)} = 0$. Since $f(0) = -9$, the vertex is $(0, -9)$.



The graph is below the x-axis when $-3 < x < 3$. Since the inequality is strict, the solution set is $\{x \mid -3 < x < 3\}$ or, using interval notation, $(-3, 3)$.

12. $x^2 - 1 < 0$

We graph the function $f(x) = x^2 - 1$. The intercepts are


y-intercept: $f(0) = -1$

x-intercepts: $x^2 - 1 = 0$

$(x+1)(x-1) = 0$

$x = -1, x = 1$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(1)} = 0$. Since $f(0) = -1$, the vertex is $(0, -1)$.



The graph is below the x-axis when $-1 < x < 1$. Since the inequality is strict, the solution set is $\{x \mid -1 < x < 1\}$ or, using interval notation, $(-1, 1)$.

13. $x^2 + x > 12$

$x^2 + x - 12 > 0$

We graph the function $f(x) = x^2 + x - 12$.


y-intercept: $f(0) = -12$

x-intercepts: $x^2 + x - 12 = 0$

$(x+4)(x-3) = 0$

$x = -4, x = 3$

The vertex is at $x = \frac{-b}{2a} = \frac{-(1)}{2(1)} = -\frac{1}{2}$. Since $f\left(-\frac{1}{2}\right) = -\frac{49}{4}$, the vertex is $\left(-\frac{1}{2}, -\frac{49}{4}\right)$.



The graph is above the x-axis when $x < -4$ or $x > 3$. Since the inequality is strict, the solution set is $\{x \mid x < -4 \text{ or } x > 3\}$ or, using interval notation, $(-\infty, -4) \cup (3, \infty)$.

14. $x^2 + 7x < -12$

$x^2 + 7x + 12 < 0$

We graph the function $f(x) = x^2 + 7x + 12$.

y-intercept: $f(0) = 12$

x-intercepts: $x^2 + 7x + 12 = 0$

$(x+4)(x+3) = 0$


$x = -4, x = -3$

The vertex is at $x = \frac{-b}{2a} = \frac{-(7)}{2(1)} = -\frac{7}{2}$. Since

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$f\left(-\frac{7}{2}\right)=-\frac{1}{4}$, the vertex is $\left(-\frac{1}{2},-\frac{1}{4}\right)$.



The graph is below the x-axis when $-4<x<-3$. Since the inequality is strict, the solution set is $\{x\,|\,-4<x<-3\}$ or, using interval notation, $(-4,-3)$.

15. $2x^2<5x+3$

$2x^2-5x-3<0$

We graph the function $f(x)=2x^2-5x-3$. The intercepts are


y-intercept: $f(0)=-3$

x-intercepts: $2x^2-5x-3=0$

$(2x+1)(x-3)=0$

$x=-\frac{1}{2}, x=3$

The vertex is at $x=\frac{-b}{2a}=\frac{-(-5)}{2(2)}=\frac{5}{4}$. Since $f\left(\frac{5}{4}\right)=-\frac{49}{8}$, the vertex is $\left(\frac{5}{4},-\frac{49}{8}\right)$.



The graph is below the x-axis when $-\frac{1}{2}<x<3$. Since the inequality is strict, the solution set is $\left\{x\left|-\frac{1}{2}<x<3\right.\right\}$ or, using interval notation, $\left(-\frac{1}{2},\ 3\right)$.

16. $6x^2<6+5x$

$6x^2-5x-6<0$

We graph the function $f(x)=6x^2-5x-6$. The intercepts are


y-intercept: $f(0)=-6$

x-intercepts: $6x^2-5x-6=0$

$(3x+2)(2x-3)=0$

$x=-\frac{2}{3}, x=\frac{3}{2}$

The vertex is at $x=\frac{-b}{2a}=\frac{-(-5)}{2(6)}=\frac{5}{12}$. Since $f\left(\frac{5}{12}\right)=-\frac{169}{24}$, the vertex is $\left(\frac{5}{12},-\frac{169}{24}\right)$.



The graph is below the x-axis when $-\frac{2}{3}<x<\frac{3}{2}$. Since the inequality is strict, the solution set is $\left\{x\left|-\frac{2}{3}<x<\frac{3}{2}\right.\right\}$ or, using interval notation, $\left(-\frac{2}{3},\ \frac{3}{2}\right)$.

17. $x^2-x+1\le 0$

We graph the function $f(x)=x^2-x+1$. The intercepts are

y-intercept: $f(0)=1$

x-intercepts: $x=\frac{-(-1)\pm\sqrt{(-1)^2-4(1)(1)}}{2(1)}$


$=\frac{1\pm\sqrt{-3}}{2}$ (not real)

Therefore, f has no x-intercepts.

The vertex is at $x=\frac{-b}{2a}=\frac{-(-1)}{2(1)}=\frac{1}{2}$. Since $f\left(\frac{1}{2}\right)=\frac{3}{4}$, the vertex is $\left(\frac{1}{2},\frac{3}{4}\right)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual



The graph is never below the x-axis. Thus, there is no real solution.

18. $x^2 + 2x + 4 > 0$

We graph the function $f(x) = x^2 + 2x + 4$.


y-intercept: $f(0) = 4$

x-intercepts: $x = \dfrac{-(2) \pm \sqrt{(2)^2 - 4(1)(4)}}{2(1)}$

$= \dfrac{-2 \pm \sqrt{-12}}{2}$ (not real)

Therefore, f has no x-intercepts.

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(2)}{2(1)} = -1$. Since $f(-1) = 3$, the vertex is $(-1, 3)$.



The graph is always above the x-axis. Thus, the solution is all real numbers or using interval notation, $(-\infty, \infty)$.

19. $4x^2 + 9 < 6x$

$4x^2 - 6x + 9 < 0$

We graph the function $f(x) = 4x^2 - 6x + 9$.


y-intercept: $f(0) = 9$

x-intercepts: $x = \dfrac{-(-6) \pm \sqrt{(-6)^2 - 4(4)(9)}}{2(4)}$

$= \dfrac{6 \pm \sqrt{-108}}{8}$ (not real)

Therefore, f has no x-intercepts.

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-6)}{2(4)} = \dfrac{6}{8} = \dfrac{3}{4}$. Since $f\left(\dfrac{3}{4}\right) = \dfrac{27}{4}$, the vertex is $\left(\dfrac{3}{4}, \dfrac{27}{4}\right)$.



The graph is never below the x-axis. Thus, there is no real solution.

20. $25x^2 + 16 < 40x$

$25x^2 - 40x + 16 < 0$

We graph the function $f(x) = 25x^2 - 40x + 16$.

y-intercept: $f(0) = 16$

x-intercepts: $25x^2 - 40x + 16 = 0$


$(5x - 4)^2 = 0$

$5x - 4 = 0$

$x = \dfrac{4}{5}$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-40)}{2(25)} = \dfrac{40}{50} = \dfrac{4}{5}$.

Since $f\left(\dfrac{4}{5}\right) = 0$, the vertex is $\left(\dfrac{4}{5}, 0\right)$.



The graph is never below the x-axis. Thus, there is no real solution.

21. $6(x^2 - 1) > 5x$

$6x^2 - 6 > 5x$

$6x^2 - 5x - 6 > 0$

We graph the function $f(x) = 6x^2 - 5x - 6$.

y-intercept: $f(0) = -6$

x-intercepts: $6x^2 - 5x - 6 = 0$


$(3x + 2)(2x - 3) = 0$

$x = -\dfrac{2}{3}, x = \dfrac{3}{2}$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-5)}{2(6)} = \dfrac{5}{12}$. Since $f\left(\dfrac{5}{12}\right) = -\dfrac{169}{24}$, the vertex is $\left(\dfrac{5}{12}, -\dfrac{169}{24}\right)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual



The graph is above the x-axis when $x < -\frac{2}{3}$ or $x > \frac{3}{2}$. Since the inequality is strict, solution set is $\left\{ x \middle| x < -\frac{2}{3} \text{ or } x > \frac{3}{2} \right\}$ or, using interval notation, $\left(-\infty, -\frac{2}{3}\right) \cup \left(\frac{3}{2}, \infty\right)$.

22. $2\left(2x^2 - 3x\right) > -9$

$4x^2 - 6x > -9$

$4x^2 - 6x + 9 > 0$

We graph the function $f(x) = 4x^2 - 6x + 9$.


y-intercept: $f(0) = 9$

x-intercepts: $x = \dfrac{-(-6) \pm \sqrt{(-6)^2 - 4(4)(9)}}{2(4)}$

$= \dfrac{6 \pm \sqrt{-108}}{8}$ (not real)

Therefore, f has no x-intercepts.

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-6)}{2(4)} = \dfrac{6}{8} = \dfrac{3}{4}$. Since $f\left(\frac{3}{4}\right) = \frac{27}{4}$, the vertex is $\left(\frac{3}{4}, \frac{27}{4}\right)$.



The graph is always above the x-axis. Thus, the solution set is all real numbers or, using interval notation, $(-\infty, \infty)$.

23. The domain of the expression $f(x) = \sqrt{x^2 - 16}$ includes all values for which $x^2 - 16 \geq 0$.

We graph the function $p(x) = x^2 - 16$. The intercepts of p are


y-intercept: $p(0) = -6$

x-intercepts: $x^2 - 16 = 0$

$(x+4)(x-4) = 0$

$x = -4, x = 4$

The vertex of p is at $x = \dfrac{-b}{2a} = \dfrac{-(0)}{2(1)} = 0$. Since $p(0) = -16$, the vertex is $(0, -16)$.



The graph of p is above the x-axis when $x < -4$ or $x > 4$. Since the inequality is not strict, the solution set of $x^2 - 16 \geq 0$ is $\{x \mid x \leq -4 \text{ or } x \geq 4\}$. Thus, the domain of f is also $\{x \mid x \leq -4 \text{ or } x \geq 4\}$ or, using interval notation, $(-\infty, -4] \cup [4, \infty)$.

24. The domain of the expression $f(x) = \sqrt{x - 3x^2}$ includes all values for which $x - 3x^2 \geq 0$.

We graph the function $p(x) = x - 3x^2$. The intercepts of p are

y-intercept: $p(0) = -6$


x-intercepts: $x - 3x^2 = 0$

$x(1 - 3x) = 0$

$x = 0, x = \dfrac{1}{3}$.

The vertex of p is at $x = \dfrac{-b}{2a} = \dfrac{-(1)}{2(-3)} = \dfrac{-1}{-6} = \dfrac{1}{6}$.

Since $p\left(\frac{1}{6}\right) = \frac{1}{12}$, the vertex is $\left(\frac{1}{6}, \frac{1}{12}\right)$.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

The graph of p is above the x-axis when $0 < x < \frac{1}{3}$. Since the inequality is not strict, the solution set of $x - 3x^2 \ge 0$ is $\left\{x \,\middle|\, 0 \le x \le \frac{1}{3}\right\}$. Thus, the domain of f is also $\left\{x \,\middle|\, 0 \le x \le \frac{1}{3}\right\}$ or, using interval notation, $\left[0, \frac{1}{3}\right]$.

25. $f(x) = x^2 - 1; \quad g(x) = 3x + 3$

a.
$$f(x) = 0$$
$$x^2 - 1 = 0$$
$$(x-1)(x+1) = 0$$
$$x = 1; x = -1$$
Solution set: $\{-1, 1\}$.

b.
$$g(x) = 0$$
$$3x + 3 = 0$$
$$3x = -3$$
$$x = -1$$
Solution set: $\{-1\}$.

c.
$$f(x) = g(x)$$
$$x^2 - 1 = 3x + 3$$
$$x^2 - 3x - 4 = 0$$
$$(x-4)(x+1) = 0$$
$$x = 4; x = -1$$
Solution set: $\{-1, 4\}$.

d. $f(x) > 0$

We graph the function $f(x) = x^2 - 1$.


y-intercept: $f(0) = -1$

x-intercepts: $x^2 - 1 = 0$

$(x+1)(x-1) = 0$

$x = -1, x = 1$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(1)} = 0$. Since $f(0) = -1$, the vertex is $(0, -1)$.



The graph is above the x-axis when $x < -1$ or $x > 1$. Since the inequality is strict, the solution set is $\{x \mid x < -1 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, -1) \cup (1, \infty)$.

e.
$$g(x) \le 0$$
$$3x + 3 \le 0$$
$$3x \le -3$$
$$x \le -1$$
The solution set is $\{x \mid x \le -1\}$ or, using interval notation, $(-\infty, -1]$.

f.
$$f(x) > g(x)$$
$$x^2 - 1 > 3x + 3$$
$$x^2 - 3x - 4 > 0$$
We graph the function $p(x) = x^2 - 3x - 4$.

The intercepts of p are


y-intercept: $p(0) = -4$

x-intercepts: $x^2 - 3x - 4 = 0$

$(x-4)(x+1) = 0$

$x = 4, x = -1$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-3)}{2(1)} = \frac{3}{2}$. Since $p\left(\frac{3}{2}\right) = -\frac{25}{4}$, the vertex is $\left(\frac{3}{2}, -\frac{25}{4}\right)$.



The graph of p is above the x-axis when $x < -1$ or $x > 4$. Since the inequality is strict, the solution set is $\{x \mid x < -1 \text{ or } x > 4\}$ or, using interval notation, $(-\infty, -1) \cup (4, \infty)$.

g.
$$f(x) \ge 1$$
$$x^2 - 1 \ge 1$$
$$x^2 - 2 \ge 0$$
We graph the function $p(x) = x^2 - 2$. The intercepts of p are

y-intercept: $p(0) = -2$

x-intercepts: $x^2 - 2 = 0$


$x^2 = 2$

$x = \pm\sqrt{2}$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(1)} = 0$. Since $p(0) = -2$, the vertex is $(0, -2)$.



The graph of p is above the x-axis when $x < -\sqrt{2}$ or $x > \sqrt{2}$. Since the inequality is not strict, the solution set is $\left\{ x \middle| x \le -\sqrt{2} \text{ or } x \ge \sqrt{2} \right\}$ or, using interval notation, $\left(-\infty, -\sqrt{2}\right] \cup \left[\sqrt{2}, \infty\right)$.

26. $f(x) = -x^2 + 3; \quad g(x) = -3x + 3$

a.
$$f(x) = 0$$
$$-x^2 + 3 = 0$$
$$x^2 = 3$$
$$x = \pm\sqrt{3}$$

Solution set: $\left\{-\sqrt{3}, \sqrt{3}\right\}$.

b.
$$g(x) = 0$$
$$-3x + 3 = 0$$
$$-3x = -3$$
$$x = 1$$

Solution set: $\{1\}$.

c.
$$f(x) = g(x)$$
$$-x^2 + 3 = -3x + 3$$
$$0 = x^2 - 3x$$
$$0 = x(x - 3)$$
$$x = 0; x = 3$$

Solution set: $\{0, 3\}$.

d. $f(x) > 0$


We graph the function $f(x) = -x^2 + 3$.

y-intercept: $f(0) = 3$

x-intercepts: $-x^2 + 3 = 0$

$$x^2 = 3$$
$$x = \pm\sqrt{3}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since $f(0) = 3$, the vertex is $(0, 3)$.



The graph is above the x-axis when $-\sqrt{3} < x < \sqrt{3}$. Since the inequality is strict, the solution set is $\left\{ x \middle| -\sqrt{3} < x < \sqrt{3} \right\}$ or, using interval notation, $\left(-\sqrt{3}, \sqrt{3}\right)$.

e.
$$g(x) \le 0$$
$$-3x + 3 \le 0$$
$$-3x \le -3$$
$$x \ge 1$$

The solution set is $\left\{ x \middle| x \ge 1 \right\}$ or, using interval notation, $\left[1, \infty\right)$.

f.
$$f(x) > g(x)$$
$$-x^2 + 3 > -3x + 3$$
$$-x^2 + 3x > 0$$

We graph the function $p(x) = -x^2 + 3x$.

The intercepts of p are


y-intercept: $p(0) = 0$

x-intercepts: $-x^2 + 3x = 0$

$$-x(x - 3) = 0$$
$$x = 0; \; x = 3$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(3)}{2(-1)} = \frac{-3}{-2} = \frac{3}{2}$.

Since $p\left(\frac{3}{2}\right) = \frac{9}{4}$, the vertex is $\left(\frac{3}{2}, \frac{9}{4}\right)$.



The graph of p is above the x-axis when $0 < x < 3$. Since the inequality is strict, the solution set is $\left\{ x \middle| 0 < x < 3 \right\}$ or, using interval notation, $(0, 3)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

g. $f(x) \ge 1$

$-x^2 + 3 \ge 1$

$-x^2 + 2 \ge 0$

We graph the function $p(x) = -x^2 + 2$. The intercepts of p are


y-intercept: $p(0) = 2$

x-intercepts: $-x^2 + 2 = 0$

$x^2 = 2$

$x = \pm\sqrt{2}$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(0)}{2(-1)} = 0$. Since $p(0) = 2$, the vertex is $(0, 2)$.



The graph of p is above the x-axis when $-\sqrt{2} < x < \sqrt{2}$. Since the inequality is not strict, the solution set is $\left\{ x \middle| -\sqrt{2} \le x \le \sqrt{2} \right\}$ or, using interval notation, $\left[-\sqrt{2}, \sqrt{2} \right]$.

27. $f(x) = -x^2 + 1; \quad g(x) = 4x + 1$

a. $f(x) = 0$

$-x^2 + 1 = 0$

$1 - x^2 = 0$

$(1 - x)(1 + x) = 0$

$x = 1; \; x = -1$

Solution set: $\{-1, 1\}$.

b. $g(x) = 0$

$4x + 1 = 0$

$4x = -1$

$x = -\dfrac{1}{4}$

Solution set: $\left\{ -\dfrac{1}{4} \right\}$.

c. $f(x) = g(x)$

$-x^2 + 1 = 4x + 1$

$0 = x^2 + 4x$

$0 = x(x + 4)$

$x = 0; \; x - 4$

Solution set: $\{-4, 0\}$.

d. $f(x) > 0$

We graph the function $f(x) = -x^2 + 1$.

y-intercept: $f(0) = 1$


x-intercepts: $-x^2 + 1 = 0$

$x^2 - 1 = 0$

$(x + 1)(x - 1) = 0$

$x = -1; x = 1$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(0)}{2(-1)} = 0$. Since $f(0) = 1$, the vertex is $(0, 1)$.



The graph is above the x-axis when $-1 < x < 1$. Since the inequality is strict, the solution set is $\left\{ x \middle| -1 < x < 1 \right\}$ or, using interval notation, $(-1, 1)$.

e. $g(x) \le 0$

$4x + 1 \le 0$

$4x \le -1$

$x \le -\dfrac{1}{4}$

The solution set is $\left\{ x \middle| x \le -\dfrac{1}{4} \right\}$ or, using interval notation, $\left(-\infty, -\dfrac{1}{4} \right]$.

f. $f(x) > g(x)$

$-x^2 + 1 > 4x + 1$

$-x^2 - 4x > 0$

We graph the function $p(x) = -x^2 - 4x$.

The intercepts of p are

y-intercept: $p(0) = 0$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


x-intercepts: $-x^2 - 4x = 0$

$-x(x+4) = 0$

$x = 0; x = -4$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-4)}{2(-1)} = \dfrac{4}{-2} = -2$.

Since $p(-2) = 4$, the vertex is $(-2, 4)$.




The graph of p is above the x-axis when $-4 < x < 0$. Since the inequality is strict, the solution set is $\{x \mid -4 < x < 0\}$ or, using interval notation, $(-4, 0)$.

g. $f(x) \geq 1$

$-x^2 + 1 \geq 1$

$-x^2 \geq 0$

We graph the function $p(x) = -x^2$. The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(0)}{2(-1)} = 0$. Since $p(0) = 0$, the vertex is $(0, 0)$. Since $a = -1 < 0$, the parabola opens downward.



The graph of p is never above the x-axis, but it does touch the x-axis at $x = 0$. Since the inequality is not strict, the solution set is $\{0\}$.

28. $f(x) = -x^2 + 4; \quad g(x) = -x - 2$

a. $f(x) = 0$

$-x^2 + 4 = 0$

$x^2 - 4 = 0$

$(x+2)(x-2) = 0$

$x = -2; x = 2$

Solution set: $\{-2, 2\}$.

b. $g(x) = 0$

$-x - 2 = 0$

$-2 = x$

Solution set: $\{-2\}$.

c. $f(x) = g(x)$

$-x^2 + 4 = -x - 2$

$0 = x^2 - x - 6$

$0 = (x-3)(x+2)$

$x = 3; x = -2$

Solution set: $\{-2, 3\}$.

d. $f(x) > 0$

$-x^2 + 4 > 0$

We graph the function $f(x) = -x^2 + 4$.

y-intercept: $f(0) = 4$


x-intercepts: $-x^2 + 4 = 0$

$x^2 - 4 = 0$

$(x+2)(x-2) = 0$

$x = -2; x = 2$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(0)}{2(-1)} = 0$. Since $f(0) = 4$, the vertex is $(0, 4)$.



The graph is above the x-axis when $-2 < x < 2$. Since the inequality is strict, the solution set is $\{x \mid -2 < x < 2\}$ or, using interval notation, $(-2, 2)$.

e. $g(x) \leq 0$

$-x - 2 \leq 0$

$-x \leq 2$

$x \geq -2$

The solution set is $\{x \mid x \geq -2\}$ or, using interval notation, $[-2, \infty)$.

f. $f(x) > g(x)$

$-x^2 + 4 > -x - 2$

$-x^2 + x + 6 > 0$

We graph the function $p(x) = -x^2 + x + 6$.

The intercepts of p are

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

y-intercept: $p(0) = 6$

x-intercepts: $-x^2 + x + 6 = 0$


$x^2 - x - 6 = 0$

$(x+2)(x-3) = 0$

$x = -2; x = 3$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(1)}{2(-1)} = \dfrac{-1}{-2} = \dfrac{1}{2}$.

Since $p\left(\dfrac{1}{2}\right) = \dfrac{25}{4}$, the vertex is $\left(\dfrac{1}{2}, \dfrac{25}{4}\right)$.



The graph of p is above the x-axis when $-2 < x < 3$. Since the inequality is strict, the solution set is $\{x \mid -2 < x < 3\}$ or, using interval notation, $(-2, 3)$.

g. $f(x) \geq 1$

$-x^2 + 4 > 1$

$-x^2 + 3 > 0$

We graph the function $p(x) = -x^2 + 3$. The intercepts of p are


y-intercept: $p(0) = 3$

x-intercepts: $-x^2 + 3 = 0$

$x^2 = 3$

$x = \pm\sqrt{3}$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(0)}{2(-1)} = 0$. Since $p(0) = 3$, the vertex is $(0, 3)$.



The graph of p is above the x-axis when $-\sqrt{3} < x < \sqrt{3}$. Since the inequality is not strict, the solution set is $\left\{x \mid -\sqrt{3} \leq x \leq \sqrt{3}\right\}$ or, using interval notation, $\left[-\sqrt{3}, \sqrt{3}\right]$.

29. $f(x) = x^2 - 4; \quad g(x) = -x^2 + 4$

a. $f(x) = 0$

$x^2 - 4 = 0$

$(x-2)(x+2) = 0$

$x = 2; x = -2$

Solution set: $\{-2, 2\}$.

b. $g(x) = 0$

$-x^2 + 4 = 0$

$x^2 - 4 = 0$

$(x+2)(x-2) = 0$

$x = -2; x = 2$

Solution set: $\{-2, 2\}$.

c. $f(x) = g(x)$

$x^2 - 4 = -x^2 + 4$

$2x^2 - 8 = 0$

$2(x-2)(x+2) = 0$

$x = 2; x = -2$

Solution set: $\{-2, 2\}$.

d. $f(x) > 0$

$x^2 - 4 > 0$

We graph the function $f(x) = x^2 - 4$.


y-intercept: $f(0) = -4$

x-intercepts: $x^2 - 4 = 0$

$(x+2)(x-2) = 0$

$x = -2; x = 2$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(0)}{2(-1)} = 0$. Since $f(0) = -4$, the vertex is $(0, -4)$.



The graph is above the x-axis when $x < -2$ or $x > 2$. Since the inequality is strict, the solution set is $\{x \mid x < -2 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -2) \cup (2, \infty)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

e. $g(x) \le 0$

$-x^2 + 4 \le 0$

We graph the function $g(x) = -x^2 + 4$.

y-intercept: $g(0) = 4$


x-intercepts: $-x^2 + 4 = 0$

$x^2 - 4 = 0$

$(x+2)(x-2) = 0$

$x = -2; x = 2$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since $g(0) = 4$, the vertex is $(0, 4)$.



The graph is below the x-axis when $x < -2$ or $x > 2$. Since the inequality is not strict, the solution set is $\{x \mid x \le -2 \text{ or } x \ge 2\}$ or, using interval notation, $(-\infty, -2] \cup [2, \infty)$.

f. $f(x) > g(x)$

$x^2 - 4 > -x^2 + 4$

$2x^2 - 8 > 0$

We graph the function $p(x) = 2x^2 - 8$.


y-intercept: $p(0) = -8$

x-intercepts: $2x^2 - 8 = 0$

$2(x+2)(x-2) = 0$

$x = -2; x = 2$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(2)} = 0$. Since $p(0) = -8$, the vertex is $(0, -8)$.



The graph is above the x-axis when $x < -2$ or $x > 2$. Since the inequality is strict, the solution set is $\{x \mid x < -2 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -2) \cup (2, \infty)$.

g. $f(x) \ge 1$

$x^2 - 4 \ge 1$

$x^2 - 5 \ge 0$

We graph the function $p(x) = x^2 - 5$.


y-intercept: $p(0) = -5$

x-intercepts: $x^2 - 5 = 0$

$x^2 = 5$

$x = \pm\sqrt{5}$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(1)} = 0$. Since $p(0) = -5$, the vertex is $(0, -5)$.



The graph of p is above the x-axis when $x < -\sqrt{5}$ or $x > \sqrt{5}$. Since the inequality is not strict, the solution set is $\{x \mid x \le -\sqrt{5} \text{ or } x \ge \sqrt{5}\}$ or, using interval notation, $(-\infty, -\sqrt{5}] \cup [\sqrt{5}, \infty)$.

30. $f(x) = x^2 - 2x + 1$; $g(x) = -x^2 + 1$

a. $f(x) = 0$

$x^2 - 2x + 1 = 0$

$(x-1)^2 = 0$

$x - 1 = 0$

$x = 1$

Solution set: $\{1\}$.

b. $g(x) = 0$

$-x^2 + 1 = 0$

$x^2 - 1 = 0$

$(x+1)(x-1) = 0$

$x = -1; x = 1$

Solution set: $\{-1, 1\}$.

c. $f(x) = g(x)$

$x^2 - 2x + 1 = -x^2 + 1$

$2x^2 - 2x = 0$

$2x(x-1) = 0$

$x = 0, x = 1$

Solution set: $\{0, 1\}$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

d. $f(x) > 0$

$x^2 - 2x + 1 > 0$

We graph the function $f(x) = x^2 - 2x + 1$.

y-intercept: $f(0) = 1$

x-intercepts: $x^2 - 2x + 1 = 0$


$(x-1)^2 = 0$

$x - 1 = 0$

$x = 1$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-2)}{2(1)} = \dfrac{2}{2} = 1$.

Since $f(1) = 0$, the vertex is $(1, 0)$.



The graph is above the x-axis when $x < 1$ or $x > 1$. Since the inequality is strict, the solution set is $\{x \mid x < 1 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, 1) \cup (1, \infty)$.

e. $g(x) \le 0$

$-x^2 + 1 \le 0$

We graph the function $g(x) = -x^2 + 1$.

y-intercept: $g(0) = 1$


x-intercepts: $-x^2 + 1 = 0$

$x^2 - 1 = 0$

$(x+1)(x-1) = 0$

$x = -1; x = 1$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(0)}{2(-1)} = 0$. Since $g(0) = 1$, the vertex is $(0, 1)$.



The graph is below the x-axis when $x < -1$ or $x > 1$. Since the inequality is not strict, the solution set is $\{x \mid x \le -1 \text{ or } x \ge 1\}$ or, using interval notation, $(-\infty, -1] \cup [1, \infty)$.

f. $f(x) > g(x)$

$x^2 - 2x + 1 > -x^2 + 1$

$2x^2 - 2x > 0$

We graph the function $p(x) = 2x^2 - 2x$.

y-intercept: $p(0) = 0$


x-intercepts: $2x^2 - 2x = 0$

$2x(x-1) = 0$

$x = 0; x = 1$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-2)}{2(2)} = \dfrac{2}{4} = \dfrac{1}{2}$.

Since $p\left(\dfrac{1}{2}\right) = \dfrac{1}{2}$, the vertex is $\left(\dfrac{1}{2}, \dfrac{1}{2}\right)$.



The graph is above the x-axis when $x < 0$ or $x > 1$. Since the inequality is strict, the solution set is $\{x \mid x < 0 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, 0) \cup (1, \infty)$.

g. $f(x) \ge 1$

$x^2 - 2x + 1 \ge 1$

$x^2 - 2x \ge 0$

We graph the function $p(x) = x^2 - 2x$.

y-intercept: $p(0) = 0$


x-intercepts: $x^2 - 2x = 0$

$x(x-2) = 0$

$x = 0; x = 2$

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-2)}{2(1)} = \dfrac{2}{2} = 1$.

Since $p(1) = -1$, the vertex is $(1, -1)$.



The graph of p is above the x-axis when $x < 0$ or $x > 2$. Since the inequality is not strict, the solution set is $\{x \mid x \le 0 \text{ or } x \ge 2\}$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

or, using interval notation,
$(-\infty, 0] \cup [2, \infty)$.


31. $f(x) = x^2 - x - 2; \quad g(x) = x^2 + x - 2$

a.
$$f(x) = 0$$
$$x^2 - x - 2 = 0$$
$$(x-2)(x+1) = 0$$
$$x = 2, x = -1$$
Solution set: $\{-1, 2\}$.

b.
$$g(x) = 0$$
$$x^2 + x - 2 = 0$$
$$(x+2)(x-1) = 0$$
$$x = -2; x = 1$$
Solution set: $\{-2, 1\}$.


c.
$$f(x) = g(x)$$
$$x^2 - x - 2 = x^2 + x - 2$$
$$-2x = 0$$
$$x = 0$$
Solution set: $\{0\}$.

d.
$$f(x) > 0$$
$$x^2 - x - 2 > 0$$
We graph the function $f(x) = x^2 - x - 2$.
y-intercept: $f(0) = -2$
x-intercepts: $x^2 - x - 2 = 0$
$$(x-2)(x+1) = 0$$
$$x = 2; x = -1$$
The vertex is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(1)} = \frac{1}{2}$. Since $f\left(\frac{1}{2}\right) = -\frac{9}{4}$, the vertex is $\left(\frac{1}{2}, -\frac{9}{4}\right)$.



The graph is above the x-axis when $x < -1$ or $x > 2$. Since the inequality is strict, the solution set is $\{x \mid x < -1 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -1) \cup (2, \infty)$.

e.
$$g(x) \le 0$$
$$x^2 + x - 2 \le 0$$
We graph the function $g(x) = x^2 + x - 2$.
y-intercept: $g(0) = -2$
x-intercepts: $x^2 + x - 2 = 0$
$$(x+2)(x-1) = 0$$
$$x = -2; x = 1$$
The vertex is at $x = \frac{-b}{2a} = \frac{-(1)}{2(1)} = -\frac{1}{2}$. Since $f\left(-\frac{1}{2}\right) = -\frac{7}{4}$, the vertex is $\left(-\frac{1}{2}, -\frac{7}{4}\right)$.




The graph is below the x-axis when $-2 < x < 1$. Since the inequality is not strict, the solution set is $\{x \mid -2 \le x \le 1\}$ or, using interval notation, $[-2, 1]$.

f.
$$f(x) > g(x)$$
$$x^2 - x - 2 > x^2 + x - 2$$
$$-2x > 0$$
$$x < 0$$
The solution set is $\{x \mid x < 0\}$ or, using interval notation, $(-\infty, 0)$.

g.
$$f(x) \ge 1$$
$$x^2 - x - 2 \ge 1$$
$$x^2 - x - 3 \ge 0$$
We graph the function $p(x) = x^2 - x - 3$.
y-intercept: $p(0) = -3$
x-intercepts: $x^2 - x - 3 = 0$
$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(-3)}}{2(1)}$$
$$= \frac{1 \pm \sqrt{1+12}}{2} = \frac{1 \pm \sqrt{13}}{2}$$
$$x \approx -1.30 \text{ or } x \approx 2.30$$
The vertex is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(1)} = \frac{1}{2}$. Since $p\left(\frac{1}{2}\right) = -\frac{13}{4}$, the vertex is $\left(\frac{1}{2}, -\frac{13}{4}\right)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual



The graph of p is above the x-axis when $x < \frac{1-\sqrt{13}}{2}$ or $x > \frac{1+\sqrt{13}}{2}$. Since the inequality is not strict, the solution set is $\left\{x \middle| x \le \frac{1-\sqrt{13}}{2} \text{ or } x \ge \frac{1+\sqrt{13}}{2}\right\}$ or, using interval notation, $\left(-\infty, \frac{1-\sqrt{13}}{2}\right] \cup \left[\frac{1+\sqrt{13}}{2}, \infty\right)$.

32. $f(x) = -x^2 - x + 1; \quad g(x) = -x^2 + x + 6$

a.
$$f(x) = 0$$
$$-x^2 - x + 1 = 0$$
$$x^2 + x - 1 = 0$$
$$x = \frac{-(1) \pm \sqrt{(1)^2 - 4(1)(-1)}}{2(1)}$$
$$= \frac{-1 \pm \sqrt{1+4}}{2} = \frac{-1 \pm \sqrt{5}}{2}$$

Solution set: $\left\{\frac{-1-\sqrt{5}}{2}, \frac{-1+\sqrt{5}}{2}\right\}$.

b.
$$g(x) = 0$$
$$-x^2 + x + 6 = 0$$
$$x^2 - x - 6 = 0$$
$$(x-3)(x+2) = 0$$
$$x = 3; x = -2$$

Solution set: $\{-2, 3\}$.

c.
$$f(x) = g(x)$$
$$-x^2 - x + 1 = -x^2 + x + 6$$
$$-2x - 5 = 0$$
$$-2x = 5$$
$$x = -\frac{5}{2}$$

Solution set: $\left\{-\frac{5}{2}\right\}$.

d.
$$f(x) > 0$$
$$-x^2 - x + 1 > 0$$


We graph the function $f(x) = -x^2 - x + 1$.

y-intercept: $f(0) = -1$

x-intercepts: $-x^2 - x + 2 = 0$
$$x^2 + x - 2 = 0$$
$$x = \frac{-(1) \pm \sqrt{(1)^2 - 4(1)(-1)}}{2(1)}$$
$$= \frac{-1 \pm \sqrt{1+4}}{2} = \frac{-1 \pm \sqrt{5}}{2}$$
$$x \approx -1.62 \text{ or } x \approx 0.62$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(-1)} = \frac{1}{-2} = -\frac{1}{2}$.

Since $f\left(-\frac{1}{2}\right) = \frac{5}{4}$, the vertex is $\left(-\frac{1}{2}, \frac{5}{4}\right)$.



The graph is above the x-axis when $\frac{-1-\sqrt{5}}{2} < x < \frac{-1+\sqrt{5}}{2}$. Since the inequality is strict, the solution set is $\left\{x \middle| \frac{-1-\sqrt{5}}{2} < x < \frac{-1+\sqrt{5}}{2}\right\}$ or, using interval notation, $\left(\frac{-1-\sqrt{5}}{2}, \frac{-1+\sqrt{5}}{2}\right)$.

e.
$$g(x) \le 0$$
$$-x^2 + x + 6 \le 0$$

We graph the function $g(x) = -x^2 + x + 6$.

y-intercept: $g(0) = 6$


x-intercepts: $-x^2 + x + 6 = 0$
$$x^2 - x - 6 = 0$$
$$(x-3)(x+2) = 0$$
$$x = 3; x = -2$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(1)}{2(-1)} = \frac{-1}{-2} = \frac{1}{2}$.

Since $f\left(\frac{1}{2}\right) = \frac{25}{4}$, the vertex is $\left(\frac{1}{2}, \frac{25}{4}\right)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual



The graph is below the x-axis when $x < -2$ or $x > 3$. Since the inequality is not strict, the solution set is $\{x \mid x \le -2 \text{ or } x \ge 3\}$ or, using interval notation, $(-\infty, 2] \cup [3, \infty)$.

f.
$$f(x) > g(x)$$
$$-x^2 - x + 1 > -x^2 + x + 6$$
$$-2x > 5$$
$$x < -\frac{5}{2}$$

The solution set is $\{x \mid x < -\frac{5}{2}\}$ or, using interval notation, $\left(-\infty, -\frac{5}{2}\right)$.

g.
$$f(x) \ge 1$$
$$-x^2 - x + 1 \ge 1$$
$$-x^2 - x \ge 0$$

We graph the function $p(x) = -x^2 - x$.


y-intercept: $p(0) = 0$

x-intercepts: $-x^2 - x = 0$

$$-x(x+1) = 0$$
$$x = 0; x = -1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(-1)} = \frac{1}{-2} = -\frac{1}{2}$.

Since $p\left(-\frac{1}{2}\right) = \frac{1}{4}$, the vertex is $\left(-\frac{1}{2}, \frac{1}{4}\right)$.



The graph of p is above the x-axis when $-1 < x < 0$. Since the inequality is not strict, the solution set is $\{x \mid -1 \le x \le 0\}$ or, using interval notation, $[-1, 0]$.

33. a. The ball strikes the ground when $s(t) = 80t - 16t^2 = 0$.

$$80t - 16t^2 = 0$$
$$16t(5 - t) = 0$$
$$t = 0, t = 5$$

The ball strikes the ground after 5 seconds.

b. Find the values of t for which

$$80t - 16t^2 > 96$$
$$-16t^2 + 80t - 96 > 0$$

We graph the function $f(t) = -16t^2 + 80t - 96$. The intercepts are


y-intercept: $f(0) = -96$

t-intercepts: $-16t^2 + 80t - 96 = 0$

$$-16(t^2 - 5t + 6) = 0$$
$$16(t-2)(t-3) = 0$$
$$t = 2, t = 3$$

The vertex is at $t = \frac{-b}{2a} = \frac{-(80)}{2(-16)} = 2.5$.

Since $f(2.5) = 4$, the vertex is $(2.5, 4)$.



The graph of f is above the t-axis when $2 < t < 3$. Since the inequality is strict, the solution set is $\{t \mid 2 < t < 3\}$ or, using interval notation, $(2, 3)$. The ball is more than 96 feet above the ground for times between 2 and 3 seconds.

34. a. The ball strikes the ground when $s(t) = 96t - 16t^2 = 0$.

$$96t - 16t^2 = 0$$
$$16t(6 - t) = 0$$
$$t = 0, t = 6$$

The ball strikes the ground after 6 seconds.

b. Find the values of t for which

$$96t - 16t^2 > 128$$
$$-16t^2 + 96t - 128 > 0$$

We graph $f(t) = -16t^2 + 96t - 128$. The intercepts are

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

y-intercept: $f(0) = -128$


t-intercepts: $-16t^2 + 96t - 128 = 0$

$16(t^2 - 6t + 8) = 0$

$-16(t-4)(t-2) = 0$

$t = 4,\ t = 2$

The vertex is at $t = \dfrac{-b}{2a} = \dfrac{-(96)}{2(-16)} = 3$. Since $f(3) = 16$, the vertex is $(3, 16)$.



The graph of f is above the t-axis when $2 < t < 4$. Since the inequality is strict, the solution set is $\{t \mid 2 < t < 4\}$ or, using interval notation, $(2, 4)$. The ball is more than 128 feet above the ground for times between 2 and 4 seconds.

35. a. $R(p) = -4p^2 + 4000p = 0$

$-4p(p - 1000) = 0$

$p = 0,\ p = 1000$

Thus, the revenue equals zero when the price is \$0 or \$1000.

b. Find the values of p for which

$-4p^2 + 4000p > 800{,}000$

$-4p^2 + 4000p - 800{,}000 > 0$

We graph $f(p) = -4p^2 + 4000p - 800{,}000$.

The intercepts are

y-intercept: $f(0) = -800{,}000$

p-intercepts:

$-4p^2 + 4000p - 800000 = 0$

$p^2 - 1000p + 200000 = 0$

$$p = \frac{-(-1000) \pm \sqrt{(-1000)^2 - 4(1)(200000)}}{2(1)}$$

$$= \frac{1000 \pm \sqrt{200000}}{2}$$


$$= \frac{1000 \pm 200\sqrt{5}}{2}$$

$$= 500 \pm 100\sqrt{5}$$

$p \approx 276.39;\ p \approx 723.61$.

The vertex is at $p = \dfrac{-b}{2a} = \dfrac{-(4000)}{2(-4)} = 500$.

Since $f(500) = 200{,}000$, the vertex is $(500, 200000)$.



The graph of f is above the p-axis when $276.39 < p < 723.61$. Since the inequality is strict, the solution set is $\{p \mid 276.39 < p < 723.61\}$ or, using interval notation, $(276.39, 723.61)$. The revenue is more than \$800,000 for prices between \$276.39 and \$723.61.

36. a. $R(p) = -\dfrac{1}{2}p^2 + 1900p = 0$

$-\dfrac{1}{2}p(p - 3800) = 0$

$p = 0,\ p = 3800$

Thus, the revenue equals zero when the price is \$0 or \$3800.

b. Find the values of p for which

$-\dfrac{1}{2}p^2 + 1900p > 1200000$

$-\dfrac{1}{2}p^2 + 1900p - 1200000 > 0$

We graph $f(p) = -\dfrac{1}{2}p^2 + 1900p - 1200000$.

The intercepts are

y-intercept: $f(0) = -1{,}200{,}000$

p-intercepts: $-\dfrac{1}{2}p^2 + 1900p - 1200000 = 0$

$p^2 - 3800p + 2400000 = 0$

$(p - 800)(p - 3000) = 0$


$p = 800;\ p = 3000$

The vertex is at $p = \dfrac{-b}{2a} = \dfrac{-(-1900)}{2(1/2)} = 1900$.

Since $f(1900) = 605{,}000$, the vertex is $(1900, 605000)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual



The graph of f is above the p-axis when $800 < p < 3000$. Since the inequality is strict, the solution set is $\{p \mid 800 < p < 3000\}$ or, using interval notation, $(800, 3000)$. The revenue is more than \$1,200,000 for prices between \$800 and \$3000.

37. $y = cx - \left(1 + c^2\right)\left(\frac{g}{2}\right)\left(\frac{x}{v}\right)^2$

a. Since the round must clear a hill 200 meters high, this mean $y > 200$.

Now $x = 2000$, $v = 897$, and $g = 9.81$.

$$c(2000) - \left(1 + c^2\right)\left(\frac{9.81}{2}\right)\left(\frac{2000}{897}\right)^2 > 200$$

$$2000c - 24.3845\left(1 + c^2\right) > 200$$

$$2000c - 24.3845 - 24.3845c^2 > 200$$

$$-24.3845c^2 + 2000c - 224.3845 > 0$$

We graph $f(c) = -24.3845c^2 + 2000c - 224.3845$.

The intercepts are

y-intercept: $f(0) = -224.3845$

c-intercepts:


$$-24.3845c^2 + 2000c - 224.3845 = 0$$

$$c = \frac{-2000 \pm \sqrt{(2000)^2 - 4(-24.3845)(-224.3845)}}{2(-24.3845)}$$

$$= \frac{-2000 \pm \sqrt{3{,}978{,}113.985}}{-48.769}$$

$c \approx 0.112$ or $c \approx 81.907$

The vertex is at $c = \frac{-b}{2a} = \frac{-(2000)}{2(-24.3845)} = 41.010$. Since $f(41.010) \approx 40{,}785.273$, the vertex is $(41.010, 40785.273)$.



The graph of f is above the c-axis when $0.112 < c < 81.907$. Since the inequality is strict, the solution set is $\{c \mid 0.112 < c < 81.907\}$ or, using interval notation, $(0.112, 81.907)$.

b. Since the round is to be on the ground $y = 0$.

Note, 75 km = 75,000 m. So, $x = 75{,}000$, $v = 897$, and $g = 9.81$.

$$c(75{,}000) - \left(1 + c^2\right)\left(\frac{9.81}{2}\right)\left(\frac{75{,}000}{897}\right)^2 = 0$$

$$75{,}000c - 34{,}290.724\left(1 + c^2\right) = 0$$

$$75{,}000c - 34{,}290.724 - 34{,}290.724c^2 = 0$$

$$-34{,}290.724c^2 + 75{,}000c - 34{,}290.724 = 0$$

We graph $f(c) = -34{,}290.724c^2 + 75{,}000c - 34{,}290.724$.

The intercepts are

y-intercept: $f(0) = -34{,}290.724$

c-intercepts:

$$-34{,}290.724c^2 + 75{,}000c - 34{,}290.724 = 0$$

$$c = \frac{-(75{,}000) \pm \sqrt{(75{,}000)^2 - 4(-34{,}290.724)(-34{,}290.724)}}{2(-34{,}290.724)}$$

$$= \frac{-75{,}000 \pm \sqrt{921{,}584{,}990.2}}{-68{,}581.448}$$

$c \approx 0.651$ or $c \approx 1.536$

It is possible to hit the target 75 kilometers away so long as $c \approx 0.651$ or $c \approx 1.536$.

38. $W = \frac{1}{2}kx^2$; $\tilde{W} = \frac{w}{2g}v^2$; $x \geq 0$

Note $v = 25$ mph $= \frac{110}{3}$ ft/sec. For $k = 9450$, $w = 4000$, $g = 32.2$, and $v = \frac{110}{3}$, we solve

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$$W > \tilde{W}$$

$$\frac{1}{2}(9450)x^2 > \frac{4000}{2(32.2)}\left(\frac{110}{3}\right)^2$$

$$4725x^2 > 83,505.866$$

$$x^2 > 17.6732$$

$$x^2 - 17.6732 > 0$$

We graph $f(x) = x^2 - 17.6732$. The intercepts are

y-intercept: $f(0) = -17.6732$


x-intercepts: $x^2 - 17.6732 = 0$

$$x^2 = 17.6732$$

$$x = \pm\sqrt{17.6732}$$

$$x \approx \pm 4.2$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(1)} = 0$. Since $f(0) = -17.6732$, the vertex is $(0, -17.6732)$.



The graph of f is above the x-axis when $x < -4.2$ or $x > 4.2$. Since we are restricted to $x \geq 0$, we disregard $x < -4.2$, so the solution is $x > 4.2$. Therefore, the spring must be able to compress at least 4.3 feet in order to stop the car safely.

39. $(x-4)^2 \leq 0$

We graph the function $f(x) = (x-4)^2$.


y-intercept: $f(0) = 16$

x-intercepts: $(x-4)^2 = 0$

$$x - 4 = 0$$

$$x = 4$$

The vertex is the vertex is $(4, 0)$.



The graph is never below the x-axis. Since the inequality is not strict, the only solution comes from the x-intercept. Therefore, the given inequality has exactly one real solution, namely $x = 4$.

40. $(x-2)^2 > 0$

We graph the function $f(x) = (x-2)^2$.


y-intercept: $f(0) = 4$

x-intercepts: $(x-2)^2 = 0$

$$x - 2 = 0$$

$$x = 2$$

The vertex is the vertex is $(2, 0)$.



The graph is above the x-axis when $x < 2$ or $x > 2$. Since the inequality is strict, the solution set is $\{x \mid x < 2 \text{ or } x > 2\}$. Therefore, the given inequality has exactly one real number that is not a solution, namely $x \neq 2$.


41. Solving $x^2 + x + 1 > 0$

We graph the function $f(x) = x^2 + x + 1$.

y-intercept: $f(0) = 1$

x-intercepts: $b^2 - 4ac = 1^2 - 4(1)(1) = -3$, so f has no x-intercepts.

The vertex is at $x = \frac{-b}{2a} = \frac{-(1)}{2(1)} = -\frac{1}{2}$. Since $f\left(-\frac{1}{2}\right) = \frac{3}{4}$, the vertex is $\left(-\frac{1}{2}, \frac{3}{4}\right)$.



The graph is always above the x-axis. Thus, the solution is the set of all real numbers or, using interval notation, $(-\infty, \infty)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

42. Solving $x^2 - x + 1 < 0$

We graph the function $f(x) = x^2 - x + 1$.

y-intercept: $f(0) = 1$

x-intercepts: $b^2 - 4ac = (-1)^2 - 4(1)(1) = -3$, so f has no x-intercepts.

The vertex is at $x = \dfrac{-b}{2a} = \dfrac{-(-1)}{2(1)} = \dfrac{1}{2}$. Since $f\left(-\dfrac{1}{2}\right) = \dfrac{3}{4}$, the vertex is $\left(-\dfrac{1}{2}, \dfrac{3}{4}\right)$.

The graph is never below the x-axis. Thus, the inequality has no solution. That is, the solution set is $\{\ \}$ or $\varnothing$.

43. The x-intercepts are included when the original inequality is not strict (when it contains an equal sign with the inequality).

44. Since the radical cannot be negative we determine what makes the radicand a nonnegative number.

$10 - 2x \ge 0$

$-2x \ge -10$

$x \le 5$

So the domain is: $\{x \mid x \le 5\}$.

45. a. $0 = \dfrac{2}{3}x - 6$

$6 = \dfrac{2}{3}x$


$x = 9$

$y = \dfrac{2}{3}(0) - 6$

$= -6$

The intercepts are: $(9,0), (0,-6)$

b.



46. $f(-x) = \dfrac{-(-x)}{(-x)^2 + 9}$

$= -\dfrac{-x}{x^2 + 9} = -f(x)$

Since $f(-x) = -f(x)$ then the function is odd.

47. $6x - 3y = 10$ $\qquad$ $2x + y = -8$

$-3y = -6x + 10$ $\qquad$ $y = -2x - 8$

$y = 2x - \dfrac{10}{3}$

Since the slopes are not equal and are not opposite reciprocals, the graphs are neither.

Section 2.6

1. $R = 3x$

2. Use LIN REGression to get $y = 1.7826x + 4.0652$

3. a. $R(x) = x\left(-\dfrac{1}{6}x + 100\right) = -\dfrac{1}{6}x^2 + 100x$

b. The quantity sold price cannot be negative, so $x \ge 0$. Similarly, the price should be positive, so $p > 0$.

$-\dfrac{1}{6}x + 100 > 0$

$-\dfrac{1}{6}x > -100$

$x < 600$

Thus, the implied domain for R is $\{x \mid 0 \le x < 600\}$ or $[0, 600)$.

c. $R(200) = -\dfrac{1}{6}(200)^2 + 100(200)$

$= \dfrac{-20000}{3} + 20000$

$= \dfrac{40000}{3} \approx \$13,333.33$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

d. $x = \dfrac{-b}{2a} = \dfrac{-100}{2\left(-\frac{1}{6}\right)} = \dfrac{-100}{\left(-\frac{1}{3}\right)} = \dfrac{300}{1} = 300$

The maximum revenue is

$$R(300) = -\frac{1}{6}(300)^2 + 100(300)$$
$$= -15000 + 30000$$
$$= \$15,000$$

e. $p = -\dfrac{1}{6}(300) + 100 = -50 + 100 = \$50$

4. a. $R(x) = x\left(-\dfrac{1}{3}x + 100\right) = -\dfrac{1}{3}x^2 + 100x$

b. The quantity sold price cannot be negative, so $x \geq 0$. Similarly, the price should be positive, so $p > 0$.

$$-\frac{1}{3}x + 100 > 0$$
$$-\frac{1}{3}x > -100$$
$$x < 300$$

Thus, the implied domain for R is $\{x \mid 0 \leq x < 300\}$ or $[0, 300)$.

c. $R(100) = -\dfrac{1}{3}(100)^2 + 100(100)$

$$= \frac{-10000}{3} + 10000$$
$$= \frac{20000}{3} \approx \$6,666.67$$

d. $x = \dfrac{-b}{2a} = \dfrac{-100}{2\left(-\frac{1}{3}\right)} = \dfrac{-100}{\left(-\frac{2}{3}\right)} = \dfrac{300}{2} = 150$

The maximum revenue is

$$R(150) = -\frac{1}{3}(150)^2 + 100(150)$$
$$= -7500 + 15000 = \$7,500$$

e. $p = -\dfrac{1}{3}(150) + 100 = -50 + 100 = \$50$

5. a. $R(x) = p(-5p + 100) = -5p^2 + 100p$

b. $R(15) = -5(17)^2 + 100(17)$

$$= -1445 + 1700 = \$255$$


c. $p = \dfrac{-b}{2a} = \dfrac{-100}{2(-5)} = \dfrac{-100}{(-10)} = 10$

The maximum revenue is

$$R(50) = -5(10)^2 + 100(50)$$
$$= -500 + 1000 = \$500$$

d. $x = -5(10) + 100 = 50$

e. Graph $R = -5p^2 + 100p$ and $R = 480$. Find where the graphs intersect by solving $480 = -5p^2 + 100x$.



$$5p^2 - 100p + 480 = 0$$
$$p^2 - 20p + 96 = 0$$
$$(p - 8)(p - 12) = 0$$
$$p = 8, p = 12$$

The company should charge between \$8 and \$12.

6. a. $R(x) = p(-20p + 500) = -20p^2 + 500p$

b. $R(24) = -20(24)^2 + 500(24)$

$$= -11528 + 12000 = \$480$$


c. $p = \dfrac{-b}{2a} = \dfrac{-500}{2(-20)} = \dfrac{-500}{(-40)} = \$12.50$.

The maximum revenue is

$$R(12.5) = -20(12.5)^2 + 500(12.5)$$
$$= -3125 + 6250 = \$3125$$

d. $x = -20(12.50) + 500 = 250$

e. Graph $R = -20p^2 + 500p$ and $R = 3000$. Find where the graphs intersect by solving $3000 = -20p^2 + 500p$.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$$20p^2 - 500p + 3000 = 0$$
$$p^2 - 25p + 150 = 0$$
$$(p-10)(p-15) = 0$$
$$p = 10, p = 15$$

The company should charge between \$10 and \$15.

7. a. Let w = width and l = length of the rectangular area.
Solving $P = 2w + 2l = 400$ for l:
$l = \dfrac{400 - 2w}{2} = 200 - w$.
Then $A(w) = (200 - w)w = 200w - w^2$
$= -w^2 + 200w$

b. $w = \dfrac{-b}{2a} = \dfrac{-200}{2(-1)} = \dfrac{-200}{-2} = 100$ yards

c. $A(100) = -100^2 + 200(100)$
$= -10000 + 20000$
$= 10{,}000 \text{ yd}^2$

8. a. Let x = width and y = width of the rectangle.
Solving $P = 2x + 2y = 3000$ for y:
$y = \dfrac{3000 - 2x}{2} = 1500 - x.$
Then $A(x) = (1500 - x)x$
$= 1500x - x^2$
$= -x^2 + 1500x.$

b. $x = \dfrac{-b}{2a} = \dfrac{-1500}{2(-1)} = \dfrac{-1500}{-2} = 750$ feet

c. $A(750) = -750^2 + 1500(750)$
$= -562500 + 1125000$
$= 562{,}500 \text{ ft}^2$

9. Let x = width and y = length of the rectangle.
Solving $P = 2x + y = 4000$ for y:
$y = 4000 - 2x$.
Then $A(x) = (4000 - 2x)x$
$= 4000x - 2x^2$
$= -2x^2 + 4000x$
$x = \dfrac{-b}{2a} = \dfrac{-4000}{2(-2)} = \dfrac{-4000}{-4} = 1000$ meters
maximizes area.
$A(1000) = -2(1000)^2 + 4000(1000)$.
$= -2000000 + 4000000$
$= 2{,}000{,}000$
The largest area that can be enclosed is 2,000,000 square meters.

10. Let x = width and y = length of the rectangle.
$2x + y = 2000$
$y = 2000 - 2x$
Then $A(x) = (2000 - 2x)x$
$= 2000x - 2x^2$
$= -2x^2 + 2000x$
$x = \dfrac{-b}{2a} = \dfrac{-2000}{2(-2)} = \dfrac{-2000}{-4} = 500$ meters
maximizes area.
$A(500) = -2(500)^2 + 2000(500)$
$= -500{,}000 + 1{,}000{,}000$
$= 500{,}000$
The largest area that can be enclosed is 500,000 square meters.

11. $h(x) = \dfrac{-32x^2}{(50)^2} + x + 200 = -\dfrac{8}{625}x^2 + x + 200$

a. $a = -\dfrac{8}{625}, b = 1, c = 200.$
The maximum height occurs when
$x = \dfrac{-b}{2a} = \dfrac{-1}{2(-8/625)} = \dfrac{625}{16} \approx 39$ feet from base of the cliff.


b. The maximum height is
$h\left(\dfrac{625}{16}\right) = \dfrac{-8}{625}\left(\dfrac{625}{16}\right)^2 + \dfrac{625}{16} + 200$
$= \dfrac{7025}{32} \approx 219.5$ feet.

c. Solving when $h(x) = 0$:
$-\dfrac{8}{625}x^2 + x + 200 = 0$
$x = \dfrac{-1 \pm \sqrt{1^2 - 4(-8/625)(200)}}{2(-8/625)}$
$x \approx \dfrac{-1 \pm \sqrt{11.24}}{-0.0256}$
$x \approx -91.90$ or $x \approx 170$
Since the distance cannot be negative, the projectile strikes the water approximately 170 feet from the base of the cliff.


Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


d.



e. Using the MAXIMUM function



Using the ZERO function



f. $-\frac{8}{625}x^2 + x + 200 = 100$

$-\frac{8}{625}x^2 + x + 100 = 0$

$x = \frac{\sqrt{1^2 - 4(-8/625)(100)}}{2(-8/625)} = \frac{-1 \pm \sqrt{6.12}}{-0.0256}$

$x \approx -57.57$ or $x \approx 135.70$

Since the distance cannot be negative, the projectile is 100 feet above the water when it is approximately 135.7 feet from the base of the cliff.

12. a. $h(x) = \frac{-32x^2}{(100)^2} + x = -\frac{2}{625}x^2 + x$

$a = -\frac{2}{625}, b = 1, c = 0.$

The maximum height occurs when

$x = \frac{-b}{2a} = \frac{-1}{2(-2/625)} = \frac{625}{4} = 156.25$ feet

b. The maximum height is

$h\left(\frac{625}{4}\right) = \frac{-2}{625}\left(\frac{625}{4}\right)^2 + \frac{625}{4}$

$= \frac{625}{8} = 78.125$ feet

c. Solving when $h(x) = 0$:

$-\frac{2}{625}x^2 + x = 0$

$x\left(-\frac{2}{625}x + 1\right) = 0$


$x = 0$ or $-\frac{2}{625}x + 1 = 0$

$x = 0$ or $1 = \frac{2}{625}x$


$x = 0$ or $x = \frac{625}{2} = 312.5$

Since the distance cannot be zero, the projectile lands 312.5 feet from where it was fired.


d.



e. Using the MAXIMUM function



Using the ZERO function



f. Solving when $h(x) = 50$:

$-\frac{2}{625}x^2 + x = 50$

$-\frac{2}{625}x^2 + x - 50 = 0$

$x = \frac{-1 \pm \sqrt{1^2 - 4(-2/625)(-50)}}{2(-2/625)}$

$= \frac{-1 \pm \sqrt{0.36}}{-0.0064} \approx \frac{-1 \pm 0.6}{-0.0064}$

$x = 62.5$ or $x = 250$

The projectile is 50 feet above the ground 62.5 feet and 250 feet from where it was fired.

Copyright © 2019 Pearson Education, Inc.


Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

13. Locate the origin at the point where the cable touches the road. Then the equation of the parabola is of the form: $y = ax^2$, where $a > 0$. Since the point (200, 75) is on the parabola, we can find the constant a:

Since $75 = a(200)^2$, then $a = \dfrac{75}{200^2} = 0.001875$.


When $x = 100$, we have:

$y = 0.001875(100)^2 = 18.75$ meters.



14. Locate the origin at the point directly under the highest point of the arch. Then the equation of the parabola is of the form: $y = -ax^2 + k$, where $a > 0$. Since the maximum height is 25 feet, when $x = 0$, $y = k = 25$. Since the point (60, 0) is on the parabola, we can find the constant a: Since $0 = -a(60)^2 + 25$ then $a = \dfrac{25}{60^2}$. The equation of the parabola is:

$h(x) = -\dfrac{25}{60^2}x^2 + 25$.



At $x = 10$:

$h(10) = -\dfrac{25}{60^2}(10)^2 + 25 = -\dfrac{25}{36} + 25 \approx 24.3$ ft.

At $x = 20$:

$h(20) = -\dfrac{25}{60^2}(20)^2 + 25 = -\dfrac{25}{9} + 25 \approx 22.2$ ft.

At $x = 40$:


$h(40) = -\dfrac{25}{60^2}(40)^2 + 25 = -\dfrac{100}{9} + 25 \approx 13.9$ ft.

15. a. Let x = the depth of the gutter and y the width of the gutter. Then $A = xy$ is the cross-sectional area of the gutter. Since the aluminum sheets for the gutter are 12 inches wide, we have $2x + y = 12$. Solving for y: $y = 12 - 2x$. The area is to be maximized, so: $A = xy = x(12 - 2x) = -2x^2 + 12x$. This equation is a parabola opening down; thus, it has a maximum

when $x = \dfrac{-b}{2a} = \dfrac{-12}{2(-2)} = \dfrac{-12}{-4} = 3$.

Thus, a depth of 3 inches produces a maximum cross-sectional area.

b. Graph $A = -2x^2 + 12x$ and $A = 16$. Find where the graphs intersect by solving $16 = -2x^2 + 12x$.



$$2x^2 - 12x + 16 = 0$$
$$x^2 - 6x + 8 = 0$$
$$(x-4)(x-2) = 0$$
$$x = 4, x = 2$$

The graph of $A = -2x^2 + 12x$ is above the graph of $A = 16$ where the depth is between 2 and 4 inches.

16. Let $x =$ width of the window and $y =$ height of the rectangular part of the window. The perimeter of the window is: $x + 2y + \dfrac{\pi x}{2} = 20$.

Solving for y: $y = \dfrac{40 - 2x - \pi x}{4}$.

The area of the window is:

$$A(x) = x\left(\frac{40 - 2x - \pi x}{4}\right) + \frac{1}{2}\pi\left(\frac{x}{2}\right)^2$$
$$= 10x - \frac{x^2}{2} - \frac{\pi x^2}{4} + \frac{\pi x^2}{8}$$
$$= \left(-\frac{1}{2} - \frac{\pi}{8}\right)x^2 + 10x.$$

This equation is a parabola opening down; thus, it has a maximum when

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$$x = \frac{-b}{2a} = \frac{-10}{2\left(-\frac{1}{2} - \frac{\pi}{8}\right)} = \frac{10}{\left(1 + \frac{\pi}{4}\right)} \approx 5.6 \text{ feet}$$

$$y = \frac{40 - 2(5.60) - \pi(5.60)}{4} \approx 2.8 \text{ feet}$$

The width of the window is about 5.6 feet and the height of the rectangular part is approximately 2.8 feet. The radius of the semicircle is roughly 2.8 feet, so the total height is about 5.6 feet.

17. Let x = the width of the rectangle or the diameter of the semicircle and let y = the length of the rectangle. The perimeter of each semicircle is $\frac{\pi x}{2}$.

The perimeter of the track is given by: $\frac{\pi x}{2} + \frac{\pi x}{2} + y + y = 1500$.

Solving for x:

$$\pi x + 2y = 1500$$
$$\pi x = 1500 - 2y$$
$$x = \frac{1500 - 2y}{\pi}$$

The area of the rectangle is:

$$A = xy = \left(\frac{1500 - 2y}{\pi}\right) y = \frac{-2}{\pi} y^2 + \frac{1500}{\pi} y.$$

This equation is a parabola opening down; thus, it has a maximum when

$$y = \frac{-b}{2a} = \frac{\frac{-1500}{\pi}}{2\left(\frac{-2}{\pi}\right)} = \frac{-1500}{-4} = 375.$$

Thus, $x = \frac{1500 - 2(375)}{\pi} = \frac{750}{\pi} \approx 238.73$

The dimensions for the rectangle with maximum area are $\frac{750}{\pi} \approx 238.73$ meters by 375 meters.

18. Let x = width of the window and y = height of the rectangular part of the window. The perimeter of the window is:

$$3x + 2y = 16$$
$$y = \frac{16 - 3x}{2}$$

The area of the window is

$$A(x) = x\left(\frac{16 - 3x}{2}\right) + \frac{\sqrt{3}}{4}x^2 = 8x - \frac{3}{2}x^2 + \frac{\sqrt{3}}{4}x^2$$
$$= \left(-\frac{3}{2} + \frac{\sqrt{3}}{4}\right)x^2 + 8x$$

This equation is a parabola opening down; thus, it has a maximum when

$$x = \frac{-b}{2a} = \frac{-8}{2\left(-\frac{3}{2} + \frac{\sqrt{3}}{4}\right)} = \frac{-8}{-3 + \frac{\sqrt{3}}{2}} = \frac{-16}{-6 + \sqrt{3}} \approx 3.75 \text{ ft.}$$

The window is approximately 3.75 feet wide.

$$y = \frac{16 - 3\left(\frac{-16}{-6 + \sqrt{3}}\right)}{2} = \frac{16 + \frac{48}{-6 + \sqrt{3}}}{2} = 8 + \frac{24}{-6 + \sqrt{3}}$$

The height of the equilateral triangle is

$$\frac{\sqrt{3}}{2}\left(\frac{-16}{-6 + \sqrt{3}}\right) = \frac{-8\sqrt{3}}{-6 + \sqrt{3}} \text{ feet, so the total height is}$$

$$8 + \frac{24}{-6 + \sqrt{3}} + \frac{-8\sqrt{3}}{-6 + \sqrt{3}} \approx 5.62 \text{ feet.}$$

19. We are given: $V(x) = kx(a - x) = -kx^2 + akx$.

The reaction rate is a maximum when:

$$x = \frac{-b}{2a} = \frac{-ak}{2(-k)} = \frac{ak}{2k} = \frac{a}{2}.$$

20. We have:

$$a(-h)^2 + b(-h) + c = ah^2 - bh + c = y_0$$
$$a(0)^2 + b(0) + c = c = y_1$$
$$a(h)^2 + b(h) + c = ah^2 + bh + c = y_2$$

Equating the two equations for the area, we have:

$$y_0 + 4y_1 + y_2 = ah^2 - bh + c + 4c + ah^2 + bh + c$$
$$= 2ah^2 + 6c.$$

Therefore,

$$\text{Area} = \frac{h}{3}\left(2ah^2 + 6c\right) = \frac{h}{3}\left(y_0 + 4y_1 + y_2\right) \text{ sq. units.}$$

21. $f(x) = -5x^2 + 8,\ h = 1$

$$\text{Area} = \frac{h}{3}\left(2ah^2 + 6c\right) = \frac{1}{3}\left(2(-5)(1)^2 + 6(8)\right)$$
$$= \frac{1}{3}(-10 + 48) = \frac{38}{3} \text{ sq. units}$$

22. $f(x) = 2x^2 + 8,\quad h = 2$

$$\text{Area} = \frac{h}{3}(2ah^2 + 6c) = \frac{2}{3}\left(2(2)(2)^2 + 6(8)\right)$$
$$= \frac{2}{3}(16 + 48) = \frac{2}{3}(64) = \frac{128}{3} \text{ sq. units}$$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


23. $f(x) = x^2 + 3x + 5,\ \ h = 4$

$$\text{Area} = \frac{h}{3}\left(2ah^2 + 6c\right) = \frac{4}{3}\left(2(1)(4)^2 + 6(5)\right)$$
$$= \frac{4}{3}(32 + 30) = \frac{248}{3} \text{ sq. units}$$

24. $f(x) = -x^2 + x + 4,\quad h = 1$


$$\text{Area} = \frac{h}{3}(2ah^2 + 6c) = \frac{1}{3}\left(2(-1)(1)^2 + 6(4)\right)$$
$$= \frac{1}{3}(-2 + 24) = \frac{1}{3}(22) = \frac{22}{3} \text{ sq. units}$$

25. a.



From the graph, the data appear to follow a quadratic relation with $a < 0$.

b. Using the QUADratic REGression program




$I(x) = -58.56x^2 + 5301.617x - 46,236.523$

c. $x = \dfrac{-b}{2a} = \dfrac{-5301.617}{2(-58.56)} \approx 45.3$


An individual will earn the most income at about 45.3 years of age.

d. The maximum income will be: $I(48.0) =$
$-58.56(45.3)^2 + 5301.617(45.3) - 46,236.523$
$\approx \$73,756$

e.




26. a.



From the graph, the data appear to follow a quadratic relation with $a < 0$.

b. Using the QUADratic REGression program




$h(x) = -0.0037x^2 + 1.0318x + 5.6667$

c. $x = \dfrac{-b}{2a} = \dfrac{-1.0318}{2(-0.0037)} \approx 139.4$


The ball will travel about 139.4 feet before it reaches its maximum height.

d. The maximum height will be: $h(139.4) =$
$-0.0037(139.4)^2 + 1.0318(139.4) + 5.6667$
≈ 77.6 feet

e.




27. a.



From the graph, the data appear to be linearly related with $m > 0$.

b. Using the LINear REGression program



$R(x) = 1.321x + 920.161$


Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

c. $R(875) = 1.321(875) + 920.161 \approx 2076$


The rent for an 875 square-foot apartment in San Diego will be about \$2076 per month.

28. a.



From the graph, the data appear to follow a quadratic relation with $a < 0$.

b. Using the QUADratic REGression program




$M(s) = -0.017s^2 + 1.935s - 25.341$

c. $M(63) = -0.017(63)^2 + 1.935(63) - 25.341$

≈ 29.1


A Camry traveling 63 miles per hour will get about 29.1 miles per gallon.

29. a.



From the graph, the data appear to follow a quadratic relation with $a < 0$.

b. Using the QUADratic REGression program




$B(a) = -0.563a^2 + 32.520a - 368.118$

c. $B(35) = -0.563(35)^2 + 32.520(35) - 368.118$

≈ 80.4


The birthrate of 35-year-old women is about 80.4 per 1000.

30. a.



From the graph, the data appear to be linearly related with $m > 0$.

b. Using the LINear REGression program



$C(x) = 0.233x - 2.037$

c. $C(80) = 0.233(80) - 2.037 \approx 16.6$

When the temperature is $80°\text{F}$, there will be about 16.6 chirps per second.

31. Answers will vary. One possibility follows: If the price is \$140, no one will buy the calculators, thus making the revenue \$0.

32. $m = \dfrac{2-(-2)}{-5-1} = \dfrac{4}{-6} = -\dfrac{2}{3}$

$$y - y_1 = m(x - x_1)$$
$$y - (-2) = -\frac{2}{3}(x - 1)$$
$$y + 2 = -\frac{2}{3}x + \frac{2}{3}$$
$$y = -\frac{2}{3}x - \frac{4}{3}$$

or

$$3y = -2x - 4$$
$$2x + 3y = -4$$

33. $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

$= \sqrt{((-1) - 4)^2 + (5 - (-7))^2}$

$= \sqrt{(-5)^2 + (12)^2}$

$= \sqrt{25 + 144} = \sqrt{169} = 13$

34. $(x - h)^2 + (y - k)^2 = r^2$

$(x - (-6))^2 + (y - 0)^2 = (\sqrt{7})^2$

$(x + 6)^2 + y^2 = 7$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

35. $3(0)^2 - 4y = 48$

$-4y = 48$

$y = -12$

The y intercept is $(0, -12)$

$3x^2 - 4(0) = 48$

$3x^2 = 48$

$x^2 = 16$

$x = \pm 4$

The x intercepts are: $(4, 0), (-4, 0)$

Section 2.7

1. Integers: $\{-3, 0\}$

Rationals: $\left\{-3, 0, \frac{6}{5}\right\}$

2. True; the set of real numbers consists of all rational and irrational numbers.

3. $10 - 5i$

4. $2 - 5i$

5. True

6. $9i$

7. $2 + 3i$

8. True


9. $f(x) = 0$

$x^2 + 4 = 0$

$x^2 = -4$

$x = \pm\sqrt{-4} = \pm 2i$

The zero are $-2i$ and $2i$.




10. $f(x) = 0$

$x^2 - 9 = 0$

$x^2 = 9$

$x = \pm\sqrt{9} = \pm 3$

The zeros are -3 and 3.




11. $f(x) = 0$

$x^2 - 16 = 0$

$x^2 = 16$

$x = \pm\sqrt{16} = \pm 4$

The zeros are -4 and 4.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


12. $f(x)=0$

$x^2+25=0$

$x^2=-25$

$x=\pm\sqrt{-25}=\pm 5i$

The zeros are $-5i$ and $5i$.



13. $f(x)=0$


$x^2-6x+13=0$

$a=1, b=-6, c=13,$

$b^2-4ac=(-6)^2-4(1)(13)=36-52=-16$

$x=\dfrac{-(-6)\pm\sqrt{-16}}{2(1)}=\dfrac{6\pm 4i}{2}=3\pm 2i$

The zeros are $3-2i$ and $3+2i$.



14. $f(x)=0$


$x^2+4x+8=0$

$a=1, b=4, c=8$

$b^2-4ac=4^2-4(1)(8)=16-32=-16$

$x=\dfrac{-4\pm\sqrt{-16}}{2(1)}=\dfrac{-4\pm 4i}{2}=-2\pm 2i$

The zeros are $-2-2i$ and $-2+2i$.



15. $f(x)=0$


$x^2-6x+10=0$

$a=1, b=-6, c=10$

$b^2-4ac=(-6)^2-4(1)(10)=36-40=-4$

$x=\dfrac{-(-6)\pm\sqrt{-4}}{2(1)}=\dfrac{6\pm 2i}{2}=3\pm i$

The zeros are $3-i$ and $3+i$.



16. $f(x)=0$


$x^2-2x+5=0$

$a=1, b=-2, c=5$

$b^2-4ac=(-2)^2-4(1)(5)=4-20=-16$

$x=\dfrac{-(-2)\pm\sqrt{-16}}{2(1)}=\dfrac{2\pm 4i}{2}=1\pm 2i$

The zeros are $1-2i$ and $1+2i$.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

17. $f(x)=0$


$x^2-4x+1=0$

$a=1, b=-4, c=1$

$b^2-4ac=(-4)^2-4(1)(1)=16-4=12$

$x=\dfrac{-(-4)\pm\sqrt{12}}{2(1)}=\dfrac{4\pm2\sqrt{3}}{2}=2\pm\sqrt{3}$

The zeros are $2-\sqrt{3}$ and $2+\sqrt{3}$, or approximately 0.27 and 3.73.



18. $f(x)=0$


$x^2+6x+1=0$

$a=1, b=6, c=1$

$b^2-4ac=6^2-4(1)(1)=36-4=32$

$x=\dfrac{-6\pm\sqrt{32}}{2(1)}=\dfrac{-6\pm4\sqrt{2}}{2}=-3\pm2\sqrt{2}$

The zeros are $-3-2\sqrt{2}$ and $-3+2\sqrt{2}$, or approximately -5.83 and -0.17.



19. $f(x)=0$


$2x^2+2x+1=0$

$a=2, b=2, c=1$

$b^2-4ac=(2)^2-4(2)(1)=4-8=-4$

$x=\dfrac{-2\pm\sqrt{-4}}{2(2)}=\dfrac{-2\pm2i}{4}=-\dfrac{1}{2}\pm\dfrac{1}{2}i$

The zeros are $-\dfrac{1}{2}-\dfrac{1}{2}i$ and $-\dfrac{1}{2}+\dfrac{1}{2}i$.



20. $f(x)=0$


$3x^2+6x+4=0$

$a=3, b=6, c=4$

$b^2-4ac=(6)^2-4(3)(4)=36-48=-12$

$x=\dfrac{-6\pm\sqrt{-12}}{2(3)}=\dfrac{-6\pm2\sqrt{3}i}{6}=-1\pm\dfrac{\sqrt{3}}{3}i$

The zeros are $-1-\dfrac{\sqrt{3}}{3}i$ and $-1+\dfrac{\sqrt{3}}{3}i$.



21. $f(x)=0$


$x^2+x+1=0$

$a=1, b=1, c=1,$

$b^2-4ac=1^2-4(1)(1)=1-4=-3$

$x=\dfrac{-1\pm\sqrt{-3}}{2(1)}=\dfrac{-1\pm\sqrt{3}\,i}{2}=-\dfrac{1}{2}\pm\dfrac{\sqrt{3}}{2}i$

The zeros are $-\dfrac{1}{2}-\dfrac{\sqrt{3}}{2}i$ and $-\dfrac{1}{2}+\dfrac{\sqrt{3}}{2}i$.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

22. $f(x) = 0$


$x^2 - x + 1 = 0$

$a = 1, b = -1, c = 1$

$b^2 - 4ac = (-1)^2 - 4(1)(1) = 1 - 4 = -3$

$x = \dfrac{-(-1) \pm \sqrt{-3}}{2(1)} = \dfrac{1 \pm \sqrt{3}\,i}{2} = \dfrac{1}{2} \pm \dfrac{\sqrt{3}}{2} i$

The zeros are $\dfrac{1}{2} - \dfrac{\sqrt{3}}{2} i$ and $\dfrac{1}{2} + \dfrac{\sqrt{3}}{2} i$.



23. $f(x) = 0$


$-2x^2 + 8x + 1 = 0$

$a = -2, b = 8, c = 1$

$b^2 - 4ac = 8^2 - 4(-2)(1) = 64 + 8 = 72$

$x = \dfrac{-8 \pm \sqrt{72}}{2(-2)} = \dfrac{-8 \pm 6\sqrt{2}}{-4} = \dfrac{4 \pm 3\sqrt{2}}{2} = 2 \pm \dfrac{3\sqrt{2}}{2}$

The zeros are $\dfrac{4 - 3\sqrt{2}}{2}$ and $\dfrac{4 + 3\sqrt{2}}{2}$, or approximately -0.12 and 4.12.



24. $f(x) = 0$


$-3x^2 + 6x + 1 = 0$

$a = -3, b = 6, c = 1$

$b^2 - 4ac = 6^2 - 4(-3)(1) = 36 + 12 = 48$

$x = \dfrac{-6 \pm \sqrt{48}}{2(-3)} = \dfrac{-6 \pm 4\sqrt{3}}{-6} = \dfrac{3 \pm 2\sqrt{3}}{3} = 1 \pm \dfrac{2\sqrt{3}}{3}$

The zeros are $\dfrac{3 - 2\sqrt{3}}{3}$ and $\dfrac{3 + 2\sqrt{3}}{3}$, or approximately -0.15 and 2.15.



25. $3x^2 - 3x + 4 = 0$

$a = 3, b = -3, c = 4$

$b^2 - 4ac = (-3)^2 - 4(3)(4) = 9 - 48 = -39$

The equation has two complex solutions that are conjugates of each other.

26. $2x^2 - 4x + 1 = 0$

$a = 2, b = -4, c = 1$

$b^2 - 4ac = (-4)^2 - 4(2)(1) = 16 - 8 = 8$

The equation has two unequal real number solutions.

27. $2x^2 + 3x - 4 = 0$

$a = 2, b = 3, c = -4$

$b^2 - 4ac = 3^2 - 4(2)(-4) = 9 + 32 = 41$

The equation has two unequal real solutions.

28. $x^2 + 2x + 6 = 0$

$a = 1, b = 2, c = 6$

$b^2 - 4ac = (2)^2 - 4(1)(6) = 4 - 24 = -20$

The equation has two complex solutions that are conjugates of each other.

29. $9x^2 - 12x + 4 = 0$

$a = 9, b = -12, c = 4$

$b^2 - 4ac = (-12)^2 - 4(9)(4) = 144 - 144 = 0$

The equation has a repeated real solution.

30. $4x^2 + 12x + 9 = 0$

$a = 4, b = 12, c = 9$

$b^2 - 4ac = 12^2 - 4(4)(9) = 144 - 144 = 0$

The equation has a repeated real solution.

31. $t^4 - 16 = 0$

$(t^2 - 4)(t^2 + 4) = 0$

$t^2 = 4 \quad t^2 = -4$

$t = \pm 2 \quad t = \pm 2i$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

32. $y^4 - 81 = 0$

$(y^2 - 9)(y^2 + 9) = 0$

$y^2 = 9 \quad y^2 = -9$

$y = \pm 3 \quad y = \pm 3i$

33. $F(x) = x^6 - 9x^3 + 8 = 0$

$(x^3 - 8)(x^3 - 1) = 0$

$(x-2)(x^2 + 2x + 4)(x-1)(x^2 + x + 1) = 0$

$x^2 + 2x + 4 = 0 \to a = 1, b = 2, c = 4$

$$x = \frac{-2 \pm \sqrt{2^2 - 4(4)}}{2(1)} = \frac{-2 \pm \sqrt{-12}}{2} = \frac{-2 \pm 2i\sqrt{3}}{2}$$

$$= -1 \pm \sqrt{3}i$$

$x^2 + x + 1 = 0 \to a = 1, b = 1, c = 1$

$$x = \frac{-1 \pm \sqrt{1^2 - 4(1)}}{2(1)} = \frac{-1 \pm \sqrt{-3}}{2} = \frac{-1 \pm i\sqrt{3}}{2}$$

$$= -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$$

The solution set is $\left\{-1 \pm i\sqrt{3}, -\frac{1}{2} \pm i\frac{\sqrt{3}}{2}, 2, 1\right\}$

34. $P(z) = z^6 + 28z^3 + 27 = 0$

$(z^3 + 27)(z^3 + 1) = 0$

$(z+3)(z^2 - 3z + 9)(z+1)(z^2 - z + 1) = 0$

$z^2 - 3z + 9 = 0$

$a = 1, b = -3, c = 9$

$$z = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(9)}}{2(1)} = \frac{3 \pm \sqrt{-27}}{2}$$

$$= \frac{3 \pm 3i\sqrt{3}}{2} = \frac{3}{2} \pm \frac{3\sqrt{3}}{2}i$$

$z^2 - z + 1 = 0 \to a = 1, b = -1, c = 1$

$$z = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)}}{2(1)} = \frac{1 \pm \sqrt{-3}}{2} = \frac{1 \pm i\sqrt{3}}{2}$$

$$= \frac{1}{2} \pm \frac{\sqrt{3}}{2}i$$

The solution set is $\left\{\frac{3}{2} \pm \frac{3\sqrt{3}}{2}i, \frac{1}{2} \pm \frac{\sqrt{3}}{2}i, -3, -1\right\}$

35. $f(x) = \frac{x}{x+1} \quad g(x) = \frac{x+2}{x}$

$$(g - f)(x) = \frac{x+2}{x} - \frac{x}{x+1}$$

$$= \frac{(x+2)(x+1)}{x(x+1)} - \frac{x(x)}{x(x+1)}$$

$$= \frac{x^2 + 3x + 2}{x(x+1)} - \frac{x^2}{x(x+1)}$$

$$= \frac{x^2 + 3x + 2 - x^2}{x(x+1)}$$

$$= \frac{3x+2}{x(x+1)}$$

Domain: $\{x \mid x \neq -1, x \neq 0\}$


36. a. Domain: $[-3,3]$ Range: $[-2,2]$

b. Intercepts: $(-3,0), (0,0), (3,0)$

c. Symmetric with respect to the orgin.

d. The relation is a function. It passes the vertical line test.

37.



Local maximum: (0,0)
Local Minima: (-2.12,-20.25), (2.12,-20.25)
Increasing: (-2.12,0), (2.12,4)
Decreasing: (-4, -2.12), (0,2.12)

38. $y = \frac{k}{x^2}$

$24 = \frac{k}{5^2} = \frac{k}{25}$

$k = 600$


$y = \frac{600}{x^2}$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Section 2.8

1. $x \geq -2$



2. The distance on a number line from the origin to a is $|a|$ for any real number a.

3. $4x - 3 = 9$

$4x = 12$

$x = 3$

The solution set is $\{3\}$.

4. $3x - 2 > 7$

$3x > 9$

$x > 3$

The solution set is $\{x \mid x > 3\}$ or, using interval notation, $(3, \infty)$.


5. $-1 < 2x + 5 < 13$

$-6 < 2x < 8$

$-3 < x < 4$

The solution set is $\{x \mid -3 < x < 4\}$ or, using interval notation, $(-3, 4)$.

6. To graph $f(x) = |x - 3|$, shift the graph of $y = |x|$ to the right 3 units.



7. $-a$; a

8. $-a < u < a$

9. $\leq$

10. True

11. False. Any real number will be a solution of $|x| > -2$ since the absolute value of any real number is positive.

12. False. $|u| > a$ is equivalent to $u < -a$ or $u > a$.

13. a. Since the graphs of f and g intersect at the points $(-9, 6)$ and $(3, 6)$, the solution set of $f(x) = g(x)$ is $\{-9, 3\}$.

b. Since the graph of f is below the graph of g when x is between -9 and 3, the solution set of $f(x) \leq g(x)$ is $\{x \mid -9 \leq x \leq 3\}$ or, using interval notation, $[-9, 3]$.

c. Since the graph of f is above the graph of g to the left of $x = -9$ and to the right of $x = 3$, the solution set of $f(x) > g(x)$ is $\{x \mid x < -9 \text{ or } x > 3\}$ or, using interval notation, $(-\infty, -9) \cup (3, \infty)$.

14. a. Since the graphs of f and g intersect at the points $(0, 2)$ and $(4, 2)$, the solution set of $f(x) = g(x)$ is $\{0, 4\}$.

b. Since the graph of f is below the graph of g when x is between 0 and 4, the solution set of $f(x) \leq g(x)$ is $\{x \mid 0 \leq x \leq 4\}$ or, using interval notation, $[0, 4]$.

c. Since the graph of f is above the graph of g to the left of $x = 0$ and to the right of $x = 4$, the solution set of $f(x) > g(x)$ is $\{x \mid x < 0 \text{ or } x > 4\}$ or, using interval notation, $(-\infty, 0) \cup (4, \infty)$.

15. a. Since the graphs of f and g intersect at the points $(-2, 5)$ and $(3, 5)$, the solution set of $f(x) = g(x)$ is $\{-2, 3\}$.

b. Since the graph of f is above the graph of g to the left of $x = -2$ and to the right of $x = 3$, the solution set of $f(x) \geq g(x)$ is $\{x \mid x \leq -2 \text{ or } x \geq 3\}$ or, using interval notation, $(-\infty, -2] \cup [3, \infty)$.

c. Since the graph of f is below the graph of g when x is between -2 and 3, the solution set of $f(x) < g(x)$ is $\{x \mid -2 < x < 3\}$ or, using interval notation, $(-2, 3)$.

16. a. Since the graphs of f and g intersect at the points $(-4, 7)$ and $(3, 7)$, the solution set of $f(x) = g(x)$ is $\{-4, 3\}$.

b. Since the graph of f is above the graph of g to the left of $x = -4$ and to the right of $x = 3$, the solution set of $f(x) \geq g(x)$ is $\{x \mid x \leq -4 \text{ or } x \geq 3\}$ or, using interval notation, $(-\infty, -4] \cup [3, \infty)$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

c. Since the graph of f is below the graph of g when x is between -4 and 3, the solution set of $f(x) < g(x)$ is $\{x \mid -4 < x < 3\}$ or, using interval notation, $(-4, 3)$.

17. $|x| = 6$

$x = 6$ or $x = -6$

The solution set is $\{-6, 6\}$.

18. $|x| = 12$

$x = 12$ or $x = -12$

The solution set is $\{-12, 12\}$.

19. $|2x + 3| = 5$

$2x + 3 = 5$ or $2x + 3 = -5$

$2x = 2$ or $2x = -8$

$x = 1$ or $x = -4$

The solution set is $\{-4, 1\}$.

20. $|3x - 1| = 2$

$3x - 1 = 2$ or $3x - 1 = -2$

$3x = 3$ or $3x = -1$

$x = 1$ or $x = -\frac{1}{3}$

The solution set is $\left\{-\frac{1}{3}, 1\right\}$.

21. $|1 - 4t| + 8 = 13$

$|1 - 4t| = 5$

$1 - 4t = 5$ or $1 - 4t = -5$

$-4t = 4$ or $-4t = -6$

$t = -1$ or $t = \frac{3}{2}$

The solution set is $\left\{-1, \frac{3}{2}\right\}$.

22. $|1 - 2z| + 6 = 9$

$|1 - 2z| = 3$

$1 - 2z = 3$ or $1 - 2z = -3$

$-2z = 2$ or $-2z = -4$

$z = -1$ or $z = 2$

The solution set is $\{-1, 2\}$.

23. $|-2x| = 8$

$-2x = 8$ or $-2x = -8$

$x = -4$ or $x = 4$

The solution set is $\{-4, 4\}$.

24. $|-x| = 1$

$-x = 1$ or $-x = -1$

The solution set is $\{-1, 1\}$.

25. $4 - |2x| = 3$

$-|2x| = -1$

$|2x| = 1$

$2x = 1$ or $2x = -1$

$x = \frac{1}{2}$ or $x = -\frac{1}{2}$

The solution set is $\left\{-\frac{1}{2}, \frac{1}{2}\right\}$.

26. $5 - \left|\frac{1}{2}x\right| = 3$

$-\left|\frac{1}{2}x\right| = -2$

$\left|\frac{1}{2}x\right| = 2$

$\frac{1}{2}x = 2$ or $\frac{1}{2}x = -2$

$x = 4$ or $x = -4$

The solution set is $\{-4, 4\}$.

27. $\frac{2}{3}|x| = 9$

$|x| = \frac{27}{2}$

$x = \frac{27}{2}$ or $x = -\frac{27}{2}$

The solution set is $\left\{-\frac{27}{2}, \frac{27}{2}\right\}$.

28. $\frac{3}{4}|x| = 9$

$|x| = 12$

$x = 12$ or $x = -12$

The solution set is $\{-12, 12\}$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

29. $\left|\frac{x}{3}+\frac{2}{5}\right|=2$

$\frac{x}{3}+\frac{2}{5}=2 \quad \text{or} \quad \frac{x}{3}+\frac{2}{5}=-2$

$5x+6=30 \quad \text{or} \quad 5x+6=-30$

$5x=24 \quad \text{or} \quad 5x=-36$

$x=\frac{24}{5} \quad \text{or} \quad x=-\frac{36}{5}$

The solution set is $\left\{-\frac{36}{5}, \frac{24}{5}\right\}$.

30. $\left|\frac{x}{2}-\frac{1}{3}\right|=1$

$\frac{x}{2}-\frac{1}{3}=1 \quad \text{or} \quad \frac{x}{2}-\frac{1}{3}=-1$

$3x-2=6 \quad \text{or} \quad 3x-2=-6$

$3x=8 \quad \text{or} \quad 3x=-4$

$x=\frac{8}{3} \quad \text{or} \quad x=-\frac{4}{3}$

The solution set is $\left\{-\frac{4}{3}, \frac{8}{3}\right\}$.

31. $|u-2|=-\frac{1}{2}$

No solution, since absolute value always yields a non-negative number.

32. $|2-v|=-1$

No solution, since absolute value always yields a non-negative number.

33. $\left|x^2-9\right|=0$

$x^2-9=0$

$x^2=9$

$x=\pm 3$

The solution set is $\{-3, 3\}$.

34. $\left|x^2-16\right|=0$

$x^2-16=0$

$x^2=16$

$x=\pm 4$

The solution set is $\{-4, 4\}$.

35. $\left|x^2-2x\right|=3$

$x^2-2x=3 \quad \text{or} \quad x^2-2x=-3$

$x^2-2x-3=0 \quad \text{or} \quad x^2-2x+3=0$

$(x-3)(x+1)=0 \quad \text{or} \quad x^2-2x+3=0$

$x=\frac{2\pm\sqrt{4-12}}{2}$

$=\frac{2\pm\sqrt{-8}}{2}=1\pm\sqrt{2}i$

$x=3 \text{ or } x=-1$

The solution set is $\left\{-1, 3, 1-\sqrt{2}i, 1+\sqrt{2}i\right\}$.

36. $\left|x^2+x\right|=12$

$x^2+x=12 \text{ or } x^2+x=-12$

$x^2+x-12=0 \quad \text{or} \quad x^2+x+12=0$

$(x-3)(x+4)=0 \quad \text{or} \quad x^2+x+3=0$

$x=\frac{-1\pm\sqrt{1-48}}{2}$

$\frac{-1\pm\sqrt{-47}}{2}=-\frac{1}{2}\pm\frac{\sqrt{47}}{2}i$

$x=3 \text{ or } x=-4$

The solution set is

$\left\{-4, 3, -\frac{1}{2}-\frac{\sqrt{47}}{2}i, -\frac{1}{2}+\frac{\sqrt{47}}{2}i\right\}$.

37. $\left|x^2+x-1\right|=1$

$x^2+x-1=1 \quad \text{or} \quad x^2+x-1=-1$

$x^2+x-2=0 \quad \text{or} \quad x^2+x=0$

$(x-1)(x+2)=0 \quad \text{or} \quad x(x+1)=0$

$x=1, x=-2 \quad \text{or} \quad x=0, x=-1$

The solution set is $\{-2, -1, 0, 1\}$.

38. $\left|x^2+3x-2\right|=2$

$x^2+3x-2=2 \quad \text{or} \quad x^2+3x-2=-2$

$x^2+3x=4 \quad \text{or} \quad x^2+3x=0$

$x^2+3x-4=0 \text{ or } x(x+3)=0$

$(x+4)(x-1)=0 \quad \text{or} \quad x=0, x=-3$

$x=-4, x=1$

The solution set is $\{-4, -3, 0, 1\}$.


Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

39. $|x|<6$

$-6<x<6$


$\{x|-6<x<6\}$ or $(-6,6)$



40. $|x|<9$

$-9<x<9$


$\{x|-9<x<9\}$ or $(-9,9)$



41. $|x|>4$

$x<-4$ or $x>4$


$\{x|x<-4 \text{ or } x>4\}$ or $(-\infty,-4)\cup(4,\infty)$



42. $|x|>1$

$x<-1$ or $x>1$

$\{x|x<-1 \text{ or } x>1\}$ or $(-\infty,-1)\cup(1,\infty)$




43. $|2x|<8$

$-8<2x<8$

$-4<x<4$

$\{x|-4<x<4\}$ or $(-4,4)$



44. $|3x|<15$

$-15<3x<15$

$-5<x<5$

$\{x|-5<x<5\}$ or $(-5,5)$




45. $|3x|>12$

$3x<-12$ or $3x>12$

$x<-4$ or $x>4$

$\{x|x<-4 \text{ or } x>4\}$ or $(-\infty,-4)\cup(4,\infty)$




46. $|2x|>6$

$2x<-6$ or $2x>6$

$x<-3$ or $x>3$

$\{x|x<-3 \text{ or } x>3\}$ or $(-\infty,-3)\cup(3,\infty)$




47. $|x-2|+2<3$

$|x-2|<1$

$-1<x-2<1$

$1<x<3$

$\{x|1<x<3\}$ or $(1,3)$




48. $|x+4|+3<5$

$|x+4|<2$

$-2<x+4<2$

$-6<x<-2$

$\{x|-6<x<-2\}$ or $(-6,-2)$




49. $|3t-2|\le 4$

$-4\le 3t-2\le 4$

$-2\le 3t\le 6$

$-\frac{2}{3}\le t\le 2$

$\left\{t\left|-\frac{2}{3}\le t\le 2\right.\right\}$ or $\left[-\frac{2}{3},2\right]$



50. $|2u+5|\le 7$

$-7\le 2u+5\le 7$

$-12\le 2u\le 2$

$-6\le u\le 1$

$\{u|-6\le u\le 1\}$ or $[-6,1]$



Copyright © 2019 Pearson Education, Inc.


Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

51. $|x-3| \geq 2$

$x-3 \leq -2$ or $x-3 \geq 2$

$x \leq 1$ or $x \geq 5$

$\{x \mid x \leq 1 \text{ or } x \geq 5\}$ or $(-\infty, 1] \cup [5, \infty)$




52. $|x+4| \geq 2$

$x+4 \leq -2$ or $x+4 \geq 2$

$x \leq -6$ or $x \geq -2$

$\{x \mid x \leq -6 \text{ or } x \geq -2\}$ or $(-\infty, -6] \cup [-2, \infty)$



53. $|1-4x|-7 < -2$

$|1-4x| < 5$


$-5 < 1-4x < 5$

$-6 < -4x < 4$

$\frac{-6}{-4} > x > \frac{4}{-4}$

$\frac{3}{2} > x > -1$ or $-1 < x < \frac{3}{2}$

$\left\{x \mid -1 < x < \frac{3}{2}\right\}$ or $\left(-1, \frac{3}{2}\right)$



54. $|1-2x|-4 < -1$

$|1-2x| < 3$


$-3 < 1-2x < 3$

$-4 < -2x < 2$

$\frac{-4}{-2} > x > \frac{2}{-2}$

$2 > x > -1$ or $-1 < x < 2$

$\{x \mid -1 < x < 2\}$ or $(-1, 2)$



55. $|1-2x| > |-3|$


$|1-2x| > 3$

$1-2x < -3$ or $1-2x > 3$

$-2x < -4$ or $-2x > 2$

$x > 2$ or $x < -1$

$\{x \mid x < -1 \text{ or } x > 2\}$ or $(-\infty, -1) \cup (2, \infty)$



56. $|2-3x| > |-1|$


$|2-3x| > 1$

$2-3x < -1$ or $2-3x > 1$

$-3x < -3$ or $-3x > -1$


$x > 1$ or $x < \frac{1}{3}$

$\left\{x \mid x < \frac{1}{3} \text{ or } x > 1\right\}$ or $\left(-\infty, \frac{1}{3}\right) \cup (1, \infty)$



57. $|2x+1| < -1$

No solution since absolute value is always non-negative.



58. $|3x-4| \geq 0$

All real numbers since absolute value is always non-negative.

$\{x \mid x \text{ is any real number}\}$ or $(-\infty, \infty)$



59. $|(3x-2)-7| < \frac{1}{2}$

$|3x-9| < \frac{1}{2}$

$-\frac{1}{2} < 3x-9 < \frac{1}{2}$

$\frac{17}{2} < 3x < \frac{19}{2}$

$\frac{17}{6} < x < \frac{19}{6}$

$\left\{x \mid \frac{17}{6} < x < \frac{19}{6}\right\}$ or $\left(\frac{17}{6}, \frac{19}{6}\right)$




Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

60. $\left|(4x-1)-11\right| < \frac{1}{4}$

$$|4x-12| < \frac{1}{4}$$
$$-\frac{1}{4} < 4x-12 < \frac{1}{4}$$
$$\frac{47}{4} < 4x < \frac{49}{4}$$
$$\frac{47}{16} < x < \frac{49}{16}$$

$\left\{x \mid \frac{47}{16} < x < \frac{49}{16}\right\}$ or $\left(\frac{47}{16}, \frac{49}{16}\right)$



61. $5-|x-1| > 2$

$$-|x-1| > -3$$
$$|x-1| < 3$$
$$-3 < x-1 < 3$$
$$-2 < x < 4$$

$\{x \mid -2 < x < 4\}$ or $(-2, 4)$



62. $6-|x+3| \geq 2$

$$-|x+3| \geq -4$$
$$|x+3| \leq 4$$
$$-4 \leq x+3 \leq 4$$
$$-7 \leq x \leq 1$$

$\{x \mid -7 \leq x \leq 1\}$ or $[-7, 1]$



63. a. $f(x) = g(x)$

$$-3|5x-2| = -9$$
$$|5x-2| = 3$$
$$5x-2 = 3 \quad \text{or} \quad 5x-2 = -3$$
$$5x = 5 \quad \text{or} \quad 5x = -1$$
$$x = 1 \quad \text{or} \quad x = -\frac{1}{5}$$

b. $f(x) > g(x)$

$$-3|5x-2| > -9$$
$$|5x-2| < 3$$
$$-3 < 5x-2 < 3$$
$$-1 < 5x < 5$$
$$-\frac{1}{5} < x < 1$$

$\left\{x \mid -\frac{1}{5} < x < 1\right\}$ or $\left(-\frac{1}{5}, 1\right)$

c. $f(x) \leq g(x)$

$$-3|5x-2| \leq -9$$
$$|5x-2| \geq 3$$
$$5x-2 \geq 3 \quad \text{or} \quad 5x-2 \leq -3$$
$$5x \geq 5 \quad \text{or} \quad 5x \leq -1$$
$$x \geq 1 \quad \text{or} \quad x \leq -\frac{1}{5}$$

$\left\{x \mid x \leq -\frac{1}{5} \text{ or } x \geq 1\right\}$ or $\left(-\infty, -\frac{1}{5}\right] \cup [1, \infty)$

64. a. $f(x) = g(x)$

$$-2|2x-3| = -12$$
$$|2x-3| = 6$$
$$2x-3 = 6 \quad \text{or} \quad 2x-3 = -6$$
$$2x = 9 \quad \text{or} \quad 2x = -3$$
$$x = \frac{9}{2} \quad \text{or} \quad x = -\frac{3}{2}$$

b. $f(x) = g(x)$

$$-2|2x-3| \geq -12$$
$$|2x-3| \leq 6$$
$$-6 \leq 2x-3 \leq 6$$
$$-3 \leq 2x \leq 9$$
$$-\frac{3}{2} \leq x \leq \frac{9}{2}$$

$\left\{x \mid -\frac{3}{2} \leq x \leq \frac{9}{2}\right\}$ or $\left[-\frac{3}{2}, \frac{9}{2}\right]$

c. $f(x) = g(x)$

$$-2|2x-3| < -12$$
$$|2x-3| > 6$$


Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$$2x-3>6 \quad \text{or} \quad 2x-3<-6$$
$$2x>9 \quad \text{or} \quad 2x<-3$$
$$x>\frac{9}{2} \quad \text{or} \quad x<-\frac{3}{2}$$
$$\left\{x \mid x<-\tfrac{3}{2} \text{ or } x>\tfrac{9}{2}\right\} \text{ or } \left(-\infty,-\tfrac{3}{2}\right)\cup\left(\tfrac{9}{2},\infty\right)$$

65. a.
$$f(x)=g(x)$$
$$|-3x+2|=x+10$$
$$-3x+2=x+10 \quad \text{or} \quad -3x+2=-(x+10)$$
$$-4x=8 \quad \text{or} \quad -3x+2=-x-10$$
$$x=-2 \quad \text{or} \quad -2x=-12$$
$$\text{or} \quad x=6$$

b.




Look at the graph of $f(x)$ and $g(x)$ and see where the graph of $f(x)\ge g(x)$. We see that this occurs where $x\le -2$ or $x\ge 6$. So the solution set is: $\{x \mid x\le -2 \text{ or } x\ge 6\}$ or $(-\infty,-2]\cup[6,\infty)$.

c. Look at the graph of $f(x)$ and $g(x)$ and see where the graph of $f(x)<g(x)$. We see that this occurs where x is between -2 and 6. So the solution set is: $\{x \mid -2<x<6\}$ or $(-2,6)$.

66. a.
$$f(x)=g(x)$$
$$|4x-3|=x+2$$
$$4x-3=x+2 \quad 4x-3=-(x+2)$$
$$3x=5 \quad \text{or} \quad 4x-3=-x-2$$
$$x=\frac{5}{3} \quad \text{or} \quad 5x=1$$
$$\text{or} \quad x=\frac{1}{5}$$

b.



Look at the graph of $f(x)$ and $g(x)$ and see where the graph of $f(x)>g(x)$. We see that this occurs where $x<\frac{1}{5}$ or $x>\frac{5}{3}$. So the solution set is: $\left\{x \mid x<\frac{1}{5} \text{ or } x>\frac{5}{3}\right\}$ or $\left(-\infty,\frac{1}{5}\right)\cup\left(\frac{5}{3},\infty\right)$

c. Look at the graph of $f(x)$ and $g(x)$ and see where the graph of $f(x)\le g(x)$. We see that this occurs where x is between $\frac{1}{5}$ and $\frac{5}{3}$. So the solution set is: $\left\{x \mid \frac{1}{5}\le x\le\frac{5}{3}\right\}$ or $\left[\frac{1}{5},\frac{5}{3}\right]$.

67.
$$|x-10|<2$$
$$-2<x-10<2$$
$$8<x<12$$
Solution set: $\{x \mid 8<x<12\}$ or $(8,12)$

68.
$$|x-(-6)|<3$$
$$|x+6|<3$$
$$-3<x+6<3$$
$$-9<x<-3$$
Solution set: $\{x \mid -9<x<-3\}$ or $(-9,-3)$

69.
$$|2x-(-1)|>5$$
$$|2x+1|>5$$
$$2x+1<-5 \quad \text{or} \quad 2x+1>5$$
$$2x<-6 \quad \text{or} \quad 2x>4$$
$$x<-3 \quad \text{or} \quad x>2$$
Solution set: $\{x \mid x<-3 \text{ or } x>2\}$ or $(-\infty,-3)\cup(2,\infty)$

70.
$$|2x-3|>1$$
$$2x-3<-1 \quad \text{or} \quad 2x-3>1$$
$$2x<2 \quad \text{or} \quad 2x>4$$
$$x<1 \quad \text{or} \quad x>2$$
Solution set: $\{x \mid x<1 \text{ or } x>2\}$ or $(-\infty,1)\cup(2,\infty)$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

71. $|x-5.7| \le 0.0005$

$-0.0005 < x-5.7 < 0.0005$

$5.6995 < x < 5.7005$

The acceptable lengths of the rod is from 5.6995 inches to 5.7005 inches.

72. $|x-6.125| \le 0.0005$

$-0.0005 < x-6.125 < 0.0005$

$6.1245 < x < 6.1255$

The acceptable lengths of the rod is from 6.1245 inches to 6.1255 inches.

73. $\left|\frac{x-100}{15}\right| > 1.96$

$\frac{x-100}{15} < -1.96$ or $\frac{x-100}{15} > 1.96$

$x-100 < -29.4$ or $x-100 > 29.4$

$x < 70.6$ or $x > 129.4$

Since IQ scores are whole numbers, any IQ less than 71 or greater than 129 would be considered unusual.

74. $\left|\frac{x-266}{16}\right| > 1.96$

$\frac{x-266}{16} < -1.96$ or $\frac{x-266}{16} > 1.96$

$x-266 < -31.36$ or $x-266 > 31.36$

$x < 234.64$ or $x > 297.36$

Pregnancies less than 235 days long or greater than 297 days long would be considered unusual.

75. $|5x+1|+7 = 5$

$|5x+1| = -2$

No matter what real number is substituted for x, the absolute value expression on the left side of the equation must always be zero or larger. Thus, it can never equal -2.

76. $|2x+5|+3 > 1 \Rightarrow |2x+5| > -2$

No matter what real number is substituted for x, the absolute value expression on the left side of the equation must always be zero or larger. Thus, it will always be larger than -2. Thus, the solution is the set of all real numbers.

77. $|2x-1| \le 0$

No matter what real number is substituted for x, the absolute value expression on the left side of the equation must always be zero or larger. Thus, the only solution to the inequality above will be when the absolute value expression equals 0:

$|2x-1| = 0$

$2x-1 = 0$

$2x = 1$

$x = \frac{1}{2}$

Thus, the solution set is $\left\{\frac{1}{2}\right\}$.

78. $f(x) = |2x-7|$

$f(-4) = |2(-4)-7|$

$= |-8-7| = |-15| = 15$


79. $2(x+4)+x < 4(x+2)$

$2x+8+x < 4x+8$

$3x+8 < 4x+8$

$-x < 0$

$x > 0$



80. $(5-i)(3+2i) =$

$15+10i-3i-2i^2 =$

$15+7i+2 = 17+7i$

81. a. Intercepts: (0,0), (4,0)

b. Domain: $[-2,5]$, Range: $[-2,4]$

c. Increasing: $[3,5]$:Decreasing: $[-2,1]$

Constant: $[1,3]$

d. Neither

Chapter 2 Review Exercises


1. $f(x) = 2x-5$

a. Slope = 2; y-intercept = -5

b. Plot the point $(0,-5)$. Use the slope to find an additional point by moving 1 unit to the right and 2 units up.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual



c. Domain and Range: $(-\infty, \infty)$


d. Average rate of change = slope = 2

e. Increasing

2. $h(x) = \frac{4}{5}x - 6$

a. Slope = $\frac{4}{5}$; y-intercept = -6

b. Plot the point $(0, -6)$. Use the slope to find an additional point by moving 5 units to the right and 4 units up.



c. Domain and Range: $(-\infty, \infty)$


d. Average rate of change = slope = $\frac{4}{5}$

e. Increasing

3. $G(x) = 4$

a. Slope = 0; y-intercept = 4

b. Plot the point (0, 4) and draw a horizontal line through it.



c. Domain: $(-\infty, \infty)$

Range: $\{y \mid y = 4\}$

d. Average rate of change = slope = 0

e. Constant


4. $f(x) = 2x + 14$

zero: $f(x) = 2x + 14 = 0$

$2x = -14$

$x = -7$

y-intercept = 14



5.

x	$y = f(x)$	Avg. rate of change $= \frac{\Delta y}{\Delta x}$
−2	−7	
0	3	$\frac{3-(-7)}{0-(-2)} = \frac{10}{2} = 5$
1	8	$\frac{8-3}{1-0} = \frac{5}{1} = 5$
3	18	$\frac{18-8}{3-1} = \frac{10}{2} = 5$
6	33	$\frac{33-18}{6-3} = \frac{15}{3} = 5$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

This is a linear function with slope = 5, since the average rate of change is constant at 5. To find the equation of the line, we use the point-slope formula and one of the points.

$$y - y_1 = m(x - x_1)$$
$$y - 3 = 5(x - 0)$$
$$y = 5x + 3$$

6.

x	$y = f(x)$	Avg. rate of change $= \frac{\Delta y}{\Delta x}$
–1	–3	
0	4	$\frac{4-(-3)}{0-(-1)} = \frac{7}{1} = 7$
1	7	$\frac{7-4}{1-0} = \frac{3}{1} = 3$
2	6	
3	1	

This is not a linear function, since the average rate of change is not constant.

7.
$$f(x) = 0$$
$$x^2 + x - 72 = 0$$
$$(x+9)(x-8) = 0$$
$$x + 9 = 0 \quad \text{or} \quad x - 8 = 0$$
$$x = -9 \qquad x = 8$$

The zeros of $f(x) = x^2 + x - 72$ are -9 and 8.

The x-intercepts of the graph of f are -9 and 8.

8.
$$P(t) = 0$$
$$6t^2 - 13t - 5 = 0$$
$$(3t+1)(2t-5) = 0$$
$$3t + 1 = 0 \quad \text{or} \quad 2t - 5 = 0$$
$$t = -\frac{1}{3} \qquad t = \frac{5}{2}$$

The zeros of $P(t) = 6t^2 - 13t - 5$ are $-\frac{1}{3}$ and $\frac{5}{2}$.

The t-intercepts of the graph of P are $-\frac{1}{3}$ and $\frac{5}{2}$.

9.
$$g(x) = 0$$
$$(x-3)^2 - 4 = 0$$
$$(x-3)^2 = 4$$
$$x - 3 = \pm\sqrt{4}$$
$$x - 3 = \pm 2$$
$$x = 3 \pm 2$$
$$x = 3 - 2 = 1 \quad \text{or} \quad x = 3 + 2 = 5$$

The zeros of $g(x) = (x-3)^2 - 4$ are 1 and 5. The x-intercepts of the graph of g are 1 and 5.

10.
$$h(x) = 0$$
$$9x^2 + 6x + 1 = 0$$
$$(3x+1)(3x+1) = 0$$
$$3x + 1 = 0 \quad \text{or} \quad 3x + 1 = 0$$
$$x = -\frac{1}{3} \qquad x = -\frac{1}{3}$$

The only zero of $h(x) = 9x^2 + 6x + 1$ is $-\frac{1}{3}$.

The only x-intercept of the graph of h is $-\frac{1}{3}$.

11.
$$G(x) = 0$$
$$2x^2 - 4x - 1 = 0$$
$$x^2 - 2x - \frac{1}{2} = 0$$
$$x^2 - 2x = \frac{1}{2}$$
$$x^2 - 2x + 1 = \frac{1}{2} + 1$$
$$(x-1)^2 = \frac{3}{2}$$
$$x - 1 = \pm\sqrt{\frac{3}{2}} = \pm\frac{\sqrt{3}}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \pm\frac{\sqrt{6}}{2}$$
$$x = 1 \pm \frac{\sqrt{6}}{2} = \frac{2 \pm \sqrt{6}}{2}$$

The zeros of $G(x) = 2x^2 - 4x - 1$ are $\frac{2-\sqrt{6}}{2}$ and $\frac{2+\sqrt{6}}{2}$. The x-intercepts of the graph of G are $\frac{2-\sqrt{6}}{2}$ and $\frac{2+\sqrt{6}}{2}$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

12. $f(x)=0$

$-2x^2+x+1=0$

$2x^2-x-1=0$

$(2x+1)(x-1)=0$

$2x+1=0 \quad \text{or} \quad x-1=0$

$x=-\frac{1}{2} \qquad x=1$

The zeros of $f(x)=-2x^2+x+1$ are $-\frac{1}{2}$ and 1.

The x-intercepts of the graph of f are $-\frac{1}{2}$ and 1.

13. $f(x)=g(x)$

$(x-3)^2=16$


$x-3=\pm\sqrt{16}=\pm 4$

$x=3\pm 4$

$x=3-4=-1 \quad \text{or} \quad x=3+4=7$

The solution set is $\{-1, 7\}$.

The x-coordinates of the points of intersection are -1 and 7. The y-coordinates are $g(-1)=16$ and $g(7)=16$. The graphs of the f and g intersect at the points $(-1, 16)$ and $(7, 16)$.



14. $f(x)=g(x)$

$x^2+4x-5=4x-1$

$x^2-4=0$


$(x+2)(x-2)=0$

$x+2=0 \quad \text{or} \quad x-2=0$

$x=-2 \qquad x=2$

The solution set is $\{-2, 2\}$.

The x-coordinates of the points of intersection are -2 and 2. The y-coordinates are $g(-2)=4(-2)-1=-8-1=-9$ and $g(2)=4(2)-1=8-1=7$. The graphs of the f and g intersect at the points $(-2,-9)$ and $(2, 7)$.



15. $f(x)=0$

$x^4-5x^2+4=0$

$(x^2-4)(x^2-1)=0$

$x^2-4=0 \quad \text{or} \quad x^2-1=0$

$x=\pm 2 \quad \text{or} \quad x=\pm 1$

The zeros of $f(x)=x^4-5x^2+4$ are -2, -1, 1, and 2. The x-intercepts of the graph of f are -2, -1, 1, and 2.

16. $F(x)=0$

$(x-3)^2-2(x-3)-48=0$

Let $u=x-3 \rightarrow u^2=(x-3)^2$

$u^2-2u-48=0$

$(u+6)(u-8)=0$

$u+6=0 \quad \text{or} \quad u-8=0$

$u=-6 \qquad u=8$

$x-3=-6 \qquad x-3=8$

$x=-3 \qquad x=11$

The zeros of $F(x)=(x-3)^2-2(x-3)-48$ are -3 and 11. The x-intercepts of the graph of F are -3 and 11.

17. $h(x)=0$

$3x-13\sqrt{x}-10=0$

Let $u=\sqrt{x} \rightarrow u^2=x$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$3u^2 - 13u - 10 = 0$

$(3u + 2)(u - 5) = 0$

$3u + 2 = 0$ or $u - 5 = 0$

$u = -\frac{2}{3}$ $\quad$ $u = 5$

$\sqrt{x} = -\frac{2}{3}$ $\quad$ $\sqrt{x} = 5$

$x =$ not real $\quad$ $x = 5^2 = 25$

Check: $h(25) = 3(25) - 13\sqrt{25} - 10$

$= 3(25) - 13(5) - 10$

$= 75 - 65 - 10 = 0$

The only zero of $h(x) = 3x - 13\sqrt{x} - 10$ is 25.

The only x-intercept of the graph of h is 25.

18. $f(x) = 0$

$\left(\frac{1}{x}\right)^2 - 4\left(\frac{1}{x}\right) - 12 = 0$

Let $u = \frac{1}{x} \rightarrow u^2 = \left(\frac{1}{x}\right)^2$

$u^2 - 4u - 12 = 0$

$(u + 2)(u - 6) = 0$

$u + 2 = 0$ or $u - 6 = 0$

$u = -2$ $\quad$ $u = 6$


$\frac{1}{x} = -2$ $\quad$ $\frac{1}{x} = 6$

$x = -\frac{1}{2}$ $\quad$ $x = \frac{1}{6}$

The zeros of $f(x) = \left(\frac{1}{x}\right)^2 - 4\left(\frac{1}{x}\right) - 12$ are $-\frac{1}{2}$ and $\frac{1}{6}$. The x-intercepts of the graph of f are $-\frac{1}{2}$ and $\frac{1}{6}$.


19. $f(x) = (x - 2)^2 + 2$

Using the graph of $y = x^2$, shift right 2 units, then shift up 2 units.




20. $f(x) = -(x - 4)^2$

Using the graph of $y = x^2$, shift the graph 4 units right, then reflect about the x-axis.



21. $f(x) = 2(x + 1)^2 + 4$

Using the graph of $y = x^2$, stretch vertically by a factor of 2, then shift 1 unit left, then shift 4 units up.



22. a. $f(x) = (x - 2)^2 + 2$

$= x^2 - 4x + 4 + 2$

$= x^2 - 4x + 6$


$a = 1, b = -4, c = 6$. Since $a = 1 > 0$, the graph opens up. The x-coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{-4}{2(1)} = \frac{4}{2} = 2$.

The y-coordinate of the vertex is $f\left(-\frac{b}{2a}\right) = f(2) = (2)^2 - 4(2) + 6 = 2$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Thus, the vertex is (2, 2).
The axis of symmetry is the line $x = 2$.
The discriminant is:
$b^2 - 4ac = (-4)^2 - 4(1)(6) = -8 < 0$, so the graph has no x-intercepts.
The y-intercept is $f(0) = 6$.



b. Domain: $(-\infty, \infty)$. Range: $[2, \infty)$.

c. Decreasing on $(-\infty, 2]$; increasing on $[2, \infty)$.


23. a. $f(x) = \frac{1}{4}x^2 - 16$

$a = \frac{1}{4}, b = 0, c = -16$. Since $a = \frac{1}{4} > 0$, the graph opens up. The x-coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{-0}{2\left(\frac{1}{4}\right)} = -\frac{0}{\frac{1}{2}} = 0$.

The y-coordinate of the vertex is
$f\left(-\frac{b}{2a}\right) = f(0) = \frac{1}{4}(0)^2 - 16 = -16$.
Thus, the vertex is (0, –16).
The axis of symmetry is the line $x = 0$.
The discriminant is:
$b^2 - 4ac = (0)^2 - 4\left(\frac{1}{4}\right)(-16) = 16 > 0$, so the graph has two x-intercepts.
The x-intercepts are found by solving:

$$\frac{1}{4}x^2 - 16 = 0$$
$$x^2 - 64 = 0$$
$$x^2 = 64$$
$$x = 8 \text{ or } x = -8$$

The x-intercepts are –8 and 8.
The y-intercept is $f(0) = -16$.



b. Domain: $(-\infty, \infty)$. Range: $[-16, \infty)$.

c. Decreasing on $(-\infty, 0]$; increasing on $[0, \infty)$.

24. a. $f(x) = -4x^2 + 4x$

$a = -4, b = 4, c = 0$. Since $a = -4 < 0$, the graph opens down. The x-coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{4}{2(-4)} = -\frac{4}{-8} = \frac{1}{2}$.

The y-coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f\left(\frac{1}{2}\right) = -4\left(\frac{1}{2}\right)^2 + 4\left(\frac{1}{2}\right)$$
$$= -1 + 2 = 1$$

Thus, the vertex is $\left(\frac{1}{2}, 1\right)$.


The axis of symmetry is the line $x = \frac{1}{2}$.
The discriminant is:
$b^2 - 4ac = 4^2 - 4(-4)(0) = 16 > 0$, so the graph has two x-intercepts.
The x-intercepts are found by solving:

$$-4x^2 + 4x = 0$$
$$-4x(x - 1) = 0$$
$$x = 0 \text{ or } x = 1$$

The x-intercepts are 0 and 1.
The y-intercept is $f(0) = -4(0)^2 + 4(0) = 0$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual



b. Domain: $(-\infty, \infty)$. Range: $(-\infty, 1]$.

c. Increasing on $\left(-\infty, \frac{1}{2}\right]$; decreasing on $\left[\frac{1}{2}, \infty\right)$.

25. a. $f(x) = \frac{9}{2}x^2 + 3x + 1$

$a = \frac{9}{2}, b = 3, c = 1$. Since $a = \frac{9}{2} > 0$, the graph opens up. The x-coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{3}{2\left(\frac{9}{2}\right)} = -\frac{3}{9} = -\frac{1}{3}$.

The y-coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f\left(-\frac{1}{3}\right) = \frac{9}{2}\left(-\frac{1}{3}\right)^2 + 3\left(-\frac{1}{3}\right) + 1$$
$$= \frac{1}{2} - 1 + 1 = \frac{1}{2}$$


Thus, the vertex is $\left(-\frac{1}{3}, \frac{1}{2}\right)$.

The axis of symmetry is the line $x = -\frac{1}{3}$.

The discriminant is:

$b^2 - 4ac = 3^2 - 4\left(\frac{9}{2}\right)(1) = 9 - 18 = -9 < 0$,

so the graph has no x-intercepts. The y-intercept is $f(0) = \frac{9}{2}(0)^2 + 3(0) + 1 = 1$.



b. Domain: $(-\infty, \infty)$. Range: $\left[\frac{1}{2}, \infty\right)$.

c. Decreasing on $\left(-\infty, -\frac{1}{3}\right]$; increasing on $\left[-\frac{1}{3}, \infty\right)$.

26. a. $f(x) = 3x^2 + 4x - 1$

$a = 3, b = 4, c = -1$. Since $a = 3 > 0$, the graph opens up. The x-coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{4}{2(3)} = -\frac{4}{6} = -\frac{2}{3}$.

The y-coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f\left(-\frac{2}{3}\right) = 3\left(-\frac{2}{3}\right)^2 + 4\left(-\frac{2}{3}\right) - 1$$
$$= \frac{4}{3} - \frac{8}{3} - 1 = -\frac{7}{3}$$

Thus, the vertex is $\left(-\frac{2}{3}, -\frac{7}{3}\right)$.

The axis of symmetry is the line $x = -\frac{2}{3}$.

The discriminant is:

$b^2 - 4ac = (4)^2 - 4(3)(-1) = 28 > 0$, so the graph has two x-intercepts.

The x-intercepts are found by solving:

$3x^2 + 4x - 1 = 0$.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \pm \sqrt{28}}{2(3)}$$
$$= \frac{-4 \pm 2\sqrt{7}}{6} = \frac{-2 \pm \sqrt{7}}{3}$$


The x-intercepts are $\frac{-2 - \sqrt{7}}{3} \approx -1.55$ and

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$\frac{-2+\sqrt{7}}{3} \approx 0.22$.

The y-intercept is $f(0) = 3(0)^2 + 4(0) - 1 = -1$.



b. Domain: $(-\infty, \infty)$. Range: $\left[-\frac{7}{3}, \infty\right)$.

c. Decreasing on $\left(-\infty, -\frac{2}{3}\right]$; increasing on $\left[-\frac{2}{3}, \infty\right)$.

27. $f(x) = 3x^2 - 6x + 4$

$a = 3, b = -6, c = 4$. Since $a = 3 > 0$, the graph opens up, so the vertex is a minimum point. The minimum occurs at

$$x = -\frac{b}{2a} = -\frac{-6}{2(3)} = \frac{6}{6} = 1.$$

The minimum value is

$$f\left(-\frac{b}{2a}\right) = f(1) = 3(1)^2 - 6(1) + 4$$
$$= 3 - 6 + 4 = 1$$

28. $f(x) = -x^2 + 8x - 4$

$a = -1, b = 8, c = -4$. Since $a = -1 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at

$$x = -\frac{b}{2a} = -\frac{8}{2(-1)} = -\frac{8}{-2} = 4.$$

The maximum value is

$$f\left(-\frac{b}{2a}\right) = f(4) = -(4)^2 + 8(4) - 4$$
$$= -16 + 32 - 4 = 12$$

29. $f(x) = -3x^2 + 12x + 4$

$a = -3, b = 12, c = 4$. Since $a = -3 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at

$$x = -\frac{b}{2a} = -\frac{12}{2(-3)} = -\frac{12}{-6} = 2.$$

The maximum value is

$$f\left(-\frac{b}{2a}\right) = f(2) = -3(2)^2 + 12(2) + 4$$
$$= -12 + 24 + 4 = 16$$

30. Consider the form $y = a(x-h)^2 + k$. The vertex is $(2, -4)$ so we have $h = 2$ and $k = -4$. The function also contains the point $(0, -16)$. Substituting these values for x, y, h, and k, we can solve for a:

$$-16 = a(0 - (2))^2 + (-4)$$
$$-16 = a(-2)^2 - 4$$
$$-16 = 4a - 4$$
$$-12 = 4a$$
$$a = -3$$

The quadratic function is

$$f(x) = -3(x-2)^2 - 4 = -3x^2 + 12x - 16.$$

31. Use the form $f(x) = a(x-h)^2 + k$.

The vertex is $(-1, 2)$, so $h = -1$ and $k = 2$.

$f(x) = a(x+1)^2 + 2$.

Since the graph passes through $(1, 6)$, $f(1) = 6$.

$$6 = a(1+1)^2 + 2$$
$$6 = a(2)^2 + 2$$
$$6 = 4a + 2$$
$$4 = 4a$$
$$1 = a$$
$$f(x) = (x+1)^2 + 2$$
$$= (x^2 + 2x + 1) + 2$$
$$= x^2 + 2x + 3$$

32. $x^2 + 6x - 16 < 0$

$f(x) = x^2 + 6x - 16$

$x^2 + 6x - 16 = 0$

$(x+8)(x-2) = 0$

$x = -8, x = 2$ are the zeros of f.

Interval	$(-\infty, -8)$	$(-8, 2)$	$(2, \infty)$
Test Number	−9	0	3
Value of f	11	−16	11
Conclusion	Positive	Negative	Positive

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

The solution set is $\{x \mid -8 < x < 2\}$ or, using interval notation, $(-8, 2)$.

33. $3x^2 \geq 14x + 5$

$3x^2 - 14x - 5 \geq 0$

$f(x) = 3x^2 - 14x - 5$

$3x^2 - 14x - 5 = 0$

$(3x + 1)(x - 5) = 0$

$x = -\frac{1}{3}$, $x = 5$ are the zeros of f.

Interval	$\left(-\infty, -\frac{1}{3}\right)$	$\left(-\frac{1}{3}, 5\right)$	$(5, \infty)$
Test Number	-1	0	2
Value of f	12	-5	19
Conclusion	Positive	Negative	Positive

The solution set is $\left\{x \middle| x \leq -\frac{1}{3} \text{ or } x \geq 5\right\}$ or, using interval notation, $\left(-\infty, -\frac{1}{3}\right] \cup [5, \infty)$.


34. $f(x) = 0$

$x^2 + 8 = 0$

$x^2 = -8$

$x = \pm\sqrt{-8} = \pm 2\sqrt{2}\, i$

The zero are $-2\sqrt{2}\, i$ and $2\sqrt{2}\, i$.



35. $g(x) = 0$


$x^2 + 2x - 4 = 0$

$a = 1, b = 2, c = -4$

$b^2 - 4ac = 2^2 - 4(1)(-4) = 4 + 16 = 20$

$x = \frac{-2 \pm \sqrt{20}}{2(1)} = \frac{-2 \pm 2\sqrt{5}}{2} = -1 \pm \sqrt{5}$

The zeros are $-1 - \sqrt{5}$ and $-1 + \sqrt{5}$.



36. $p(x) = 0$


$-2x^2 + 4x - 3 = 0$

$a = -2, b = 4, c = -3$

$b^2 - 4ac = 4^2 - 4(-2)(-3) = 16 - 24 = -8$

$x = \frac{-4 \pm \sqrt{-8}}{2(-2)} = \frac{-4 \pm 2\sqrt{2}\, i}{-4} = 1 \pm \frac{\sqrt{2}}{2} i$

The zeros are $1 - \frac{\sqrt{2}}{2} i$ and $1 + \frac{\sqrt{2}}{2} i$.



37. $f(x) = 0$

$4x^2 + 4x + 3 = 0$

$a = 4, b = 4, c = 3$


$b^2 - 4ac = 4^2 - 4(4)(3) = 16 - 48 = -32$

$x = \frac{-4 \pm \sqrt{-32}}{2(4)} = \frac{-4 \pm 4\sqrt{2}\, i}{8} = -\frac{1}{2} \pm \frac{\sqrt{2}}{2} i$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

The zeros are $-\frac{1}{2}-\frac{\sqrt{2}}{2}i$ and $-\frac{1}{2}+\frac{\sqrt{2}}{2}i$.



38. $|2x+3|=7$

$2x+3=7$ or $2x+3=-7$

$2x=4$ or $2x=-10$

$x=2$ or $x=-5$

The solution set is $\{-5, 2\}$.

39. $|2-3x|+2=9$

$|2-3x|=7$

$2-3x=7$ or $2-3x=-7$

$-3x=5$ or $-3x=-9$

$x=-\frac{5}{3}$ or $x=3$

The solution set is $\left\{-\frac{5}{3}, 3\right\}$.

40. $|3x+4|<\frac{1}{2}$

$-\frac{1}{2}<3x+4<\frac{1}{2}$

$-\frac{9}{2}<3x<-\frac{7}{2}$

$-\frac{3}{2}<x<-\frac{7}{6}$

$\left\{x\middle|-\frac{3}{2}<x<-\frac{7}{6}\right\}$ or $\left(-\frac{3}{2}, -\frac{7}{6}\right)$




41. $|2x-5|\geq 9$

$2x-5\leq -9$ or $2x-5\geq 9$

$2x\leq -4$ or $2x\geq 14$

$x\leq -2$ or $x\geq 7$

$\{x|x\leq -2 \text{ or } x\geq 7\}$ or $(-\infty,-2]\cup[7,\infty)$



42. $2+|2-3x|\leq 4$

$|2-3x|\leq 2$

$-2\leq 2-3x\leq 2$

$-4\leq -3x\leq 0$

$\frac{4}{3}\geq x\geq 0$

$\left\{x\middle|0\leq x\leq\frac{4}{3}\right\}$ or $\left[0, \frac{4}{3}\right]$



43. $1-|2-3x|<-4$

$-|2-3x|<-5$

$|2-3x|>5$


$2-3x<-5$ or $2-3x>5$

$7<3x$ or $-3>3x$

$\frac{7}{3}<x$ or $-1>x$

$x<-1$ or $x>\frac{7}{3}$

$\left\{x\middle|x<-1 \text{ or } x>\frac{7}{3}\right\}$ or $(-\infty,-1)\cup\left(\frac{7}{3},\infty\right)$



44. a. $S(x)=0.01x+25{,}000$

b. $S(1{,}000{,}000)=0.01(1{,}000{,}000)+25{,}000$

$=10{,}000+25{,}000=35{,}000$

Bill's salary would be \$35,000.

c. $0.01x+25{,}000=100{,}000$

$0.01x=75{,}000$

$x=7{,}500{,}000$

Bill's sales would have to be \$7,500,000 in order to earn \$100,000.

d. $0.01x+25{,}000>150{,}000$

$0.01x>125{,}000$

$x>12{,}500{,}000$

Bill's sales would have to be more than

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$12,500,000 in order for his salary to exceed $150,000.

45. a. If $x = 1500 - 10p$, then $p = \dfrac{1500 - x}{10}$.

$R(p) = px = p(1500 - 10p) = -10p^2 + 1500p$

b. Domain: $\{p \mid 0 < p \le 150\}$

c. $p = \dfrac{-b}{2a} = \dfrac{-1500}{2(-10)} = \dfrac{-1500}{-20} = \$75$


d. The maximum revenue is

$R(75) = -10(75)^2 + 1500(75)$

$= -56250 + 112500 = \$56,250$

e. $x = 1500 - 10(75) = 1500 - 750 = 750$

f. Graph $R = -10p^2 + 1500p$ and $R = 56000$.



Find where the graphs intersect by solving $56000 = -10p^2 + 1500p$.

$10p^2 - 1500p + 56000 = 0$


$p^2 - 150p + 5600 = 0$

$(p - 70)(p - 80) = 0$

$p = 70, p = 80$

The company should charge between $70 and $80.

46. Let w = the width. Then $w + 2$ = the length.



By the Pythagorean Theorem we have:

$w^2 + (w+2)^2 = (10)^2$

$w^2 + w^2 + 4w + 4 = 100$

$2w^2 + 4w - 96 = 0$

$w^2 + 2w - 48 = 0$

$(w+8)(w-6) = 0$

$w = -8$ or $w = 6$

Disregard the negative answer because the width of a rectangle must be positive. Thus, the width is 6 inches, and the length is 8 inches

47. $C(x) = 4.9x^2 - 617.4x + 19,600$;

$a = 4.9, b = -617.4, c = 19,600$. Since $a = 4.9 > 0$, the graph opens up, so the vertex is a minimum point.

a. The minimum marginal cost occurs at

$x = -\dfrac{b}{2a} = -\dfrac{-617.40}{2(4.9)} = \dfrac{617.40}{9.8} = 63$.

Thus, 63 golf clubs should be manufactured in order to minimize the marginal cost.

b. The minimum marginal cost is

$C\left(-\dfrac{b}{2a}\right) = C(63)$

$= 4.9(63)^2 - (617.40)(63) + 19600$

$= \$151.90$

48. Since there are 200 feet of border, we know that $2x + 2y = 200$. The area is to be maximized, so $A = x \cdot y$. Solving the perimeter formula for y:

$2x + 2y = 200$

$2y = 200 - 2x$

$y = 100 - x$

The area function is:

$A(x) = x(100 - x) = -x^2 + 100x$

The maximum value occurs at the vertex:

$x = -\dfrac{b}{2a} = -\dfrac{100}{2(-1)} = -\dfrac{100}{-2} = 50$

The pond should be 50 feet by 50 feet for maximum area.

49. The area function is:

$A(x) = x(10 - x) = -x^2 + 10x$

The maximum value occurs at the vertex:

$x = -\dfrac{b}{2a} = -\dfrac{10}{2(-1)} = -\dfrac{10}{-2} = 5$


The maximum area is:

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$A(5) = -(5)^2 + 10(5)$

$= -25 + 50 = 25$ square units



50. Locate the origin at the point directly under the highest point of the arch. Then the equation is in the form: $y = -ax^2 + k$, where $a > 0$. Since the maximum height is 10 feet, when $x = 0$, $y = k = 10$. Since the point (10, 0) is on the parabola, we can find the constant:

$0 = -a(10)^2 + 10$

$a = \dfrac{10}{10^2} = \dfrac{1}{10} = 0.10$


The equation of the parabola is:

$y = -\dfrac{1}{10}x^2 + 10$

At $x = 8$:

$y = -\dfrac{1}{10}(8)^2 + 10 = -6.4 + 10 = 3.6$ feet

51. a.



b.

p	q	Avg. rate of change $= \dfrac{\Delta q}{\Delta p}$
75	100	
100	80	$\dfrac{80-100}{100-75} = \dfrac{-20}{25} = -0.8$
125	60	$\dfrac{60-80}{125-100} = \dfrac{-20}{25} = -0.8$
150	40	$\dfrac{40-60}{150-125} = \dfrac{-20}{25} = -0.8$

Since each input (price) corresponds to a single output (quantity demanded), we know that the quantity demanded is a function of price. Also, because the average rate of change is constant at $-\$0.8$ per LCD monitor, the function is linear.

c. From part (b), we know $m = -0.8$. Using $(p_1, q_1) = (75, 100)$, we get the equation:

$q - q_1 = m(p - p_1)$

$q - 100 = -0.8(p - 75)$

$q - 100 = -0.8p + 60$

$q = -0.8p + 160$

Using function notation, we have $q(p) = -0.8p + 160$.

d. The price cannot be negative, so $p \geq 0$. Likewise, the quantity cannot be negative, so, $q(p) \geq 0$.


$-0.8p + 160 \geq 0$

$-0.8p \geq -160$

$p \leq 200$

Thus, the implied domain for $q(p)$ is $\{p \mid 0 \leq p \leq 200\}$ or $[0, 200]$.

e.




f. If the price increases by \$1, then the quantity demanded of LCD monitors decreases by 0.8 monitor.

g. p-intercept: If the price is \$0, then 160 LCD monitors will be demanded.
q-intercept: There will be 0 LCD monitors demanded when the price is \$200.

Copyright © 2019 Pearson Education, Inc.


Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

52. a.



b. Yes, the two variables appear to have a linear relationship.


c. Using the LINear REGression program, the line of best fit is:
$y = 1.390171918x + 1.113952697$



d. $y = 1.390171918(26.5) + 1.113952697$
≈ 38.0 mm


53. a.

The data appear to be quadratic



with $a < 0$.

b. Using the QUADratic REGression program, the quadratic function of best fit is:
$y = -7.76x^2 + 411.88x + 942.72$.



The maximum revenue occurs at

$$A = \frac{-b}{2a} = \frac{-(411.88)}{2(-7.76)}$$

$$= \frac{-411.88}{-15.52} \approx \$26.5 \text{ thousand}$$


c. The maximum revenue is

$$R\left(\frac{-b}{2a}\right) = R(26.53866)$$

$$= -7.76(26.5)^2 + (411.88)(26.5) + 942.72$$

$\approx \$6408$ thousand

d.




Chapter 2 Test

1. $f(x) = -4x + 3$

a. The slope f is -4.

b. The slope is negative, so the graph is decreasing.

c. Plot the point $(0, 3)$. Use the slope to find an additional point by moving 1 unit to the right and 4 units down.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

2.

x	y	Avg. rate of change $= \frac{\Delta y}{\Delta x}$
-2	12	
-1	7	$\frac{7-12}{-1-(-2)} = \frac{-5}{1} = -5$
0	2	$\frac{2-7}{0-(-1)} = \frac{-5}{1} = -5$
1	-3	$\frac{-3-2}{1-0} = \frac{-5}{1} = -5$
2	-8	$\frac{-8-(-3)}{2-1} = \frac{-5}{1} = -5$

Since the average rate of change is constant at -5, this is a linear function with slope $= -5$. To find the equation of the line, we use the point-slope formula and one of the points.

$$y - y_1 = m(x - x_1)$$
$$y - 2 = -5(x - 0)$$
$$y = -5x + 2$$

3. $f(x) = 0$

$$3x^2 - 2x - 8 = 0$$
$$(3x + 4)(x - 2) = 0$$
$$3x + 4 = 0 \quad \text{or} \quad x - 2 = 0$$
$$x = -\frac{4}{3} \qquad x = 2$$

The zeros of f are $-\frac{4}{3}$ and 2.

4. $G(x) = 0$


$$-2x^2 + 4x + 1 = 0$$
$$a = -2, b = 4, c = 1$$
$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \pm \sqrt{4^2 - 4(-2)(1)}}{2(-2)}$$
$$= \frac{-4 \pm \sqrt{24}}{-4} = \frac{-4 \pm 2\sqrt{6}}{-4} = \frac{2 \pm \sqrt{6}}{2}$$

The zeros of G are $\frac{2-\sqrt{6}}{2}$ and $\frac{2+\sqrt{6}}{2}$.

5. $f(x) = g(x)$

$$x^2 + 3x = 5x + 3$$
$$x^2 - 2x - 3 = 0$$
$$(x + 1)(x - 3) = 0$$
$$x + 1 = 0 \quad \text{or} \quad x - 3 = 0$$
$$x = -1 \qquad x = 3$$

The solution set is $\{-1, 3\}$.



6. $f(x) = 0$

$$(x - 1)^2 + 5(x - 1) + 4 = 0$$


Let $u = x - 1 \rightarrow u^2 = (x - 1)^2$

$$u^2 + 5u + 4 = 0$$
$$(u + 4)(u + 1) = 0$$
$$u + 4 = 0 \quad \text{or} \quad u + 1 = 0$$
$$u = -4 \qquad u = -1$$
$$x - 1 = -4 \qquad x - 1 = -1$$
$$x = -3 \qquad x = 0$$

The zeros of G are -3 and 0.

7. $f(x) = (x - 3)^2 - 2$

Using the graph of $y = x^2$, shift right 3 units, then shift down 2 units.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

8. a. $f(x) = 3x^2 - 12x + 4$

$a = 3, b = -12, c = 4$. Since $a = 3 > 0$, the graph opens up.

b. The x-coordinate of the vertex is

$$x = -\frac{b}{2a} = -\frac{-12}{2(3)} = -\frac{-12}{6} = 2.$$

The y-coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f(2) = 3(2)^2 - 12(2) + 4$$
$$= 12 - 24 + 4 = -8$$

Thus, the vertex is $(2, -8)$.

c. The axis of symmetry is the line $x = 2$.


d. The discriminant is:

$b^2 - 4ac = (-12)^2 - 4(3)(4) = 96 > 0$, so the graph has two x-intercepts. The x-intercepts are found by solving: $3x^2 - 12x + 4 = 0$.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-12) \pm \sqrt{96}}{2(3)}$$
$$= \frac{12 \pm 4\sqrt{6}}{6} = \frac{6 \pm 2\sqrt{6}}{3}$$

The x-intercepts are $\frac{6 - 2\sqrt{6}}{3} \approx 0.37$ and $\frac{6 + 2\sqrt{6}}{3} \approx 3.63$. The y-intercept is $f(0) = 3(0)^2 - 12(0) + 4 = 4$.

e.



f. The domain is $(-\infty, \infty)$.

The range is $[-8, \infty)$.

g. Decreasing on $(-\infty, 2]$.

Increasing on $[2, \infty)$.

9. a. $g(x) = -2x^2 + 4x - 5$

$a = -2, b = 4, c = -5$. Since $a = -2 < 0$, the graph opens down.

b. The x-coordinate of the vertex is

$$x = -\frac{b}{2a} = -\frac{4}{2(-2)} = -\frac{4}{-4} = 1.$$

The y-coordinate of the vertex is

$$g\left(-\frac{b}{2a}\right) = g(1) = -2(1)^2 + 4(1) - 5$$
$$= -2 + 4 - 5 = -3$$


Thus, the vertex is $(1, -3)$.

c. The axis of symmetry is the line $x = 1$.

d. The discriminant is:

$b^2 - 4ac = (4)^2 - 4(-2)(-5) = -24 < 0$, so the graph has no x-intercepts. The y-intercept is $g(0) = -2(0)^2 + 4(0) - 5 = -5$.

e.



f. The domain is $(-\infty, \infty)$.

The range is $(-\infty, -3]$.

g. Increasing on $(-\infty, 1]$.

Decreasing on $[1, \infty)$.

10. Consider the form $y = a(x - h)^2 + k$. From the graph we know that the vertex is $(1, -32)$ so we have $h = 1$ and $k = -32$. The graph also passes through the point $(x, y) = (0, -30)$. Substituting these values for x, y, h, and k, we can solve for a:

$-30 = a(0 - 1)^2 + (-32)$ The quadratic function is

$-30 = a(-1)^2 - 32$

$-30 = a - 32$

$2 = a$

$f(x) = 2(x - 1)^2 - 32 = 2x^2 - 4x - 30$.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

11. $f(x) = -2x^2 + 12x + 3$

$a = -2, b = 12, c = 3$. Since $a = -2 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at

$$x = -\frac{b}{2a} = -\frac{12}{2(-2)} = -\frac{12}{-4} = 3.$$

The maximum value is

$$f(3) = -2(3)^2 + 12(3) + 3 = -18 + 36 + 3 = 21.$$

12. $x^2 - 10x + 24 \ge 0$

$f(x) = x^2 - 10x + 24$

$x^2 - 10x + 24 = 0$

$(x-4)(x-6) = 0$

$x = 4, \ x = 6$ are the zeros of f.

Interval	$(-\infty, 4)$	$(4, 6)$	$(6, \infty)$
Test Number	0	5	7
Value of f	24	-1	3
Conclusion	Positive	Negative	Positive

The solution set is $\{x \mid x \le 4 \text{ or } x \ge 6\}$ or, using interval notation, $(-\infty, 4] \cup [6, \infty)$.

13. $f(x) = 0$

$2x^2 + 4x + 5 = 0$

$a = 2, b = 4, c = 5$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \pm \sqrt{4^2 - 4(2)(5)}}{2(2)}$$

$$= \frac{-4 \pm \sqrt{-24}}{4} = \frac{-4 \pm 2\sqrt{6}\,i}{4} = -1 \pm \frac{\sqrt{6}}{2}i$$

The complex zeros of f are $-1 - \frac{\sqrt{6}}{2}i$ and $-1 + \frac{\sqrt{6}}{2}i$.

14. $|3x + 1| = 8$

$3x + 1 = 8 \quad \text{or} \quad 3x + 1 = -8$

$3x = 7 \quad \text{or} \quad 3x = -9$

$x = \frac{7}{3} \quad \text{or} \quad x = -3$

The solution set is $\left\{-3, \frac{7}{3}\right\}$.

15. $\left|\frac{x+3}{4}\right| < 2$

$-2 < \frac{x+3}{4} < 2$

$-8 < x + 3 < 8$

$-11 < x < 5$

$\{x \mid -11 < x < 5\}$ or $(-11, 5)$



16. $|2x + 3| - 4 \ge 3$


$|2x + 3| \ge 7$

$2x + 3 \le -7 \quad \text{or} \quad 2x + 3 \ge 7$

$2x \le -10 \quad \text{or} \quad 2x \ge 4$

$x \le -5 \quad \text{or} \quad x \ge 2$

$\{x \mid x \le -5 \text{ or } x \ge 2\}$ or $(-\infty, -5] \cup [2, \infty)$



17. a. $C(m) = 0.15m + 129.50$

b. $C(860) = 0.15(860) + 129.50$

$= 129 + 129.50 = 258.50$

If 860 miles are driven, the rental cost is \$258.50.

c. $C(m) = 213.80$

$0.15m + 129.50 = 213.80$

$0.15m = 84.30$

$m = 562$

The rental cost is \$213.80 if 562 miles were driven.

18. a. $R(x) = x\left(-\frac{1}{10}x + 1000\right) = -\frac{1}{10}x^2 + 1000x$

b. $R(400) = -\frac{1}{10}(400)^2 + 1000(400)$

$= -16{,}000 + 400{,}000$

$= \$384{,}000$

c. $x = \frac{-b}{2a} = \frac{-1000}{2\left(-\frac{1}{10}\right)} = \frac{-1000}{\left(-\frac{1}{5}\right)} = 5000$

The maximum revenue is

Copyright © 2019 Pearson Education, Inc.


Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

$$R(5000) = -\frac{1}{10}(5000)^2 + 1000(5000)$$
$$= -250,000 + 5,000,000$$
$$= \$2,500,000$$

Thus, 5000 units maximizes revenue at \$2,500,000.


d. $p = -\frac{1}{10}(5000) + 1000$
$= -500 + 1000$
$= \$500$

19. a. Set A:




The data appear to be linear with a negative slope.

Set B:




The data appear to be quadratic and opens up.

b. Using the LINear REGression program, the linear function of best fit is: $y = -4.234x - 2.362$.



c. Using the QUADratic REGression program, the quadratic function of best fit is: $y = 1.993x^2 + 0.289x + 2.503$.



Chapter 2 Cumulative Review

1. $P = (-1,3); Q = (4,-2)$

Distance between P and Q:

$$d(P,Q) = \sqrt{\left(4-(-1)\right)^2 + (-2-3)^2}$$
$$= \sqrt{(5)^2 + (5)^2}$$
$$= \sqrt{25+25}$$
$$= \sqrt{50} = 5\sqrt{2}$$

Midpoint between P and Q:

$$\left(\frac{-1+4}{2}, \frac{3-2}{2}\right) = \left(\frac{3}{2}, \frac{1}{2}\right) = (1.5, 0.5)$$

2. $y = x^3 - 3x + 1$

a. $(-2,-1)$: $-1 = (-2)^3 - 3(-2) + 1$
$-1 = -8 + 6 + 1$
$-1 = -1$

Yes, $(-2,-1)$ is on the graph.

b. $(2,3)$: $3 = (2)^3 - 3(2) + 1$
$3 = 8 - 6 + 1$
$3 = 3$


Yes, $(2,3)$ is on the graph.

c. $(3,1)$: $1 = (3)^3 - 3(3) + 1$
$1 = -27 - 9 + 1$
$1 \neq -35$

No, $(3,1)$ is not on the graph.

3. $5x + 3 \geq 0$
$5x \geq -3$
$x \geq -\frac{3}{5}$

The solution set is $\left\{x \,\middle|\, x \geq -\frac{3}{5}\right\}$ or $\left[-\frac{3}{5}, +\infty\right)$.




Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

4. (–1,4) and (2,–2) are points on the line.

$$\text{Slope} = \frac{-2-4}{2-(-1)} = \frac{-6}{3} = -2$$


$$y - y_1 = m(x - x_1)$$
$$y - 4 = -2(x - (-1))$$
$$y - 4 = -2(x + 1)$$
$$y - 4 = -2x - 2$$
$$y = -2x + 2$$



5. Perpendicular to $y = 2x + 1$;
Containing (3,5)

$$\text{Slope of perpendicular} = -\frac{1}{2}$$


$$y - y_1 = m(x - x_1)$$
$$y - 5 = -\frac{1}{2}(x - 3)$$
$$y - 5 = -\frac{1}{2}x + \frac{3}{2}$$
$$y = -\frac{1}{2}x + \frac{13}{2}$$



6. $x^2 + y^2 - 4x + 8y - 5 = 0$

$$x^2 - 4x + y^2 + 8y = 5$$
$$(x^2 - 4x + 4) + (y^2 + 8y + 16) = 5 + 4 + 16$$
$$(x-2)^2 + (y+4)^2 = 25$$
$$(x-2)^2 + (y+4)^2 = 5^2$$

Center: (2,–4) Radius = 5



7. Yes, this is a function since each x-value is paired with exactly one y-value.

8. $f(x) = x^2 - 4x + 1$

a. $f(2) = 2^2 - 4(2) + 1 = 4 - 8 + 1 = -3$

b. $f(x) + f(2) = x^2 - 4x + 1 + (-3)$
$= x^2 - 4x - 2$

c. $f(-x) = (-x)^2 - 4(-x) + 1 = x^2 + 4x + 1$

d. $-f(x) = -(x^2 - 4x + 1) = -x^2 + 4x - 1$

e. $f(x+2) = (x+2)^2 - 4(x+2) + 1$
$= x^2 + 4x + 4 - 4x - 8 + 1$
$= x^2 - 3$

f. $\dfrac{f(x+h) - f(x)}{h}$

$$= \frac{(x+h)^2 - 4(x+h) + 1 - (x^2 - 4x + 1)}{h}$$
$$= \frac{x^2 + 2xh + h^2 - 4x - 4h + 1 - x^2 + 4x - 1}{h}$$
$$= \frac{2xh + h^2 - 4h}{h}$$
$$= \frac{h(2x + h - 4)}{h} = 2x + h - 4$$

9. $h(z) = \dfrac{3z-1}{6z-7}$

The denominator cannot be zero:
$6z - 7 \neq 0$
$6z \neq 7$
$z \neq \frac{7}{6}$

Domain: $\left\{ z \,\middle|\, z \neq \frac{7}{6} \right\}$

10. Yes, the graph represents a function since it passes the Vertical Line Test.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

11. $f(x)=\dfrac{x}{x+4}$

a. $f(1)=\dfrac{1}{1+4}=\dfrac{1}{5}\neq\dfrac{1}{4}$, so $\left(1,\dfrac{1}{4}\right)$ is not on the graph of f.

b. $f(-2)=\dfrac{-2}{-2+4}=\dfrac{-2}{2}=-1$, so $(-2,-1)$ is a point on the graph of f.

c. Solve for x:

$$2=\frac{x}{x+4}$$
$$2x+8=x$$
$$x=-8$$

So, $(-8, 2)$ is a point on the graph of f.


12. $f(x)=\dfrac{x^2}{2x+1}$


$$f(-x)=\frac{(-x)^2}{2(-x)+1}=\frac{x^2}{-2x+1}\neq f(x) \text{ or } -f(x)$$

Therefore, f is neither even nor odd.

13. $f(x)=x^3-5x+4$ on the interval $(-4,4)$

Use MAXIMUM and MINIMUM on the graph of $y_1=x^3-5x+4$.





Local maximum is 5.30 and occurs at $x\approx -1.29$;
Local minimum is –3.30 and occurs at $x\approx 1.29$;
f is increasing on $[-4,-1.29]$ or $[1.29,4]$;
f is decreasing on $[-1.29,1.29]$.

14. $f(x)=3x+5;\quad g(x)=2x+1$

a. $f(x)=g(x)$

$$3x+5=2x+1$$
$$3x+5=2x+1$$
$$x=-4$$

b. $f(x)>g(x)$

$$3x+5>2x+1$$
$$3x+5>2x+1$$
$$x>-4$$

The solution set is $\{x|x>-4\}$ or $(-4,\infty)$.

15. **a.** Domain: $\{x\,|\,-4\le x\le 4\}$ or $[-4, 4]$

Range: $\{y\,|\,-1\le y\le 3\}$ or $[-1, 3]$

b. Intercepts: $(-1,0)$, $(0,-1)$, $(1,0)$

x-intercepts: $-1, 1$

y-intercept: -1


c. The graph is symmetric with respect to the y-axis.

d. When $x=2$, the function takes on a value of 1. Therefore, $f(2)=1$.


e. The function takes on the value 3 at $x=-4$ and $x=4$.

f. $f(x)<0$ means that the graph lies below the x-axis. This happens for x values between -1 and 1. Thus, the solution set is $\{x\,|\,-1<x<1\}$ or $(-1,1)$.

g. The graph of $y=f(x)+2$ is the graph of $y=f(x)$ but shifted up 2 units.




h. The graph of $y=f(-x)$ is the graph of $y=f(x)$ but reflected about the y-axis.



i. The graph of $y=2f(x)$ is the graph of $y=f(x)$ but stretched vertically by a factor of 2. That is, the coordinate of each point is multiplied by 2.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual



j. Since the graph is symmetric about the y-axis, the function is even.

k. The function is increasing on the open interval $(0,4)$.


Chapter 2 Projects

Project I – Internet-based Project


Answers will vary.

Project II


a.



b. The data would be best fit by a quadratic function.



$y = 0.085x^2 - 14.46x + 1069.52$



These results seem reasonable since the function fits the data well.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

c. $s_0 = 0$m

Type	Weight kg	Velocity m/sec	Equation in the form: $s(t) = -4.9t^2 + \frac{\sqrt{2}}{2}v_0t + s_0$
MG 17	10.2	905	$s(t) = -4.9t^2 + 639.93t$ Best. (It goes the highest)
MG 131	19.7	710	$s(t) = -4.9t^2 + 502.05t$
MG 151	41.5	850	$s(t) = -4.9t^2 + 601.04t$
MG 151/20	42.3	695	$s(t) = -4.9t^2 + 491.44t$
MG/FF	35.7	575	$s(t) = -4.9t^2 + 406.59t$
MK 103	145	860	$s(t) = -4.9t^2 + 608.11t$
MK 108	58	520	$s(t) = -4.9t^2 + 367.70t$
WGr 21	111	315	$s(t) = -4.9t^2 + 222.74t$

$s_0 = 200$m

Type	Weight kg	Velocity m/sec	Equation in the form: $s(t) = -4.9t^2 + \frac{\sqrt{2}}{2}v_0t + s_0$
MG 17	10.2	905	$s(t) = -4.9t^2 + 639.93t + 200$ Best. (It goes the highest)
MG 131	19.7	710	$s(t) = -4.9t^2 + 502.05t + 200$
MG 151	41.5	850	$s(t) = -4.9t^2 + 601.04t + 200$
MG 151/20	42.3	695	$s(t) = -4.9t^2 + 491.44t + 200$
MG/FF	35.7	575	$s(t) = -4.9t^2 + 406.59t + 200$
MK 103	145	860	$s(t) = -4.9t^2 + 608.11t + 200$
MK 108	58	520	$s(t) = -4.9t^2 + 367.70t + 200$
WGr 21	111	315	$s(t) = -4.9t^2 + 222.74t + 200$

$s_0 = 30$m

Type	Weight kg	Velocity m/sec	Equation in the form: $s(t) = -4.9t^2 + \frac{\sqrt{2}}{2}v_0t + s_0$
MG 17	10.2	905	$s(t) = -4.9t^2 + 639.93t + 30$ Best. (It goes the highest)
MG 131	19.7	710	$s(t) = -4.9t^2 + 502.05t + 30$
MG 151	41.5	850	$s(t) = -4.9t^2 + 601.04t + 30$
MG 151/20	42.3	695	$s(t) = -4.9t^2 + 491.44t + 30$
MG/FF	35.7	575	$s(t) = -4.9t^2 + 406.59t + 30$
MK 103	145	860	$s(t) = -4.9t^2 + 608.11t + 30$
MK 108	58	520	$s(t) = -4.9t^2 + 367.70t + 30$
WGr 21	111	315	$s(t) = -4.9t^2 + 222.74t + 30$

Notice that the gun is what makes the difference, not how high it is mounted necessarily. The only way to change the true maximum height that the projectile can go is to change the angle at which it fires.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Project III

a.

x	1	2	3	4	5
$y=-2x+5$	3	1	−1	−3	−5

b. $\dfrac{\Delta y}{\Delta x}=\dfrac{y_2-y_1}{x_2-x_1}=\dfrac{1-3}{1}=-2$

$\dfrac{\Delta y}{\Delta x}=\dfrac{y_2-y_1}{x_2-x_1}=\dfrac{-1-1}{1}=-2$

$\dfrac{\Delta y}{\Delta x}=\dfrac{y_2-y_1}{x_2-x_1}=\dfrac{-3-(-1)}{1}=-2$

$\dfrac{\Delta y}{\Delta x}=\dfrac{y_2-y_1}{x_2-x_1}=\dfrac{-5-(-3)}{1}=-2$

All of the values of $\dfrac{\Delta y}{\Delta x}$ are the same.

c.



d. $\dfrac{\Delta I}{\Delta x}=\dfrac{30633-9548}{10}=2108.50$

$\dfrac{\Delta I}{\Delta x}=\dfrac{37088-30633}{10}=645.50$

$\dfrac{\Delta I}{\Delta x}=\dfrac{41072-37088}{10}=398.40$

$\dfrac{\Delta I}{\Delta x}=\dfrac{34414-41072}{10}=-665.80$

$\dfrac{\Delta I}{\Delta x}=\dfrac{19167-34414}{10}=-1524.70$

These $\dfrac{\Delta I}{\Delta x}$ values are not all equal. The data are not linearly related.

e.

x	−2	−1	0	1	2	3	4
y	23	9	3	5	15	33	59
$\dfrac{\Delta y}{\Delta x}$		−14	−6	2	10	18	26

As x increases, $\dfrac{\Delta y}{\Delta x}$ increases. This makes sense because the parabola is increasing (going up) steeply as x increases.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

f.

x	−2	−1	0	1	2	3	4
y	23	9	3	5	15	33	59
$\frac{\Delta^2 y}{\Delta x^2}$			8	8	8	8	8

The second differences are all the same.

g. The paragraph should mention at least two observations:

1. The first differences for a linear function are all the same.
2. The second differences for a quadratic function are the same.

Project IV

a. – i. Answers will vary , depending on where the CBL is located above the bouncing ball.

j. The ratio of the heights between bounces will be the same.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Mini-Lecture 2.1
Properties of Linear Functions and Linear Models


Learning Objectives:

1. Graph Linear Functions
2. Use Average Rate of Change to Identify Linear Functions
3. Determine Whether a Linear function Is Increasing, Decreasing, or Constant
4. Find the Zero of a Linear Function
4. Build Linear Models from Verbal Descriptions

Examples:

1. Suppose that $f(x)=5x-9$ and $g(x)=-3x+7$. Solve $f(x)=g(x)$. Then graph $y=f(x)$ and $y=g(x)$ and label the point that represents the solution to the equation $f(x)=g(x)$.

2. In parts a) and b) using the following figure,



a) Solve $f(x)=12$. b) Solve $0<f(x)<12$.

3. The monthly cost C, in dollars, for renting a full-size car for a day from a particular agency is modeled by the function $C(x)=0.12x+40$, where x is the number of miles driven. Suppose that your budget for renting a car is $\$100$. What is the maximum number of miles that you can drive in one day?

4. Find a firm's break-even point if $R(x)=10x$ and $C(x)=7x+6000$.

Copyright © 2019 Pearson Education, Inc.


Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Teaching Notes:

- Review the slope-intercept form of the equation of a line.
- Emphasize the theorem for Average Rate of Change of a Linear Function in the book.
- Emphasize the need to express the answer to a verbal problem in terms of the units given in the problem.
- Discuss depreciation, supply and demand, and break-even analysis in more depth before working either the examples in the book or the examples above.

Answers:

1. $x = 2$,



2. a) $x = 6$, b) $2 < x < 6$

3. 500 miles

4. 2000 units

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Mini-Lecture 2.2
Building Linear Models from Data

Learning Objectives:

1. Draw and Interpret Scatter Diagrams
2. Distinguish between Linear and Nonlinear Relations
3. Use a Graphing Utility to Find the Line of Best Fit

Examples:

x	3	7	8	9	11	15
y	2	4	7	8	6	10

1. Draw a scatter diagram. Select two points from the scatter diagram and find the equation of the line containing the two points.

2. Graph the line on the scatter diagram.

The marketing manager for a toy company wishes to find a function that relates the demand D for a doll and p the price of the doll. The following data were obtained based on a price history of the doll. The demand is given in thousands of dolls sold per day.

Price	9.00	10.50	11.00	12.00	12.50	13
Demand	12	11	9	10	9.5	8

3. Use a graphing utility to draw a scatter diagram. Then, find and draw the line of best fit.

4. How many dolls will be demanded if the price is $11.50?

Teaching Notes:


- Many students have trouble deciding what scale to use on the x- and y-axes of their scatter plots. Remind them that the scale does not have to be the same on both axes and that the axes may show a break between zero and the first labeled tick mark.
- Use a set of data and ask different students to pick two points and find the equation of the line through the points. Use the graphing utility to draw a scatter diagram and plot each student's line on the scatter diagram. Then find the line of best fit using the graphing utility and graph it on the scatter diagram. This exercise will give students a better understanding of the line of best fit.
- To help students understand of the correlation coefficient show scatter diagrams for different data sets. Show the students data sets with correlations coefficients close to 1, -1, and 0.

Copyright © 2019 Pearson Education, Inc.


Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Answers: 1. $y = \frac{1}{2}x + \frac{1}{2}$ for points $(3,2)$ and $(7,4)$ 4. $D = 9.77$ thousand dolls


1.



2.



3.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Mini-Lecture 2.3
Quadratic Functions and Their Zeros

Learning Objectives:

1. Find the Zeros of a Quadratic Function by Factoring
2. Find the Zeros of a Quadratic Function Using the Square Root Method
3. Find the Zeros of a Quadratic Function by Completing the Square
4. Find the Zeros of a Quadratic Function Using the Quadratic Formula
5. Find the Point of Intersection of Two Functions
6. Solve Equations That are Quadratic in Form

Examples:

1. Find the zeros by factoring: $f(x) = 3x^2 + 4x - 4$

2. Find the zeros by using the square root method: $f(x) = (4x-1)^2 - 16$

3. Find the zeros by completing the square: $f(x) = x^2 + 4x - 10$

4. Find the zeros by using the quadratic formula: $f(x) = 3x^2 - 5x - 7$

5. Find the real zeros of: $f(x) = x^4 - 11x^2 + 18$

6. Find the points of intersection: $f(x) = x^2 - 4$ & $g(x) = 3 - x^2$

Teaching Notes:

- Students that do not have good skills will struggle with this section. Most students can factor pretty well, but they will commit many types of algebraic mistakes when using the other methods.
- When students use the quadratic formula, they will have trouble simplifying the rational expression. For example, reducing like this $\frac{10 \pm 5\sqrt{10}}{10} = 1 \pm 5\sqrt{10}$ is a common error.
- Completing the square will shine a light on the difficulties that students have with fractions.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Answers:

1. $\frac{2}{3}; -2$

2. $\frac{5}{4}; -\frac{3}{4}$

3. $-2 \pm \sqrt{14}$

4. $\frac{5 \pm \sqrt{109}}{6}$

5. $x = \pm\sqrt{2}; \pm 3$

6. $\left(\pm \frac{\sqrt{14}}{2}, -\frac{1}{2} \right)$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Mini-Lecture 2.4
Properties of Quadratic Functions

Learning Objectives:

1. Graph a Quadratic Function Using Transformations
2. Identify the Vertex and Axis of Symmetry of a Quadratic Function
3. Graph a Quadratic Function Using Its Vertex, Axis, and Intercepts
4. Find a Quadratic Function Given Its Vertex and One Other Point
5. Find the Maximum and Minimum Value of a Quadratic Function

Examples:


1. Find the coordinates of the vertex for the parabola defined by the given quadratic function. $f(x) = -3x^2 + 5x - 4$

2. Sketch the graph of the quadratic function by determining whether it opens up or down and by finding its vertex, axis of symmetry, y-intercepts, and x-intercepts, if any. $f(x) = 6 - 5x + x^2$

3. For the quadratic function, $f(x) = 4x^2 - 8x$,

   a) determine, without graphing, whether the function has a minimum value or a maximum value,
   b) find the minimum or maximum value.

4. Determine the quadratic function whose graph is given.



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


Teaching Notes:

- Remind students to review transformations of graphs before beginning to graph quadratic functions.
- Emphasize from the book "Steps for Graphing a Quadratic Function $f(x) = ax^2 + bx + c$, $a \neq 0$."
- Stress the use of a from the standard form to determine the direction the parabola is opening before beginning to graph it. Students need to recognize early on the benefits of knowing as much about a graph as possible before beginning to draw it.
- In addition to the intercepts, encourage students to use symmetry to find additional points on the graph of a quadratic function.
- Many students will give the x-value found with $x = -\frac{b}{2a}$ as the maximum or minimum value of the quadratic function. Emphasize that finding the maximum or minimum is a two-step process. First, find <u>where</u> it occurs (x), then find <u>what</u> it is (y).

Answers:

1. $\left(\frac{5}{6}, -\frac{23}{12}\right)$

2. opens up, vertex $\left(\frac{5}{2}, -\frac{1}{4}\right)$, axis $x = \frac{5}{2}$, x-intercepts 2 and 3, y-intercept 6



3. a) minimum b) minimum of -4

4. $f(x) = -x^2 + 4x - 8$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual


Mini-Lecture 2.5
Inequalities Involving Quadratic Functions

Learning Objectives:

1. Solve Inequalities Involving a Quadratic Function

Examples:

1. Use the figure to solve the inequality $f(x) \geq g(x)$.



2. Solve and express the solution in interval notation.

$$9x^2 - 6x + 1 < 0$$

3. Solve the inequality.

$$x(x+2) > 15$$

4. Solve.

$$f(x) > g(x). \quad f(x) = -x^2 + 2x + 3;\ g(x) = -x^2 - 2x + 8$$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Teaching Notes:

- Suggest that students review interval notation. Caution them to check for the correct use of brackets or parentheses in solutions written in interval notation.
- Suggest that students review factoring a trinomial.
- Be sure to show Option I and Option II in Example 2 in the book.

Answers:

1. $[-2,1]$

2. $\varnothing$

3. $(-\infty,-5)\cup(3,\infty)$

4. $\left(\frac{5}{4},\infty\right)$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Mini-Lecture 2.6

Building Quadratic Models from Verbal Descriptions and from Data

Learning Objectives:

1. Build Quadratic Models from Verbal Descriptions
2. Build Quadratic Models from Data

Examples:

1. Among all pairs of numbers whose sum is 50, find a pair whose product is as large as possible. What is the maximum product?

2. A person standing close to the edge of the top of a 180-foot tower throws a ball vertically upward. The quadratic function $s(t) = -16t^2 + 64t + 180$ models the ball's height above ground , $s(t)$, in feet, t seconds after it was thrown. After how many seconds does the ball reach its maximum height? What is the maximum height?

3. The price p (in dollars) and the quantity x sold of a certain product obey the demand equation $p = -\frac{1}{4}x + 120$. Find the model that expresses the revenue R as a function of x. What quantity x maximizes revenue? What is the maximum revenue?

4. The following data represent the percentage of the population in a certain country aged 40 or older whose age is x who do not have a college degree of some type.

Age, x	40	45	50	55	60	65
No college	25.4	23.2	21.8	24.5	26.1	29.8

Find a quadratic model that describes the relationship between age and percentage of the population that do not have a college degree. Use the model to predict the percentage of 53-year-olds that do not have a college degree.

Teaching Notes:

- Show students how to use MAXIMUM and MINIMUM on the graphing utility.
- Show students how to use the QUADratic REGression program on the graphing utility.
- Encourage students to review the discriminant.

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Answers:

1. $(25, 25)$, 625

2. 2 sec., 244 ft

3. $R(x) = -\frac{1}{4}x^2 + 120x$, $x = 240$, $R(240) = \$14,400$

4. $P(x) = .0296x^2 - 2.9216x + 94.6550$, 23.0%

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Mini-Lecture 2.7
Complex Zeros of a Quadratic Function

Learning Objectives:

1. Find the Complex Zeros of a Quadratic Function

Examples:

1. Find the complex zeros; graph the function; label the intercepts.

   a) $f(x) = x^2 + 5$ b) $f(x) = x^2 + 2x + 7$

   c) $f(x) = 2x^2 - 4x - 5$ d) $f(x) = x^2 - 2x + 5$

Teaching Notes:

- Have students review the quadratic formula.
- Students will need to know how to reduce radical and rational expressions.
- Remind the students that i is outside the radical when simplifying a radical expression.

Answers:


1. a) Complex zeros $x = \pm i\sqrt{5}$, y-intercept $= 5$, no x-intercepts.

   b) Complex zeros $x = -1 \pm i\sqrt{6}$, y-intercept $= 7$, no x-intercepts


   c) Complex zeros $x = \dfrac{2 \pm \sqrt{14}}{2}$, y-intercept $= -5$, x-intercepts $x = \dfrac{2 \pm \sqrt{14}}{2}$

   d) Complex zeros $x = 1 \pm 2i$, y-intercept $= 5$, no x-intercepts


a)




b)



c)



d)



Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

Mini-Lecture 2.8

Equations and Inequalities Involving the Absolute Value Function

Learning Objectives:

1. Solve Absolute Value Equations
2. Solve Absolute Value Inequalities

Examples:

1. Solve each equation.

a) $|5x-10|=15$ b) $\left|\frac{2}{3}x+6\right|=12$ c) $|4-3x|-4=1$ d) $|3-x|=-7$

2. Solve each absolute value inequality.

a) $|3x|\le 21$ b) $|4x-3|\ge 9$ c) $|2-6x|-5<1$

Teaching Notes:

- When solving absolute value equations, students will sometimes forget that there are two solutions.
- Students will often not isolate the absolute value expression before trying to solve, such as examples 1 c) and 2 c) above.
- Some students try to combine two intervals that cannot be combined, such as $-3<x>2$.

Answers:

1. a) $x=5, x=-1$ b) $x=9, x=-27$ c) $x=-\frac{1}{3}, x=3$ d) No solution

2. a) $-7\le x\le 7$ b) $x\le -\frac{3}{2}$ *or* $x\ge 3$ c) $-\frac{2}{3}<x<\frac{4}{3}$

Copyright © 2019 Pearson Education, Inc.

Precalculus Concepts Through Functions A Right Triangle Approach to Trigonometry 4th edition Sullivan Solutions Manual

4. Cannons The velocity of a projectile depends upon many factors, in particular, the weight of the ammunition.

(a) Plot a scatter diagram of the data in the table below. Let x be the weight in kilograms and let y be the velocity in meters per second.

Type	Weight (kg)	Initial Velocity (m/sec)
MG 17	10.2	905
MG 131	19.7	710
MG 151	41.5	850
MG 151/20	42.3	695
MG/FF	35.7	575
MK 103	145	860
MK 108	58	520
WGr 21	111	315

(Data and information taken from "Flugzeug-Handbuch, Ausgabe Dezember 1996: Guns and Cannons of the Jagdwaffe" at www.xs4all.nl/~rhorta/jgguns.htm)

Precalculus Concepts Through Functions A Right

(b) Determine which type of function would fit this data the best: linear or quadratic. Use a graphing utility to find the function of best fit. Are the results reasonable?

(c) Based on velocity, we can determine how high a projectile will travel before it begins to come back down. If a cannon is fired at an angle of 45° to the horizontal, then the function for the height of the projectile is given by $s(t) = -16t^2 + \frac{\sqrt{2}}{2}v_0 t + s_0$, where v_0 is the velocity at which the shell leaves the cannon (initial velocity), and s_0 is the initial height of the nose of the cannon (because cannons are not very long, we may assume that the nose and the firing pin at the back are at the same height for simplicity). Graph the function $s = s(t)$ for each of the guns described in the table. Which gun would be the best for anti-aircraft if the gun were sitting on the ground? Which would be the best to have mounted on a hilltop or on the top of a tall building? If the guns were on the turret of a ship, which would be the most effective?

Precalculus Concepts Through Functions A Right

7. **CBL Experiment** Locate the motion detector on a Calculator Based Laboratory (CBL) or a Calculator Based Ranger (CBR) above a bouncing ball.

(a) Plot the data collected in a scatter diagram with time as the independent variable.

(b) Find the quadratic function of best fit for the second bounce.

(c) Find the quadratic function of best fit for the third bounce.

(d) Find the quadratic function of best fit for the fourth bounce.

(e) Compute the maximum height for the second bounce.

(f) Compute the maximum height for the third bounce.

(g) Compute the maximum height for the fourth bounce.

(h) Compute the ratio of the maximum height of the third bounce to the maximum height of the second bounce.

(i) Compute the ratio of the maximum height of the fourth bounce to the maximum height of the third bounce.



(j) Compare the results from parts (h) and (i). What do you conclude?

Precalculus Concepts Through Functions A Right Tr

3. Suppose $f(x) = \sin x$.

(a) Build a table of values for $f(x)$ where $x = 0, \frac{\pi}{6}, \frac{\pi}{4}, \frac{\pi}{3}, \frac{\pi}{2}, \frac{2\pi}{3}, \frac{3\pi}{4}, \frac{5\pi}{6}, \pi, \frac{7\pi}{6}, \frac{5\pi}{4}, \frac{4\pi}{3}, \frac{3\pi}{2}, \frac{5\pi}{3}, \frac{7\pi}{4}, \frac{11\pi}{6}, 2\pi$. Use exact values.

(b) Find the **first differences** for each consecutive pair of values in part (a). That is, evaluate $g(x_i) = \frac{\Delta f(x_i)}{\Delta x_i} = \frac{f(x_{i+1}) - f(x_i)}{x_{i+1} - x_i}$, where $x_1 = 0$, $x_2 = \frac{\pi}{6}, \ldots, x_{17} = 2\pi$. Use your calculator to approximate each value rounded to three decimal places.

(c) Plot the points $(x_i, g(x_i))$ for $i = 1, \ldots, 16$ on a scatter diagram. What shape does the set of points give? What function does this resemble? Fit a sine curve of best fit to the points. How does that relate to your guess?

(d) Find the first differences for each consecutive pair of values in part (b). That is, evaluate $h(x_i) = \frac{\Delta g(x_i)}{\Delta x_i} = \frac{g(x_{i+1}) - g(x_i)}{x_{i+1} - x_i}$ where $x_1 = 0$, $x_2 = \frac{\pi}{6}, \ldots, x_{16} = \frac{11\pi}{6}$. This is the set of **second differences** of $f(x)$. Use your calculator to approximate each value rounded to three decimal places. Plot the points $(x_i, h(x_i))$ for $i = 1, \ldots, 15$ on a scatter diagram. What shape does the set of points give? What function does this resemble? Fit a sine curve of best fit to the points. How does that relate to your guess?

(e) Find the first differences for each consecutive pair of values in part (d). That is, evaluate $k(x_i) = \dfrac{\Delta h(x_i)}{\Delta x_i} = \dfrac{h(x_{i+1}) - h(x_i)}{x_{i+1} - x_i}$, where $x_1 = 0$, $x_2 = \dfrac{\pi}{6}, \ldots, x_{15} = \dfrac{7\pi}{4}$. This is the set of **third differences** of $f(x)$. Use your calculator to approximate each value rounded to three decimal places. Plot the points $(x_i, k(x_i))$ for $i = 1, \ldots, 14$ on a scatter diagram. What shape does the set of points give? What function does this resemble? Fit a sine curve of best fit to the points. How does that relate to your guess?

(f) Find the first differences for each consecutive pair of values in part (e). That is, evaluate $m(x_i) = \dfrac{\Delta k(x_i)}{\Delta x_i} = \dfrac{k(x_{i+1}) - k(x_i)}{x_{i+1} - x_i}$, where $x_1 = 0$, $x_2 = \dfrac{\pi}{6}, \ldots, x_{14} = \dfrac{5\pi}{3}$. This is the set of **fourth differences** of $f(x)$.

Use your calculator to approximate each value rounded to three decimal places. Plot the points $(x_i, m(x_i))$ for $i = 1, \ldots, 13$ on a scatter diagram. What shape does the set of points give? What function does this resemble? Fit a sine curve of best fit to the points. How does that relate to your guess?

(g) What pattern do you notice about the curves that you found? What happened in part (f)? Can you make a generalization about what happened as you computed the differences? Explain your answers.

Precalculus Concepts Through Functions A Right