Calculus 3rd edition Briggs Test Bank

SKU: 9780134765631-TEST-BANK

Exam

Name_____

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find the average velocity of an object which follows the position function s(t) over the given interval.

1) $s(t) = t^2 + 2t$, $[2, 7]$ 1) _____

A) $\frac{63}{5}$ B) $\frac{55}{7}$ C) 11 D) 9

2) $s(t) = 6t^3 - 8t^2 + 2$, $[-8, -1]$ 2) _____

A) 12 B) 510 C) $-\frac{12}{7}$ D) -3570

3) $s(t) = \sqrt{2t}$, $[2, 8]$ 3) _____

A) 7 B) $-\frac{3}{10}$ C) 2 D) $\frac{1}{3}$

4) $s(t) = \frac{3}{t-2}$, $[4, 7]$ 4) _____

A) $-\frac{3}{10}$ B) 2 C) $\frac{1}{3}$ D) 7

5) $s(t) = 4t^2$, $\left[0, \frac{7}{4}\right]$ 5) _____

A) $-\frac{3}{10}$ B) $\frac{1}{3}$ C) 2 D) 7

6) $s(t) = -3t^2 - t$, $[5, 6]$ 6) _____

A) $\frac{1}{2}$ B) -34 C) $-\frac{1}{6}$ D) -2

7) $s(t) = \sin(4t)$, $\left[0, \frac{\pi}{8}\right]$ 7) _____

A) $-\frac{8}{\pi}$ B) $\frac{\pi}{8}$ C) $\frac{4}{\pi}$ D) $\frac{8}{\pi}$

8) $s(t) = 5 + \tan t$, $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$ 8) _____

A) $-\frac{4}{\pi}$ B) $\frac{4}{\pi}$ C) $-\frac{16}{11}$ D) 0

Calculus 3rd edition Briggs Test Bank

Use the table to find the instantaneous velocity of y at the specified value of x.

9) x = 1. 9) _____

x	y
0	0
0.2	0.02
0.4	0.08
0.6	0.18
0.8	0.32
1.0	0.5
1.2	0.72
1.4	0.98

A) 0.5 B) 1.5 C) 2 D) 1

10) x = 1. 10) _____

x	y
0	0
0.2	0.01
0.4	0.04
0.6	0.09
0.8	0.16
1.0	0.25
1.2	0.36
1.4	0.49

A) 1 B) 0.5 C) 1.5 D) 2

11) x = 1. 11) _____

x	y
0	0
0.2	0.12
0.4	0.48
0.6	1.08
0.8	1.92
1.0	3
1.2	4.32
1.4	5.88

A) 6 B) 8 C) 4 D) 2

Calculus 3rd edition Briggs Test Bank

12) $x = 2$. 12) _____

x	y
0	10
0.5	38
1.0	58
1.5	70
2.0	74
2.5	70
3.0	58
3.5	38
4.0	10

A) –8 B) 0 C) 8 D) 4

13) $x = 1$. 13) _____

x	y
0.900	–0.05263
0.990	–0.00503
0.999	–0.0005
1.000	0.0000
1.001	0.0005
1.010	0.00498
1.100	0.04762

A) 1 B) 0 C) 0.5 D) –0.5

Find the slope of the curve for the given value of x.

14) $y = x^2 + 5x$, $x = 4$ 14) _____

A) slope is $\frac{1}{20}$ B) slope is –39 C) slope is 13 D) slope is $-\frac{4}{25}$

15) $y = x^2 + 11x - 15$, $x = 1$ 15) _____

A) slope is $\frac{1}{20}$ B) slope is –39 C) slope is $-\frac{4}{25}$ D) slope is 13

16) $y = x^3 - 9x$, $x = 1$ 16) _____

A) slope is 1 B) slope is –6 C) slope is –3 D) slope is 3

17) $y = x^3 - 3x^2 + 4$, $x = 3$ 17) _____

A) slope is 0 B) slope is 1 C) slope is –9 D) slope is 9

18) $y = -3 - x^3$, $x = 1$ 18) _____

A) slope is 0 B) slope is 3 C) slope is –1 D) slope is –3

Calculus 3rd edition Briggs Test Bank

Solve the problem.

19) Given $\lim_{x \to 0^-} f(x) = L_l$, $\lim_{x \to 0^+} f(x) = L_r$, and $L_l \neq L_r$, which of the following statements is true? 19) _____

I. $\lim_{x \to 0} f(x) = L_l$

II. $\lim_{x \to 0} f(x) = L_r$

III. $\lim_{x \to 0} f(x)$ does not exist.

A) III B) II C) none D) I

20) Given $\lim_{x \to 0^-} f(x) = L_l$, $\lim_{x \to 0^+} f(x) = L_r$, and $L_l = L_r$, which of the following statements is false? 20) _____

I. $\lim_{x \to 0} f(x) = L_l$

II. $\lim_{x \to 0} f(x) = L_r$

III. $\lim_{x \to 0} f(x)$ does not exist.

A) II B) I C) III D) none

21) If $\lim_{x \to 0} f(x) = L$, which of the following expressions are true? 21) _____

I. $\lim_{x \to 0^-} f(x)$ does not exist.

II. $\lim_{x \to 0^+} f(x)$ does not exist.

III. $\lim_{x \to 0^-} f(x) = L$

IV. $\lim_{x \to 0^+} f(x) = L$

A) II and III only B) I and II only C) I and IV only D) III and IV only

22) What conditions, when present, are sufficient to conclude that a function f(x) has a limit as x approaches some value of a? 22) _____

A) Either the limit of f(x) as x→a from the left exists or the limit of f(x) as x→a from the right exists

B) f(a) exists, the limit of f(x) as x→a from the left exists, and the limit of f(x) as x→a from the right exists.


C) The limit of f(x) as x→a from the left exists, the limit of f(x) as x→a from the right exists, and at least one of these limits is the same as f(a).

D) The limit of f(x) as x→a from the left exists, the limit of f(x) as x→a from the right exists, and these two limits are the same.

Calculus 3rd edition Briggs Test Bank


Use the graph to evaluate the limit.

23) $\lim_{x \to -1} f(x)$ 23) _____



A) ∞ B) –1 C) $\frac{1}{2}$ D) $-\frac{1}{2}$


24) $\lim_{x \to 0} f(x)$ 24) _____



A) –3 B) 0 C) 3 D) does not exist


Calculus 3rd edition Briggs Test Bank

25) $\lim_{x \to 0} f(x)$ 25) _____



A) does not exist B) 0 C) 2 D) –2


26) $\lim_{x \to 0} f(x)$ 26) _____



A) –1 B) 0 C) does not exist D) 6


Calculus 3rd edition Briggs Test Bank

27) $\lim_{x \to 0} f(x)$ 27) _____



A) 1 B) ∞ C) –1 D) does not exist


28) $\lim_{x \to 0} f(x)$ 28) _____



A) ∞ B) 1 C) –1 D) does not exist


Calculus 3rd edition Briggs Test Bank

29) $\lim_{x \to 0} f(x)$ 29) _____



A) does not exist B) –2 C) 2 D) 0


30) $\lim_{x \to 0} f(x)$ 30) _____



A) 1 B) does not exist C) 0 D) –2


Calculus 3rd edition Briggs Test Bank

31) $\lim_{x \to 0} f(x)$ 31) _____



A) 2 B) –1 C) –2 D) does not exist

32) Find $\lim_{x \to (-1)^-} f(x)$ and $\lim_{x \to (-1)^+} f(x)$ 32) _____



A) –7; –2 B) –5; –2 C) –2; –7 D) –7; –5

Calculus 3rd edition Briggs Test Bank

Use the table of values of f to estimate the limit.

33) Let $f(x) = x^2 + 8x - 2$, find $\lim_{x \to -2} f(x)$. 33) _____

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)						

A)

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)	5.043	5.364	5.396	5.404	5.436	5.763

; limit = ∞

B)

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)	16.810	17.880	17.988	18.012	18.120	19.210

; limit = 18.0

C)

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)	16.692	17.592	17.689	17.710	17.808	18.789

; limit = 17.70

D)

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)	5.043	5.364	5.396	5.404	5.436	5.763

; limit = 5.40

34) Let $f(x) = \dfrac{x - 4}{\sqrt{x} - 2}$, find $\lim_{x \to 4} f(x)$. 34) _____

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)						

A)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	1.19245	1.19925	1.19993	1.20007	1.20075	1.20745

; limit = ∞

B)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	5.07736	5.09775	5.09978	5.10022	5.10225	5.12236

; limit = 5.10

C)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	3.97484	3.99750	3.99975	4.00025	4.00250	4.02485

; limit = 4.0

D)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	1.19245	1.19925	1.19993	1.20007	1.20075	1.20745

; limit = 1.20

Calculus 3rd edition Briggs Test Bank

35) Let $f(x) = x^2 - 5$, find $\lim_{x \to 0} f(x)$. 35) _____

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)						

A)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−1.4970	−1.4999	−1.5000	−1.5000	−1.4999	−1.4970

; limit = −15.0

B)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−2.9910	−2.9999	−3.0000	−3.0000	−2.9999	−2.9910

; limit = −3.0

C)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−1.4970	−1.4999	−1.5000	−1.5000	−1.4999	−1.4970

; limit = ∞

D)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−4.9900	−4.9999	−5.0000	−5.0000	−4.9999	−4.9900

; limit = −5.0

36) Let $f(x) = \dfrac{x - 1}{x^2 + 2x - 3}$, find $\lim_{x \to 1} f(x)$. 36) _____

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)						

A)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.1564	0.1506	0.1501	0.1499	0.1494	0.1439

; limit = 0.15

B)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	−0.2564	−0.2506	−0.2501	−0.2499	−0.2494	−0.2439

; limit = −0.25

C)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.3564	0.3506	0.3501	0.3499	0.3494	0.3439

; limit = 0.35

D)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.2564	0.2506	0.2501	0.2499	0.2494	0.2439

; limit = 0.25

Calculus 3rd edition Briggs Test Bank

37) Let $f(x) = \dfrac{x^2 + 2x - 3}{x^2 + 4x - 5}$, find $\lim\limits_{x \to 1} f(x)$. 37) _____

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)						

A)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.4872	0.4987	0.4999	0.5001	0.5012	0.5122

; limit = 0.5

B)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.5610	0.5661	0.5666	0.5667	0.5672	0.5721

; limit = 0.5667

C)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.6610	0.6661	0.6666	0.6667	0.6672	0.6721

; limit = 0.6667

D)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.7610	0.7661	0.7666	0.7667	0.7672	0.7721

; limit = 0.7667

38) Let $f(x) = \dfrac{\sin(8x)}{x}$, find $\lim\limits_{x \to 0} f(x)$. 38) _____

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)		7.9914694			7.9914694	

A) limit = 7.5
B) limit = 8
C) limit = 0
D) limit does not exist

39) Let $f(\theta) = \dfrac{\cos(4\theta)}{\theta}$, find $\lim\limits_{\theta \to 0} f(\theta)$. 39) _____

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(θ)	−9.2106099					9.2106099

A) limit = 9.2106099
B) limit does not exist
C) limit = 4
D) limit = 0

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

Provide an appropriate response.

40) It can be shown that the inequalities $1 - \dfrac{x^2}{6} < \dfrac{x \sin(x)}{2 - 2\cos(x)} < 1$ hold for all values of x close to zero. What, if anything, does this tell you about $\dfrac{x \sin(x)}{2 - 2\cos(x)}$? Explain. 40) _____

Calculus 3rd edition Briggs Test Bank

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

41) Write the formal notation for the principle "the limit of a quotient is the quotient of the limits" and include a statement of any restrictions on the principle. 41) _____

A) $\lim_{x \to a} \frac{g(x)}{f(x)} = \frac{g(a)}{f(a)}$.

B) If $\lim_{x \to a} g(x) = M$ and $\lim_{x \to a} f(x) = L$, then $\lim_{x \to a} \frac{g(x)}{f(x)} = \frac{\lim_{x \to a} g(x)}{\lim_{x \to a} f(x)} = \frac{M}{L}$, provided that $f(a) \neq 0$.

C) $\lim_{x \to a} \frac{g(x)}{f(x)} = \frac{g(a)}{f(a)}$, provided that $f(a) \neq 0$.

D) If $\lim_{x \to a} g(x) = M$ and $\lim_{x \to a} f(x) = L$, then $\lim_{x \to a} \frac{g(x)}{f(x)} = \frac{\lim_{x \to a} g(x)}{\lim_{x \to a} f(x)} = \frac{M}{L}$, provided that $L \neq 0$.

42) Provide a short sentence that summarizes the general limit principle given by the formal notation $\lim_{x \to a} [f(x) \pm g(x)] = \lim_{x \to a} f(x) \pm \lim_{x \to a} g(x) = L \pm M$, given that $\lim_{x \to a} f(x) = L$ and $\lim_{x \to a} g(x) = M$. 42) _____

A) The limit of a sum or a difference is the sum or the difference of the functions.
B) The limit of a sum or a difference is the sum or the difference of the limits.
C) The sum or the difference of two functions is the sum of two limits.
D) The sum or the difference of two functions is continuous.

43) The statement "the limit of a constant times a function is the constant times the limit" follows from a combination of two fundamental limit principles. What are they? 43) _____

A) The limit of a product is the product of the limits, and a constant is continuous.
B) The limit of a product is the product of the limits, and the limit of a quotient is the quotient of the limits.
C) The limit of a constant is the constant, and the limit of a product is the product of the limits.
D) The limit of a function is a constant times a limit, and the limit of a constant is the constant.

Find the limit.

44) $\lim_{x \to 15} \sqrt{5}$ 44) _____

A) $\sqrt{15}$ B) 15 C) 5 D) $\sqrt{5}$

45) $\lim_{x \to -5} (4x - 2)$ 45) _____

A) −18 B) −22 C) 22 D) 18

46) $\lim_{x \to -9} (26 - 3x)$ 46) _____

A) −1 B) −53 C) 1 D) 53

Calculus 3rd edition Briggs Test Bank

Give an appropriate answer.

47) Let $\lim_{x \to 2} f(x) = 5$ and $\lim_{x \to 2} g(x) = -7$. Find $\lim_{x \to 2} [f(x) - g(x)]$. 47) _____

A) −2 B) 2 C) 5 D) 12

48) Let $\lim_{x \to -4} f(x) = -6$ and $\lim_{x \to -4} g(x) = 9$. Find $\lim_{x \to -4} [f(x) \cdot g(x)]$. 48) _____

A) −4 B) −54 C) 9 D) 3

49) Let $\lim_{x \to 10} f(x) = 3$ and $\lim_{x \to 10} g(x) = 9$. Find $\lim_{x \to 10} \frac{f(x)}{g(x)}$. 49) _____

A) 10 B) −6 C) 3 D) $\frac{1}{3}$

50) Let $\lim_{x \to 10} f(x) = 196$. Find $\lim_{x \to 10} \sqrt{f(x)}$. 50) _____

A) 10 B) 196 C) 14 D) 3.7417

51) Let $\lim_{x \to 4} f(x) = -4$ and $\lim_{x \to 4} g(x) = 2$. Find $\lim_{x \to 4} [f(x) + g(x)]^2$. 51) _____

A) 20 B) −2 C) 4 D) −6

52) Let $\lim_{x \to 6} f(x) = 8$. Find $\lim_{x \to 6} \sqrt[3]{f(x)}$. 52) _____

A) 6 B) 8 C) 2 D) 3

53) Let $\lim_{x \to -8} f(x) = 5$ and $\lim_{x \to -8} g(x) = -9$. Find $\lim_{x \to -8} \left[\frac{-8f(x) - 2g(x)}{1 + g(x)}\right]$. 53) _____

A) − 42 B) $\frac{11}{4}$ C) $\frac{29}{4}$ D) −8

Find the limit.

54) $\lim_{x \to 2} (x^3 + 5x^2 - 7x + 1)$ 54) _____

A) 15 B) does not exist C) 0 D) 29

55) $\lim_{x \to 2} (3x^5 - 2x^4 + 4x^3 + x^2 - 5)$ 55) _____

A) 47 B) 159 C) 95 D) 31

56) $\lim_{x \to -1} \frac{x}{3x + 2}$ 56) _____

A) $-\frac{1}{5}$ B) does not exist C) 1 D) 0

Calculus 3rd edition Briggs Test Bank

57) $\lim_{x \to 0} \frac{x^3 - 6x + 8}{x - 2}$ 57) _____

A) Does not exist B) 4 C) −4 D) 0

58) $\lim_{x \to 1} \frac{3x^2 + 7x - 2}{3x^2 - 4x - 2}$ 58) _____

A) Does not exist B) $-\frac{7}{4}$ C) 0 D) $-\frac{8}{3}$

59) $\lim_{x \to 2} (x + 1)^2(x - 1)^3$ 59) _____

A) 9 B) 27 C) 1 D) 243

60) $\lim_{x \to 4} \sqrt{x^2 + 4x + 4}$ 60) _____

A) does not exist B) 36 C) ±6 D) 6

61) $\lim_{x \to 5} \sqrt{7x + 91}$ 61) _____

A) 126 B) $-3\sqrt{14}$ C) −126 D) $3\sqrt{14}$

62) $\lim_{h \to 0} \frac{2}{\sqrt{3h + 4} + 2}$ 62) _____

A) 2 B) 1 C) 1/2 D) Does not exist

63) $\lim_{x \to 0} \frac{\sqrt{1 + x} - 1}{x}$ 63) _____

A) 0 B) 1/2 C) Does not exist D) 1/4

Determine the limit by sketching an appropriate graph.

64) $\lim_{x \to 3^-} f(x)$, where $f(x) = \begin{cases} -5x + 0 & \text{for } x < 3 \\ 3x + 1 & \text{for } x \geq 3 \end{cases}$ 64) _____

A) 1 B) −15 C) 10 D) 2

65) $\lim_{x \to 4^+} f(x)$, where $f(x) = \begin{cases} -2x + 1 & \text{for } x < 4 \\ 4x + 2 & \text{for } x \geq 4 \end{cases}$ 65) _____

A) −7 B) 18 C) 2 D) 3

66) $\lim_{x \to -4^+} f(x)$, where $f(x) = \begin{cases} x^2 + 3 & \text{for } x \neq -4 \\ 0 & \text{for } x = -4 \end{cases}$ 66) _____

A) 13 B) 19 C) 0 D) 16

Calculus 3rd edition Briggs Test Bank

67) $\lim_{x \to 5^-} f(x)$, where $f(x) = \begin{cases} \sqrt{16 - x^2} & 0 \le x < 4 \\ 4 & 4 \le x < 5 \\ 5 & x = 5 \end{cases}$ 67) _____

A) 5 B) Does not exist C) 4 D) 0

68) $\lim_{x \to -6^+} f(x)$, where $f(x) = \begin{cases} 2x & -6 \le x < 0, \text{ or } 0 < x \le 2 \\ 2 & x = 0 \\ 0 & x < -6 \text{ or } x > 2 \end{cases}$ 68) _____

A) 4 B) Does not exist C) –12 D) –0

Find the limit, if it exists.

69) $\lim_{x \to 0} \dfrac{x^3 + 12x^2 - 5x}{5x}$ 69) _____

A) 0 B) 5 C) –1 D) Does not exist

70) $\lim_{x \to 1} \dfrac{x^4 - 1}{x - 1}$ 70) _____

A) 0 B) 2 C) Does not exist D) 4

71) $\lim_{x \to 2} \dfrac{x^2 - 4}{x - 2}$ 71) _____

A) 1 B) Does not exist C) 4 D) 2

72) $\lim_{x \to -3} \dfrac{x^2 + 13x + 30}{x + 3}$ 72) _____

A) 78 B) 7 C) 13 D) Does not exist

73) $\lim_{x \to 4} \dfrac{x^2 + 5x - 36}{x - 4}$ 73) _____

A) 5 B) Does not exist C) 0 D) 13

74) $\lim_{x \to 6} \dfrac{x^2 + 4x - 60}{x^2 - 36}$ 74) _____

A) 0 B) $-\dfrac{1}{3}$ C) $\dfrac{4}{3}$ D) Does not exist

75) $\lim_{x \to 2} \dfrac{x^2 - 4}{x^2 - 3x + 2}$ 75) _____

A) Does not exist B) 2 C) 4 D) 0

76) $\lim_{x \to -6} \dfrac{x^2 + 5x - 6}{x^2 + 5x - 6}$ 76) _____

A) $\dfrac{5}{7}$ B) 1 C) Does not exist D) – 1

Calculus 3rd edition Briggs Test Bank

77) $\lim_{h \to 0} \frac{(x+h)^3 - x^3}{h}$ 77) _____

A) Does not exist B) $3x^2$ C) $3x^2 + 3xh + h^2$ D) 0

78) $\lim_{x \to 7} \frac{|7 - x|}{7 - x}$ 78) _____

A) 0 B) –1 C) Does not exist D) 1

Provide an appropriate response.

79) It can be shown that the inequalities $-x \le x \cos\left(\frac{1}{x}\right) \le x$ hold for all values of $x \ge 0$. 79) _____

Find $\lim_{x \to 0} x \cos\left(\frac{1}{x}\right)$ if it exists.

A) does not exist B) 0.0007 C) 1 D) 0

80) The inequality $1 - \frac{x^2}{2} < \frac{\sin x}{x} < 1$ holds when x is measured in radians and $|x| < 1$. 80) _____

Find $\lim_{x \to 0} \frac{\sin x}{x}$ if it exists.

A) 0.0007 B) does not exist C) 1 D) 0

81) If $x^3 \le f(x) \le x$ for x in [–1,1], find $\lim_{x \to 0} f(x)$ if it exists. 81) _____

A) 0 B) 1 C) –1 D) does not exist

Compute the values of f(x) and use them to determine the indicated limit.

82) If $f(x) = x^2 + 8x - 2$, find $\lim_{x \to 2} f(x)$. 82) _____

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)						

A)

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)	16.692	17.592	17.689	17.710	17.808	18.789

; limit = 17.70

B)

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)	5.043	5.364	5.396	5.404	5.436	5.763

; limit = 5.40

C)

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)	5.043	5.364	5.396	5.404	5.436	5.763

; limit = ∞

D)

x	1.9	1.99	1.999	2.001	2.01	2.1
f(x)	16.810	17.880	17.988	18.012	18.120	19.210

; limit = 18.0

Calculus 3rd edition Briggs Test Bank

83) If $f(x) = \dfrac{x^4 - 1}{x - 1}$, find $\lim\limits_{x \to 1} f(x)$. 83) _____

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)						

A)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	1.032	1.182	1.198	1.201	1.218	1.392

; limit = ∞

B)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	1.032	1.182	1.198	1.201	1.218	1.392

; limit = 1.210

C)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	4.595	5.046	5.095	5.105	5.154	5.677

; limit = 5.10

D)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	3.439	3.940	3.994	4.006	4.060	4.641

; limit = 4.0

84) If $f(x) = \dfrac{x^3 - 6x + 8}{x - 2}$, find $\lim\limits_{x \to 0} f(x)$. 84) _____

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)						

A)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−1.22843	−1.20298	−1.20030	−1.19970	−1.19699	−1.16858

; limit = −1.20

B)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−2.18529	−2.10895	−2.10090	−2.99910	−2.09096	−2.00574

; limit = −2.10

C)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−4.09476	−4.00995	−4.00100	−3.99900	−3.98995	−3.89526

; limit = −4.0

D)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−1.22843	−1.20298	−1.20030	−1.19970	−1.19699	−1.16858

; limit = ∞

Calculus 3rd edition Briggs Test Bank

85) If $f(x) = \dfrac{x - 4}{\sqrt{x} - 2}$, find $\lim\limits_{x \to 4} f(x)$. 85) _____

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)						

A)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	5.07736	5.09775	5.09978	5.10022	5.10225	5.12236

; limit = 5.10

B)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	1.19245	1.19925	1.19993	1.20007	1.20075	1.20745

; limit = ∞

C)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	3.97484	3.99750	3.99975	4.00025	4.00250	4.02485

; limit = 4.0

D)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	1.19245	1.19925	1.19993	1.20007	1.20075	1.20745

; limit = 1.20

86) If $f(x) = x^2 - 5$, find $\lim\limits_{x \to 0} f(x)$. 86) _____

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)						

A)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−2.9910	−2.9999	−3.0000	−3.0000	−2.9999	−2.9910

; limit = −3.0

B)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−4.9900	−4.9999	−5.0000	−5.0000	−4.9999	−4.9900

; limit = −5.0

C)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−1.4970	−1.4999	−1.5000	−1.5000	−1.4999	−1.4970

; limit = −15.0

D)

x	−0.1	−0.01	−0.001	0.001	0.01	0.1
f(x)	−1.4970	−1.4999	−1.5000	−1.5000	−1.4999	−1.4970

; limit = ∞

Calculus 3rd edition Briggs Test Bank

87) If $f(x) = \dfrac{\sqrt{x+1}}{x+1}$, find $\lim\limits_{x \to 1} f(x)$. 87) _____

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)						

A)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	2.15293	2.13799	2.13656	2.13624	2.13481	2.12106

; limit = 2.13640

B)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.21764	0.21266	0.21219	0.21208	0.21160	0.20702

; limit = ∞

C)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.72548	0.70888	0.70728	0.70693	0.70535	0.69007

; limit = 0.7071

D)

x	0.9	0.99	0.999	1.001	1.01	1.1
f(x)	0.21764	0.21266	0.21219	0.21208	0.21160	0.20702

; limit = 0.21213

88) If $f(x) = \sqrt{x} - 2$, find $\lim\limits_{x \to 4} f(x)$. 88) _____

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)						

A)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	3.9000	2.9000	1.9000	2.0000	3.0000	4.0000

; limit = 1.95

B)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	3.9000	2.9000	1.9000	2.0000	3.0000	4.0000

; limit = ∞

C)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	−0.02516	−0.00250	−0.00025	0.00025	0.00250	0.02485

; limit = 0

D)


x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	1.47736	1.49775	1.49977	1.50022	1.50225	1.52236

; limit = 1.50

Calculus 3rd edition Briggs Test Bank


For the function f whose graph is given, determine the limit.

89) Find $\lim_{x \to -1^-} f(x)$ and $\lim_{x \to -1^+} f(x)$. 89) _____



A) –2; –7 B) –7; –5 C) –5; –2 D) –7; –2


90) Find $\lim_{x \to 2^-} f(x)$ and $\lim_{x \to 2^+} f(x)$. 90) _____



A) 3; –1 B) does not exist; does not exist
C) –1; 3 D) 1; 1


Calculus 3rd edition Briggs Test Bank

91) Find $\lim_{x \to 2^+} f(x)$. 91) _____



A) 1.3 B) 4 C) 5 D) –1


92) Find $\lim_{x \to 1^-} f(x)$. 92) _____



A) –1 B) does not exist C) 2 D) $\frac{1}{2}$


Calculus 3rd edition Briggs Test Bank

93) Find $\lim_{x \to 1^+} f(x)$. 93) _____



A) 4 B) 3 C) does not exist D) $3\frac{1}{2}$


94) Find $\lim_{x \to 0} f(x)$. 94) _____



A) –2 B) 2 C) does not exist D) 0


Calculus 3rd edition Briggs Test Bank

95) Find $\lim_{x \to 0} f(x)$. 95) _____



A) 3 B) 0 C) does not exist D) –3


96) Find $\lim_{x \to 0} f(x)$. 96) _____



A) 2 B) 0 C) does not exist D) –2


Calculus 3rd edition Briggs Test Bank

97) Find $\lim_{x \to -1} f(x)$. 97) _____



A) does not exist B) –1 C) $-\frac{1}{2}$ D) $\frac{1}{2}$

98) Find $\lim_{x \to \infty} f(x)$. 98) _____



A) ∞ B) –2 C) 0 D) does not exist

Find the limit.

99) $\lim_{x \to -2} \frac{1}{x + 2}$ 99) _____

A) 1/2 B) Does not exist C) –∞ D) ∞

100) $\lim_{x \to -10^-} \frac{1}{x + 10}$ 100) _____

A) –1 B) -∞ C) ∞ D) 0

101) $\lim_{x \to 8^-} \frac{1}{(x - 8)^2}$ 101) _____

A) 0 B) –1 C) ∞ D) –∞

Calculus 3rd edition Briggs Test Bank

102) $\lim_{x \to -3^-} \frac{1}{x^2 - 9}$ 102) _____

A) ∞ B) –1 C) –∞ D) 0

103) $\lim_{x \to 2^+} \frac{3}{x^2 - 4}$ 103) _____

A) 0 B) –∞ C) 1 D) ∞

104) $\lim_{x \to (\pi/2)^+} \tan x$ 104) _____

A) 1 B) ∞ C) –∞ D) 0

105) $\lim_{x \to (-\pi/2)^-} \sec x$ 105) _____

A) –∞ B) ∞ C) 1 D) 0

106) $\lim_{x \to 0^+} (1 + \csc x)$ 106) _____

A) 1 B) ∞ C) 0 D) Does not exist

107) $\lim_{x \to 0} (1 - \cot x)$ 107) _____

A) ∞ B) 0 C) –∞ D) Does not exist

108) $\lim_{x \to -3^-} \frac{x^2 - 4x + 3}{x^3 - 9x}$ 108) _____

A) Does not exist B) 0 C) -∞ D) ∞

109) $\lim_{x \to 0^+} \frac{x^2 - 5x + 6}{x^3 - 9x}$ 109) _____

A) ∞ B) -∞ C) Does not exist D) 0

Find all vertical asymptotes of the given function.

110) $h(x) = \frac{7x}{x + 6}$ 110) _____

A) x = 7 B) none C) x = –6 D) x = 6

111) $f(x) = \frac{x + 2}{x^2 - 49}$ 111) _____

A) x = 49, x = –2 B) x = 0, x = 49
C) x = –7, x = 7 D) x = –7, x = 7, x = –2

Calculus 3rd edition Briggs Test Bank

112) $f(x) = \dfrac{x + 9}{x^2 + 36}$ 112) _____

A) $x = -6, x = -9$
B) $x = -6, x = 6$
C) none
D) $x = -6, x = 6, x = -9$

113) $g(x) = \dfrac{x + 11}{x^2 - 16x}$ 113) _____

A) $x = 0, x = 16$
B) $x = 0, x = -4, x = 4$
C) $x = -4, x = 4$
D) $x = 16, x = -11$

114) $f(x) = \dfrac{x - 1}{x^3 + 25x}$ 114) _____

A) $x = 0, x = -25$
B) $x = 0, x = -5, x = 5$
C) $x = 0$
D) $x = -5, x = 5$

115) $R(x) = \dfrac{-3x^2}{x^2 + 2x - 8}$ 115) _____

A) $x = 4, x = -2$
B) $x = -4, x = 2, x = -3$
C) $x = -4, x = 2$
D) $x = -8$

116) $R(x) = \dfrac{x - 1}{x^3 + 4x^2 - 77x}$ 116) _____

A) $x = -11, x = 0, x = 7$
B) $x = -7, x = -30, x = 11$
C) $x = -7, x = 0, x = 11$
D) $x = -11, \ x = 7$

117) $f(x) = \dfrac{-2x(x + 2)}{4x^2 - 3x - 7}$ 117) _____

A) $x = \dfrac{4}{7}, x = -1$
B) $x = -\dfrac{4}{7}, x = 1$
C) $x = \dfrac{7}{4}, x = -1$
D) $x = -\dfrac{7}{4}, x = 1$

118) $f(x) = \dfrac{x - 7}{49x - x^3}$ 118) _____

A) $x = 0, x = -7$
B) $x = 0, \ x = 7$
C) $x = -7, x = 7$
D) $x = 0, x = -7, x = 7$

119) $f(x) = \dfrac{-x^2 + 16}{x^2 + 5x + 4}$ 119) _____


A) $x = -1$
B) $x = 1, x = -4$
C) $x = -1, x = -4$
D) $x = -1, x = 4$

Calculus 3rd edition Briggs Test Bank


Choose the graph that represents the given function without using a graphing utility.

120) $f(x) = \dfrac{x}{x+3}$ 120) _____


A)




B)



C)




D)




Calculus 3rd edition Briggs Test Bank

121) $f(x) = \dfrac{x}{x^2 + x + 4}$ 121) _____


A)




B)



C)




D)




Calculus 3rd edition Briggs Test Bank

122) $f(x) = \dfrac{x^2 + 2}{x^3}$ 122) _____


A)




B)



C)




D)




Calculus 3rd edition Briggs Test Bank

123) $f(x) = \dfrac{1}{x + 1}$ 123) _____


A)




B)



C)




D)




Calculus 3rd edition Briggs Test Bank

124) $f(x) = \dfrac{x - 1}{x + 1}$ 124) _____


A)




B)



C)




D)




Calculus 3rd edition Briggs Test Bank

125) $f(x) = \dfrac{1}{(x+2)^2}$ 125) _____


A)




B)



C)




D)




Calculus 3rd edition Briggs Test Bank

126) $f(x) = \dfrac{2x^2}{4 - x^2}$ 126) _____


A)




B)



C)



D)



Find the limit.

127) $\lim\limits_{x \to -\infty} (-4x^{18} + 15)$ 127) _____

A) ∞ B) $-\infty$ C) 0 D) 15

128) $\lim\limits_{x \to -\infty} 4x^{-9}$ 128) _____

A) -4 B) ∞ C) 0 D) $-\infty$

129) $\lim\limits_{x \to \infty} 2x^8 - 15x^5$ 129) _____

A) -13 B) ∞ C) 0 D) $-\infty$

130) $\lim\limits_{x \to \infty} \dfrac{1}{x} - 4$ 130) _____

A) -5 B) -3 C) 4 D) -4

Calculus 3rd edition Briggs Test Bank

131) $\lim_{x \to -\infty} \frac{9}{6 - (4/x^2)}$ 131) _____

A) $-\infty$ B) $\frac{3}{2}$ C) $\frac{9}{2}$ D) 9

132) $\lim_{x \to -\infty} \frac{-1 + (5/x)}{2 - (1/x^2)}$ 132) _____

A) ∞ B) $-\frac{1}{2}$ C) $-\infty$ D) $\frac{1}{2}$

133) $\lim_{x \to \infty} \frac{x^2 + 5x + 6}{x^3 + 5x^2 + 11}$ 133) _____

A) 1 B) ∞ C) 0 D) $\frac{6}{11}$

134) $\lim_{x \to -\infty} \frac{-19x^2 - 4x + 7}{-10x^2 + 3x + 13}$ 134) _____

A) $\frac{7}{13}$ B) ∞ C) 1 D) $\frac{19}{10}$

135) $\lim_{x \to \infty} \frac{2x + 1}{16x - 7}$ 135) _____

A) $-\frac{1}{7}$ B) 0 C) ∞ D) $\frac{1}{8}$

136) $\lim_{x \to \infty} \frac{9x^3 - 6x^2 + 3x}{-x^3 - 2x + 7}$ 136) _____

A) 9 B) –9 C) $\frac{3}{2}$ D) ∞

137) $\lim_{x \to -\infty} \frac{6x^3 + 4x^2}{x - 5x^2}$ 137) _____

A) $-\frac{4}{5}$ B) 6 C) $-\infty$ D) ∞

138) $\lim_{x \to -\infty} \frac{\cos 5x}{x}$ 138) _____

A) $-\infty$ B) 5 C) 1 D) 0

Calculus 3rd edition Briggs Test Bank

Divide numerator and denominator by the highest power of x in the denominator to find the limit.

139) $\lim_{x\to\infty} \sqrt{\frac{49x^2}{2+9x^2}}$ 139) _____

A) $\frac{49}{9}$ B) $\frac{7}{3}$ C) $\frac{49}{2}$ D) does not exist

140) $\lim_{x\to\infty} \sqrt{\frac{64x^2+x-3}{(x-17)(x+1)}}$ 140) _____

A) ∞ B) 64 C) 0 D) 8

141) $\lim_{x\to\infty} \frac{4\sqrt{x}+x^{-1}}{3x+5}$ 141) _____

A) ∞ B) $\frac{4}{3}$ C) $\frac{1}{3}$ D) 0

142) $\lim_{x\to\infty} \frac{-4x^{-1}+3x^{-3}}{-2x^{-2}+x^{-5}}$ 142) _____

A) ∞ B) 0 C) 2 D) -∞

143) $\lim_{x\to-\infty} \frac{\sqrt[3]{x}+2x+4}{-4x+x^{2/3}-2}$ 143) _____

A) $-\frac{1}{2}$ B) 0 C) – 2 D) -∞

144) $\lim_{t\to\infty} \frac{\sqrt{100t^2-1{,}000}}{t-10}$ 144) _____

A) 100 B) does not exist C) 1,000 D) 10

145) $\lim_{t\to\infty} \frac{\sqrt{100t^2-1{,}000}}{t-10}$ 145) _____

A) 1,000 B) 10 C) does not exist D) 100

146) $\lim_{x\to\infty} \frac{5x+7}{\sqrt{7x^2+1}}$ 146) _____

A) ∞ B) 0 C) $\frac{5}{7}$ D) $\frac{5}{\sqrt{7}}$

Determine the limit.

147) $\lim_{x\to\infty} \left(\frac{12x^2+7x+1}{\sqrt{4x^4+x^3}}\right)$ 147) _____

A) 4 B) 3 C) 0 D) 6

Calculus 3rd edition Briggs Test Bank

148) $\lim_{x \to \infty} \left(x - \sqrt{x^2 - 6x}\right)$ (Hint: Multiply by $\frac{x + \sqrt{x^2 - 6x}}{x + \sqrt{x^2 - 6x}}$ first.) 148) _____

A) 0 B) 6 C) $\sqrt{6}$ D) 3

Find all horizontal asymptotes of the given function, if any.

149) $h(x) = \frac{2x - 2}{x - 7}$ 149) _____

A) $y = 7$
B) $y = 0$
C) $y = 2$
D) no horizontal asymptotes

150) $h(x) = 5 - \frac{8}{x}$ 150) _____

A) $x = 0$
B) $y = 5$
C) $y = 8$
D) no horizontal asymptotes

151) $g(x) = \frac{x^2 + 9x - 6}{x - 6}$ 151) _____

A) $y = 6$
B) $y = 1$
C) $y = 0$
D) no horizontal asymptotes

152) $h(x) = \frac{8x^2 - 3x - 5}{2x^2 - 2x + 4}$ 152) _____

A) $y = 0$
B) $y = 4$
C) $y = \frac{3}{2}$
D) no horizontal asymptotes

153) $h(x) = \frac{9x^4 - 5x^2 - 4}{3x^5 - 3x + 4}$ 153) _____

A) $y = \frac{5}{3}$
B) $y = 3$
C) $y = 0$
D) no horizontal asymptotes

154) $h(x) = \frac{6x^3 - 5x}{9x^3 - 3x + 7}$ 154) _____

A) $y = \frac{5}{3}$
B) $y = 0$
C) $y = \frac{2}{3}$
D) no horizontal asymptotes

Calculus 3rd edition Briggs Test Bank

155) $h(x) = \dfrac{8x^3 - 6x - 5}{9x^2 + 6}$ 155) _____

A) $y = 8$ B) $y = 0$

C) $y = \dfrac{8}{9}$ D) no horizontal asymptotes

156) $h(x) = \dfrac{6x + 1}{x^2 - 4}$ 156) _____

A) $y = 6$ B) no horizontal asymptotes
C) $y = -2, y = 2$ D) $y = 0$

157) $R(x) = \dfrac{-3x^2 + 1}{x^2 + 6x - 72}$ 157) _____

A) $y = -3$ B) $y = -12, y = 6$
C) $y = 0$ D) no horizontal asymptotes

158) $f(x) = \dfrac{x^2 - 4}{16x - x^4}$ 158) _____

A) no horizontal asymptotes B) $y = -4, y = 4$
C) $y = 0$ D) $y = -1$


159) $f(x) = \dfrac{49x^4 + x^2 - 7}{x - x^3}$ 159) _____

A) $y = 0$ B) $y = -49$
C) no horizontal asymptotes D) $y = -1, y = 1$

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.


Sketch the graph of a function y = f(x) that satisfies the given conditions.

160) $f(0) = 0, f(1) = 4, f(-1) = -4, \lim\limits_{x \to -\infty} f(x) = -3, \lim\limits_{x \to \infty} f(x) = 3.$ 160) _____




Calculus 3rd edition Briggs Test Bank


161) $f(0) = 0, f(1) = 2, f(-1) = 2, \lim_{x \to \pm\infty} f(x) = -2.$ 161) _____



162) $f(0) = 4, f(1) = -4, f(-1) = -4, \lim_{x \to \pm\infty} f(x) = 0.$ 162) _____




163) $f(0) = 0, \lim_{x \to \pm\infty} f(x) = 0, \lim_{x \to 6^-} f(x) = -\infty, \lim_{x \to -6^+} f(x) = -\infty, \lim_{x \to 6^+} f(x) = \infty, \lim_{x \to -6^-} f(x) = \infty.$ 163) _____



Calculus 3rd edition Briggs Test Bank


MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find all points where the function is discontinuous.

164) 


164) _____

A) x = 4, x = 2 B) x = 2 C) None D) x = 4

165) 


165) _____

A) x = −2 B) x = −2, x = 1 C) None D) x = 1

166) 


166) _____

A) x = −2, x = 0, x = 2 B) x = 2
C) x = 0, x = 2 D) x = −2, x = 0

167) 

167) _____


A) None B) x = 6 C) x = −2, x = 6 D) x = −2

168) 

168) _____


A) x = 1, x = 4, x = 5 B) x = 1, x = 5
C) None D) x = 4

Calculus 3rd edition Briggs Test Bank

169) 


169) _____

A) x = 1 B) None C) x = 0, x = 1 D) x = 0

170) 


170) _____

A) x = 0, x = 3 B) x = 0 C) x = 3 D) None

171) 

171) _____

A) x = 2 B) x = –2, x = 2 C) x = –2 D) None

172) 

172) _____


A) None B) x = 0
C) x = –2, x = 2 D) x = –2, x = 0, x = 2

Calculus 3rd edition Briggs Test Bank

Provide an appropriate response.

173) Is f continuous at f(1)? 173) _____


$$f(x) = \begin{cases} -x^2 + 1, & -1 \le x < 0 \\ 5x, & 0 < x < 1 \\ -2, & x = 1 \\ -5x + 10 & 1 < x < 3 \\ 5, & 3 < x < 5 \end{cases}$$



A) No B) Yes

174) Is f continuous at f(0)? 174) _____


$$f(x) = \begin{cases} -x^2 + 1, & -1 \le x < 0 \\ 2x, & 0 < x < 1 \\ -3, & x = 1 \\ -2x + 4 & 1 < x < 3 \\ 4, & 3 < x < 5 \end{cases}$$



A) No B) Yes

175) Is f continuous at x = 0? 175) _____

$$f(x) = \begin{cases} x^3, & -2 < x \le 0 \\ -2x, & 0 \le x < 2 \\ 7, & 2 < x \le 4 \\ 0, & x = 2 \end{cases}$$




A) No B) Yes

Calculus 3rd edition Briggs Test Bank

176) Is f continuous at x = 4? 176) _____

$$f(x) = \begin{cases} x^3, & -2 < x \le 0 \\ -4x, & 0 \le x < 2 \\ 6, & 2 < x \le 4 \\ 0, & x = 2 \end{cases}$$



A) Yes B) No

Find the intervals on which the function is continuous.

177) $y = \frac{2}{x + 3} - 5x$ 177) _____

A) discontinuous only when x = –3
B) discontinuous only when x = –8
C) discontinuous only when x = 3
D) continuous everywhere

178) $y = \frac{3}{(x + 1)^2 + 2}$ 178) _____

A) continuous everywhere
B) discontinuous only when x = 3
C) discontinuous only when x = –1
D) discontinuous only when x = –8

179) $y = \frac{x + 1}{x^2 - 11x + 28}$ 179) _____

A) discontinuous only when x = 4
B) discontinuous only when x = –7 or x = 4
C) discontinuous only when x = –4 or x = 7
D) discontinuous only when x = 4 or x = 7

180) $y = \frac{1}{x^2 - 16}$ 180) _____

A) discontinuous only when x = –4 or x = 4
B) discontinuous only when x = –16 or x = 16
C) discontinuous only when x = 16
D) discontinuous only when x = –4

181) $y = \frac{3}{|x| + 1} - \frac{x^2}{7}$ 181) _____

A) discontinuous only when x = –7 or x = –1
B) discontinuous only when x = –1
C) continuous everywhere
D) discontinuous only when x = –8

Calculus 3rd edition Briggs Test Bank

182) $y = \dfrac{\sin(5\theta)}{2\theta}$ 182) _____

A) discontinuous only when $\theta = \dfrac{\pi}{2}$

B) discontinuous only when $\theta = 0$

C) continuous everywhere

D) discontinuous only when $\theta = \pi$

183) $y = \dfrac{4\cos\theta}{\theta + 2}$ 183) _____

A) continuous everywhere

B) discontinuous only when $\theta = -2$

C) discontinuous only when $\theta = \dfrac{\pi}{2}$

D) discontinuous only when $\theta = 2$

184) $y = \sqrt{4x + 1}$ 184) _____

A) continuous on the interval $\left(-\dfrac{1}{4}, \infty\right)$

B) continuous on the interval $\left(-\infty, -\dfrac{1}{4}\right]$

C) continuous on the interval $\left[-\dfrac{1}{4}, \infty\right)$

D) continuous on the interval $\left[\dfrac{1}{4}, \infty\right)$

185) $y = \sqrt[4]{6x - 3}$ 185) _____

A) continuous on the interval $\left(\dfrac{1}{2}, \infty\right)$

B) continuous on the interval $\left[-\dfrac{1}{2}, \infty\right)$

C) continuous on the interval $\left[\dfrac{1}{2}, \infty\right)$

D) continuous on the interval $\left(-\infty, \dfrac{1}{2}\right]$

186) $y = \sqrt{x^2 - 6}$ 186) _____

A) continuous everywhere

B) continuous on the interval $[\sqrt{6}, \infty)$

C) continuous on the intervals $(-\infty, -\sqrt{6}]$ and $[\sqrt{6}, \infty)$


D) continuous on the interval $[-\sqrt{6}, \sqrt{6}]$

Calculus 3rd edition Briggs Test Bank

Provide an appropriate response.

187) Is f continuous on (–2, 4]? 187) _____

$$f(x) = \begin{cases} x^3, & -2 < x \le 0 \\ -3x, & 0 \le x < 2 \\ 7, & 2 < x \le 4 \\ 0, & x = 2 \end{cases}$$



A) Yes B) No

Find the limit, if it exists.

188) $\lim\limits_{x \to -3} (x^2 - 16 + \sqrt[3]{x^2 - 36})$ 188) _____

A) –10 B) Does not exist C) 4 D) –4

189) $\lim\limits_{x \to 7} \sqrt{x^2 + 2x + 1}$ 189) _____

A) ±8 B) 8 C) Does not exist D) 64

190) $\lim\limits_{x \to 1} \sqrt{x - 4}$ 190) _____

A) 1.73205081 B) 0 C) Does not exist D) –1.7320508

191) $\lim\limits_{x \to 4} \sqrt{x^2 - 9}$ 191) _____

A) $\pm\sqrt{7}$ B) 3.5 C) $\sqrt{7}$ D) Does not exist

192) $\lim\limits_{x \to -6^-} \sqrt{x^2 - 36}$ 192) _____

A) $6\sqrt{2}$ B) 3 C) 0 D) Does not exist

193) $\lim\limits_{x \to 5^+} \dfrac{3\sqrt{(x - 5)^3}}{x - 5}$ 193) _____

A) 3 B) $3\sqrt{5}$ C) 0 D) Does not exist

194) $\lim\limits_{t \to 1^+} \dfrac{\sqrt{(t + 9)(t - 1)^2}}{7t - 7}$ 194) _____

A) $\dfrac{1}{7}$ B) $\dfrac{\sqrt{10}}{7}$ C) 0 D) Does not exist

Calculus 3rd edition Briggs Test Bank

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

Provide an appropriate response.

195) Use the Intermediate Value Theorem to prove that $9x^3 + 3x^2 - 9x - 8 = 0$ has a solution between 1 and 2. 195) _____

196) Use the Intermediate Value Theorem to prove that $5x^4 + 8x^3 + 8x - 1 = 0$ has a solution between -3 and -2. 196) _____

197) Use the Intermediate Value Theorem to prove that $x(x - 8)^2 = 8$ has a solution between 7 and 9. 197) _____

198) Use the Intermediate Value Theorem to prove that $5 \sin x = x$ has a solution between $\frac{\pi}{2}$ and π. 198) _____

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find numbers a and b, or k, so that f is continuous at every point.

199) 199) _____

$$f(x) = \begin{cases} -27, & x < -4 \\ ax + b, & -4 \le x \le 5 \\ 45, & x > 5 \end{cases}$$

A) $a = 8, b = 85$ B) $a = -27, b = 45$ C) $a = 8, b = 5$ D) Impossible

200) 200) _____

$$f(x) = \begin{cases} x^2, & x < -1 \\ ax + b, & -1 \le x \le 3 \\ x + 6, & x > 3 \end{cases}$$

A) $a = -2, b = 3$ B) $a = 2, b = -3$ C) $a = 2, b = 3$ D) Impossible

201) 201) _____

$$f(x) = \begin{cases} 6x + 8, \text{ if } x < -10 \\ kx + 6, \text{ if } x \ge -10 \end{cases}$$

A) $k = \frac{29}{5}$ B) $k = -\frac{3}{5}$ C) $k = 8$ D) $k = \frac{3}{5}$

202) 202) _____

$$f(x) = \begin{cases} x^2, & \text{if } x \le 3 \\ x + k, & \text{if } x > 3 \end{cases}$$

A) $k = -3$ B) $k = 6$ C) $k = 12$ D) Impossible

Calculus 3rd edition Briggs Test Bank

203) 203) _____

$$f(x) = \begin{cases} x^2, & \text{if } x \le 7 \\ kx, & \text{if } x > 7 \end{cases}$$

A) $k = 49$ B) $k = 7$ C) $k = \frac{1}{7}$ D) Impossible

Solve the problem.

204) Select the correct statement for the definition of the limit: $\lim_{x \to x_0} f(x) = L$ 204) _____

means that _____

A) if given any number $\varepsilon > 0$, there exists a number $\delta > 0$, such that for all x, $0 < |x - x_0| < \varepsilon$ implies $|f(x) - L| < \delta$.

B) if given a number $\varepsilon > 0$, there exists a number $\delta > 0$, such that for all x, $0 < |x - x_0| < \delta$ implies $|f(x) - L| > \varepsilon$.

C) if given any number $\varepsilon > 0$, there exists a number $\delta > 0$, such that for all x, $0 < |x - x_0| < \delta$ implies $|f(x) - L| < \varepsilon$.

D) if given any number $\varepsilon > 0$, there exists a number $\delta > 0$, such that for all x, $0 < |x - x_0| < \varepsilon$ implies $|f(x) - L| > \delta$.

205) Identify the incorrect statements about limits. 205) _____

I. The number L is the limit of f(x) as x approaches x_0 if f(x) gets closer to L as x approaches x_0.


II. The number L is the limit of f(x) as x approaches x_0 if, for any $\varepsilon > 0$, there corresponds a $\delta > 0$ such that $|f(x) - L| < \varepsilon$ whenever $0 < |x - x_0| < \delta$.

III. The number L is the limit of f(x) as x approaches x_0 if, given any $\varepsilon > 0$, there exists a value of x for which $|f(x) - L| < \varepsilon$.

A) II and III B) I and II C) I and III D) I, II, and III

Use the graph to find a $\delta > 0$ such that for all x, $0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \varepsilon$.

206) 206) _____




NOT TO SCALE

$f(x) = 2x + 3$
$x_0 = 2$
$L = 7$
$\varepsilon = 0.2$

A) 5 B) 0.1 C) 0.4 D) 0.2

Calculus 3rd edition Briggs Test Bank

207) 207) _____




f(x) = 5x – 1
$x_0 = 1$
L = 4
ε = 0.2

NOT TO SCALE

A) 0.4 B) 0.08 C) 0.04 D) 3

208) 208) _____




f(x) = –5x – 2
$x_0 = -2$
L = 8
ε = 0.2

NOT TO SCALE

A) –0.04 B) 0.04 C) 14 D) 0.4

Calculus 3rd edition Briggs Test Bank

209) 209) _____




$f(x) = -2x + 3$
$x_0 = -2$
$L = 7$
$\varepsilon = 0.2$

NOT TO SCALE

A) 9 B) −0.1 C) 0.1 D) 0.2

210) 210) _____



$f(x) = \frac{4}{3}x + 2$
$x_0 = 1$
$L = 3.3$
$\varepsilon = 0.2$

NOT TO SCALE

A) 2.3 B) 0.1 C) −0.3 D) 0.3

Calculus 3rd edition Briggs Test Bank


211) 211) _____

$f(x) = -\frac{4}{3}x + 2$

$x_0 = -2$

$L = 4.7$

$\varepsilon = 0.2$



NOT TO SCALE

A) −0.3 B) 6.7 C) 0.3 D) 0.1


212) 212) _____

$f(x) = 2\sqrt{x}$

$x_0 = 3$

$L = 2\sqrt{3}$

$\varepsilon = \frac{1}{4}$




NOT TO SCALE

A) 0.46 B) 0.4169 C) 0.865 D) 0.4481

Calculus 3rd edition Briggs Test Bank

213) 213) _____




NOT TO SCALE

$f(x) = \sqrt{x - 3}$
$x_0 = 4$
$L = 1$
$\varepsilon = \frac{1}{4}$

A) 0.5625 B) 3 C) 0.4375 D) 1

214) 214) _____




NOT TO SCALE

$f(x) = 2x^2$
$x_0 = 1$
$L = 2$
$\varepsilon = 1$

A) 1 B) 0.22 C) 0.51 D) 0.29

Calculus 3rd edition Briggs Test Bank

215) 215) _____



$f(x) = x^2 - 3$
$x_0 = 3$
$L = 6$
$\varepsilon = 1$

NOT TO SCALE

A) 0.17 B) 0.33 C) 3 D) 0.16

A function f(x), a point x_0, the limit of f(x) as x approaches x_0, and a positive number ε is given. Find a number $\delta > 0$ such that for all x, $0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \varepsilon$.

216) $f(x) = 7x + 5$, $L = 19$, $x_0 = 2$, and $\varepsilon = 0.01$ 216) _____
A) 0.002857 B) 0.005 C) 0.007143 D) 0.001429

217) $f(x) = 7x - 2$, $L = 19$, $x_0 = 3$, and $\varepsilon = 0.01$ 217) _____
A) 0.001429 B) 0.003333 C) 0.000714 D) 0.002857

218) $f(x) = -9x + 2$, $L = -25$, $x_0 = 3$, and $\varepsilon = 0.01$ 218) _____
A) 0.004444 B) –0.003333 C) 0.001111 D) 0.002222

219) $f(x) = -9x - 8$, $L = -44$, $x_0 = 4$, and $\varepsilon = 0.01$ 219) _____
A) 0.002222 B) –0.0025 C) 0.001111 D) 0.000556

220) $f(x) = 10x^2$, $L = 640$, $x_0 = 8$, and $\varepsilon = 0.3$ 220) _____
A) 7.99812 B) 8.00187 C) 0.00188 D) 0.00187

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

Prove the limit statement

221) $\lim\limits_{x \to 5} (5x - 4) = 21$ 221) _____

222) $\lim\limits_{x \to 6} \dfrac{x^2 - 36}{x - 6} = 12$ 222) _____

223) $\lim\limits_{x \to 7} \dfrac{3x^2 - 17x - 28}{x - 7} = 25$ 223) _____

Calculus 3rd edition Briggs Test Bank

224) $\lim_{x \to 5} \frac{1}{x} = \frac{1}{5}$ 224) _____

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Use the precise definition of a limit to prove the limit. Specify a relationship between ε and δ that guarantees the limit exists.

225) $\lim_{x \to 2} x^4 = 16$ 225) _____

A) $\delta = \min\left\{1, \frac{\varepsilon}{32}\right\}$; Let $\varepsilon > 0$ and assume $0 < |x - 2| < \delta$. Then $|x^4 - 16| = |x^2 + 4|\,|x + 2|\,|x - 2| =$ $(8)(4)\frac{\varepsilon}{32} = \varepsilon$. That is, for any $\varepsilon > 0$, $|x^4 - 16| = \varepsilon$ whenever $0 < |x - 2| < \delta$, provided $0 < \delta \le \frac{\varepsilon}{32}$.

Therefore, $\lim_{x \to 2} x^4 = 16$.

B) $\delta = \min\left\{1, \frac{\varepsilon}{32}\right\}$; Let $\varepsilon > 0$ and assume $0 < |x - 2| < \delta$. Then $|x^4 - 16| = |x^2 + 4|\,|x + 2|\,|x - 2| <$ $(8)(4)\frac{\varepsilon}{32} = \varepsilon$. That is, for any $\varepsilon > 0$, $|x^4 - 16| < \varepsilon$ whenever $0 < |x-2| < \delta$, provided $0 < \delta \le \frac{\varepsilon}{32}$.

Therefore, $\lim_{x \to 2} x^4 = 16$.

C) $\delta = \min\left\{1, \frac{\varepsilon}{65}\right\}$; Let $\varepsilon > 0$ and assume $0 < |x - 2| < \delta$. Then $|x^4 - 16| = |x^2 + 4|\,|x + 2|\,|x - 2| =$ $(13)(5)\frac{\varepsilon}{65} = \varepsilon$. That is, for any $\varepsilon > 0$, $|x^4 - 16| = \varepsilon$ whenever $0 < |x - 2| < \delta$, provided $0 < \delta \le \frac{\varepsilon}{65}$.

Therefore, $\lim_{x \to 2} x^4 = 16$.

D) $\delta = \min\left\{1, \frac{\varepsilon}{65}\right\}$; Let $\varepsilon > 0$ and assume $0 < |x - 2| < \delta$. Then $|x^4 - 16| = |x^2 + 4|\,|x + 2|\,|x - 2| <$ $(13)(5)\frac{\varepsilon}{65} = \varepsilon$. That is, for any $\varepsilon > 0$, $|x^4 - 16| < \varepsilon$ whenever $0 < |x-2| < \delta$, provided $0 < \delta \le \frac{\varepsilon}{65}$.

Therefore, $\lim_{x \to 2} x^4 = 16$.

Calculus 3rd edition Briggs Test Bank

226) $\lim_{x \to 3} \frac{1}{x^2} = \frac{1}{9}$ 226) _____

A) $\delta = \min\left\{1, \frac{144\varepsilon}{7}\right\}$; Let $\varepsilon > 0$ and assume $0 < |x - 3| < \delta$. Then $\left|\frac{1}{x^2} - \frac{1}{9}\right| = \frac{|-1||x + 3||x - 3|}{|9x^2|} = \frac{7}{144}$

$\left(\frac{144\varepsilon}{7}\right) = \varepsilon$. That is, for any $\varepsilon > 0$, $\left|\frac{1}{x^2} - \frac{1}{9}\right| <= \varepsilon$ whenever $0 < |x - 3| < \delta$, provided $0 < \delta \leq \frac{144\varepsilon}{7}$.

Therefore, $\lim_{x \to 2} \frac{1}{x^2} = \frac{1}{9}$.

B) $\delta = \min\left\{1, \frac{36\varepsilon}{7}\right\}$; Let $\varepsilon > 0$ and assume $0 < |x - 3| < \delta$. Then $\left|\frac{1}{x^2} - \frac{1}{9}\right| = \frac{|-1|\,|x + 3|\,|x - 3|}{|9x^2|} < \frac{7}{36}$

$\left(\frac{36\varepsilon}{7}\right) = \varepsilon$. That is, for any $\varepsilon > 0$, $\left|\frac{1}{x^2} - \frac{1}{9}\right| < \varepsilon$ whenever $0 < |x - 3| < \delta$, provided $0 < \delta \leq \frac{36\varepsilon}{7}$.

Therefore, $\lim_{x \to 2} \frac{1}{x^2} = \frac{1}{9}$.

C) $\delta = \min\left\{1, \frac{36\varepsilon}{7}\right\}$; Let $\varepsilon > 0$ and assume $0 < |x - 3| < \delta$. Then $\left|\frac{1}{x^2} - \frac{1}{9}\right| = \frac{|-1||x + 3||x - 3|}{|9x^2|} = \frac{7}{36}$

$\left(\frac{36\varepsilon}{7}\right) = \varepsilon$. That is, for any $\varepsilon > 0$, $\left|\frac{1}{x^2} - \frac{1}{9}\right| = \varepsilon$ whenever $0 < |x - 3| < \delta$, provided $0 < \delta \leq \frac{36\varepsilon}{7}$.

Therefore, $\lim_{x \to 2} \frac{1}{x^2} = \frac{1}{9}$.

D) $\delta = \min\left\{1, \frac{144\varepsilon}{7}\right\}$; Let $\varepsilon > 0$ and assume $0 < |x - 3| < \delta$. Then $\left|\frac{1}{x^2} - \frac{1}{9}\right| = \frac{|-1|\,|x + 3||x - 3|}{|9x^2|} < \frac{7}{144}$

$\left(\frac{144\varepsilon}{7}\right) = \varepsilon$. That is, for any $\varepsilon > 0$, $\left|\frac{1}{x^2} - \frac{1}{9}\right| < \varepsilon$ whenever $0 < |x - 3| < \delta$, provided $0 < \delta \leq \frac{144\varepsilon}{7}$.

Therefore, $\lim_{x \to 2} \frac{1}{x^2} = \frac{1}{9}$.

Calculus 3rd edition Briggs Test Bank

1) C
2) B
3) D
4) A
5) D
6) B
7) D
8) B
9) D
10) B
11) A
12) B
13) C
14) C
15) D
16) B
17) D
18) D
19) A
20) C
21) D
22) D
23) C
24) D
25) B
26) B
27) D
28) D
29) B
30) D
31) C
32) C
33) B
34) C
35) D
36) D
37) C
38) B
39) B

40) Answers may vary. One possibility: $\lim_{x \to 0} 1 - \frac{x^2}{6} = \lim_{x \to 0} 1 = 1$. According to the squeeze theorem, the function $\frac{x \sin(x)}{2 - 2\cos(x)}$, which is squeezed between $1 - \frac{x^2}{6}$ and 1, must also approach 1 as x approaches 0. Thus, $\lim_{x \to 0} \frac{x \sin(x)}{2 - 2\cos(x)} = 1$.

41) D
42) B
43) C

Calculus 3rd edition Briggs Test Bank


44) D
45) B
46) D
47) D
48) B
49) D
50) C
51) C
52) C
53) B
54) A
55) C
56) C
57) C
58) D
59) A
60) D
61) D
62) C
63) B
64) B
65) B
66) B
67) C
68) C
69) C
70) D
71) C
72) B
73) D
74) C
75) C
76) B
77) B
78) C
79) D
80) C
81) A
82) D
83) D
84) C
85) C
86) B
87) C
88) C
89) A
90) A
91) B
92) C
93) B

Calculus 3rd edition Briggs Test Bank


94) C
95) B
96) C
97) D
98) A
99) B
100) B
101) C
102) A
103) D
104) C
105) B
106) B
107) D
108) C
109) B
110) C
111) C
112) C
113) A
114) C
115) C
116) A
117) C
118) A
119) A
120) A
121) C
122) D
123) D
124) A
125) C
126) D
127) B
128) C
129) B
130) D
131) B
132) B
133) C
134) D
135) D
136) B
137) D
138) D
139) B
140) D
141) D
142) A
143) A

Calculus 3rd edition Briggs Test Bank

144) D
145) B
146) D
147) D
148) D
149) C
150) B
151) D
152) B
153) C
154) C
155) D
156) D
157) A
158) C
159) C
160) Answers may vary. One possible answer:




161) Answers may vary. One possible answer:




Calculus 3rd edition Briggs Test Bank

162) Answers may vary. One possible answer:



163) Answers may vary. One possible answer:



164) D
165) D
166) A
167) B
168) C
169) B
170) C
171) B
172) B
173) A
174) A
175) B
176) A
177) A
178) A
179) D
180) A
181) C
182) B
183) B
184) C

Calculus 3rd edition Briggs Test Bank

185) C
186) C
187) B
188) A
189) B
190) C
191) C
192) C
193) C
194) B
195) Let $f(x) = 9x^3 + 3x^2 - 9x - 8$ and let $y_0 = 0$. $f(1) = -5$ and $f(2) = 58$. Since f is continuous on [1, 2] and since $y_0 = 0$ is between f(1) and f(2), by the Intermediate Value Theorem, there exists a c in the interval (1 , 2) with the property that $f(c) = 0$. Such a c is a solution to the equation $9x^3 + 3x^2 - 9x - 8 = 0$.
196) Let $f(x) = 5x^4 + 8x^3 + 8x - 1$ and let $y_0 = 0$. $f(-3) = 164$ and $f(-2) = -1$. Since f is continuous on [–3, –2] and since $y_0 = 0$ is between f(–3) and f(–2), by the Intermediate Value Theorem, there exists a c in the interval (–3, –2) with the property that $f(c) = 0$. Such a c is a solution to the equation $5x^4 + 8x^3 + 8x - 1 = 0$.
197) Let $f(x) = x(x - 8)^2$ and let $y_0 = 8$. $f(7) = 7$ and $f(9) = 9$. Since f is continuous on [7, 9] and since $y_0 = 8$ is between f(7) and f(9), by the Intermediate Value Theorem, there exists a c in the interval (7, 9) with the property that $f(c) = 8$. Such a c is a solution to the equation $x(x - 8)^2 = 8$.
198) Let $f(x) = \frac{\sin x}{x}$ and let $y_0 = \frac{1}{5}$. $f\left(\frac{\pi}{2}\right) \approx 0.6366$ and $f(\pi) = 0$. Since f is continuous on $\left[\frac{\pi}{2}, \pi\right]$ and since $y_0 = \frac{1}{5}$ is between $f\left(\frac{\pi}{2}\right)$ and $f(\pi)$, by the Intermediate Value Theorem, there exists a c in the interval $\left(\frac{\pi}{2}, \pi\right)$, with the property that $f(c) = \frac{1}{5}$. Such a c is a solution to the equation $5 \sin x = x$.
199) C
200) C
201) A
202) B
203) B
204) C
205) C
206) B
207) C
208) B
209) C
210) B
211) D
212) B
213) C
214) B
215) D
216) D
217) A
218) C
219) C
220) D

Calculus 3rd edition Briggs Test Bank

221)
Let $\varepsilon > 0$ be given. Choose $\delta = \varepsilon/5$. Then $0 < |x - 5| < \delta$ implies that

$$\begin{aligned} |(5x - 4) - 21| &= |5x - 25| \\ &= |5(x - 5)| \\ &= 5|x - 5| < 5\delta = \varepsilon \end{aligned}$$

Thus, $0 < |x - 5| < \delta$ implies that $|(5x - 4) - 21| < \varepsilon$

222) Let $\varepsilon > 0$ be given. Choose $\delta = \varepsilon$. Then $0 < |x - 6| < \delta$ implies that

$$\begin{aligned} \left|\frac{x^2 - 36}{x - 6} - 12\right| &= \left|\frac{(x - 6)(x + 6)}{x - 6} - 12\right| \\ &= |(x + 6) - 12| \quad \text{for } x \neq 6 \\ &= |x - 6| < \delta = \varepsilon \end{aligned}$$

Thus, $0 < |x - 6| < \delta$ implies that $\left|\frac{x^2 - 36}{x - 6} - 12\right| < \varepsilon$

223) Let $\varepsilon > 0$ be given. Choose $\delta = \varepsilon/3$. Then $0 < |x - 7| < \delta$ implies that

$$\begin{aligned} \left|\frac{3x^2 - 17x - 28}{x - 7} - 25\right| &= \left|\frac{(x - 7)(3x + 4)}{x - 7} - 25\right| \\ &= |(3x + 4) - 25| \quad \text{for } x \neq 7 \\ &= |3x - 21| \\ &= |3(x - 7)| \\ &= 3|x - 7| < 3\delta = \varepsilon \end{aligned}$$

Thus, $0 < |x - 7| < \delta$ implies that $\left|\frac{3x^2 - 17x - 28}{x - 7} - 25\right| < \varepsilon$

224) Let $\varepsilon > 0$ be given. Choose $\delta = \min\{5/2, 25\varepsilon/2\}$. Then $0 < |x - 5| < \delta$ implies that

$$\begin{aligned} \left|\frac{1}{x} - \frac{1}{5}\right| &= \left|\frac{5 - x}{5x}\right| \\ &= \frac{1}{|x|} \cdot \frac{1}{5} \cdot |x - 5| \\ &< \frac{1}{5/2} \cdot \frac{1}{5} \cdot \frac{25\varepsilon}{2} = \varepsilon \end{aligned}$$

Thus, $0 < |x - 5| < \delta$ implies that $\left|\frac{1}{x} - \frac{1}{5}\right| < \varepsilon$

225) D
226) B

Calculus 3rd edition Briggs Test Bank