Probability and Statistical Inference 10th edition Hogg Solutions Manual

SKU: 9780135189399-SOLUTIONS

INSTRUCTOR'S SOLUTIONS MANUAL

PROBABILITY AND STATISTICAL INFERENCE

TENTH EDITION


Robert V. Hogg

Elliot A. Tanis

Dale L. Zimmerman

Pearson

Probability and Statistical Inference 10th edition Hogg Solutions Manual



The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expressed or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson from electronic files supplied by the author.

Copyright © 2020, 2015, 2010 Pearson Education, Inc.
Publishing as Pearson, 330 Hudson Street, NY NY 10013

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.



ISBN-13: 978-0-13-518948-1
ISBN-10: 0-13-518948-9

Probability and Statistical Inference 10th edition Hogg Solutions Manual

Contents

Preface **v**

1 Probability **1**
1.1 Properties of Probability . . . 1
1.2 Methods of Enumeration . . . 2
1.3 Conditional Probability . . . 3
1.4 Independent Events . . . 4
1.5 Bayes' Theorem . . . 5

2 Discrete Distributions **7**
2.1 Random Variables of the Discrete Type . . . 7
2.2 Mathematical Expectation . . . 9
2.3 Special Mathematical Expectations . . . 11
2.4 The Binomial Distribution . . . 14
2.5 The Hypergeometric Distribution . . . 16
2.6 The Negative Binomial Distribution . . . 17
2.7 The Poisson Distribution . . . 18

3 Continuous Distributions **19**
3.1 Random Variables of the Continuous Type . . . 19
3.2 The Exponential, Gamma, and Chi-Square Distributions . . . 26
3.3 The Normal Distribution . . . 28
3.4 Additional Models . . . 30

4 Bivariate Distributions **33**
4.1 Bivariate Distributions of the Discrete Type . . . 33
4.2 The Correlation Coefficient . . . 34
4.3 Conditional Distributions . . . 36
4.4 Bivariate Distributions of the Continuous Type . . . 37
4.5 The Bivariate Normal Distribution . . . 41

5 Distributions of Functions of Random Variables **45**
5.1 Functions of One Random Variable . . . 45
5.2 Transformations of Two Random Variables . . . 47
5.3 Several Random Variables . . . 52
5.4 The Moment-Generating Function Technique . . . 53
5.5 Random Functions Associated with Normal Distributions . . . 55
5.6 The Central Limit Theorem . . . 58
5.7 Approximations for Discrete Distributions . . . 59
5.8 Chebyshev's Inequality and Convergence in Probability . . . 61
5.9 Limiting Moment-Generating Functions . . . 62

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

6 Point Estimation **63**
6.1 Descriptive Statistics . . . 63
6.2 Exploratory Data Analysis . . . 65
6.3 Order Statistics . . . 70
6.4 Maximum Likelihood Estimation . . . 73
6.5 A Simple Regression Problem . . . 76
6.6 Asymptotic Distributions of Maximum Likelihood Estimators . . . 81
6.7 Sufficient Statistics . . . 81
6.8 Bayesian Estimation . . . 84

7 Interval Estimation **87**
7.1 Confidence Intervals for Means . . . 87
7.2 Confidence Intervals for the Difference of Two Means . . . 88
7.3 Confidence Intervals For Proportions . . . 90
7.4 Sample Size . . . 91
7.5 Distribution-Free Confidence Intervals for Percentiles . . . 92
7.6 More Regression . . . 93
7.7 Resampling Methods . . . 99

8 Tests of Statistical Hypotheses **107**
8.1 Tests About One Mean . . . 107
8.2 Tests of the Equality of Two Means . . . 109
8.3 Tests for Variances . . . 111
8.4 Tests about Proportions . . . 113
8.5 Some Distribution-Free Tests . . . 114
8.6 Power of a Statistical Test . . . 118
8.7 Best Critical Regions . . . 121
8.8 Likelihood Ratio Tests . . . 124

9 More Tests **127**
9.1 Chi-Square Goodness-of-Fit Tests . . . 127
9.2 Contingency Tables . . . 130
9.3 One-Factor Analysis of Variance . . . 131
9.4 Two-Way Analysis of Variance . . . 134
9.5 General Factorial and 2^k Factorial Designs . . . 135
9.6 Tests Concerning Regression and Correlation . . . 136
9.7 Statistical Quality Control . . . 137

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

Preface

This solutions manual provides answers for the even-numbered exercises in *Probability and Statistical Inference*, tenth edition, by Robert V. Hogg, Elliot A. Tanis, and Dale L. Zimmerman. Complete solutions are given for most of these exercises. You, the instructor, may decide how many of these solutions and answers you want to make available to your students. Note that the answers for the odd-numbered exercises are given in the textbook. Our hope is that this solutions manual will be helpful to each of you in your teaching.

All of the figures in this manual were generated using *Maple*, a computer algebra system. Most of the figures were generated and many of the solutions, especially those involving data, were solved using procedures that were written by Zaven Karian from Denison University. We thank him for providing these. These procedures are available free of charge for your use. They are available for down load at http://www.math.hope.edu/tanis/. Short descriptions of these procedures are provided on the "Maple Card." Complete descriptions of these procedures are given in *Probability and Statistics: Explorations with MAPLE*, second edition, 1999, written by Zaven Karian and Elliot Tanis, published by Prentice Hall (ISBN 0-13-021536-8). You can download a slightly revised edition of this manual at http://www.math.hope.edu/tanis/MapleManual.pdf.

We also want to acknowledge the many suggestions/corrections that were made by our accuracy checker, Kyle Siegrist.

If you find an error or wish to make a suggestion, please send them to dale-zimmerman@uiowa.edu. These **errata** will be posted on http://homepage.divms.uiowa.edu/∼dzimmer/.

E.A.T.
D.L.Z.

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

Probability and Statistical Inference 10th edition Hogg Solutions Manual

Chapter 1

Probability

1.1 Properties of Probability

1.1-2 Sketch a figure and fill in the probabilities of each of the disjoint sets.

Let $A = \{\text{insure more than one car}\}$, $P(A) = 0.85$.

Let $B = \{\text{insure a sports car}\}$, $P(B) = 0.23$.

Let $C = \{\text{insure exactly one car}\}$, $P(C) = 0.15$.

It is also given that $P(A \cap B) = 0.17$. Since $A \cap C = \phi$, $P(A \cap C) = 0$. It follows that $P(A \cap B \cap C') = 0.17$. Thus $P(A' \cap B \cap C') = 0.06$ and $P(B' \cap C) = 0.09$.

1.1-4 **(a)** $S = \{$HHHH, HHHT, HHTH, HTHH, THHH, HHTT, HTTH, TTHH, HTHT, THTH, THHT, HTTT, THTT, TTHT, TTTH, TTTT$\}$;

(b) **(i)** 5/16, **(ii)** 0, **(iii)** 11/16, **(iv)** 4/16, **(v)** 4/16, **(vi)** 9/16, **(vii)** 4/16.

1.1-6 **(a)** $P(A \cup B) = 0.5 + 0.6 - 0.4 = 0.7$;

(b)
$$\begin{aligned} A &= (A \cap B') \cup (A \cap B) \\ P(A) &= P(A \cap B') + P(A \cap B) \\ 0.5 &= P(A \cap B') + 0.4 \\ P(A \cap B') &= 0.1; \end{aligned}$$

(c) $P(A' \cup B') = P[(A \cap B)'] = 1 - P(A \cap B) = 1 - 0.4 = 0.6$.

1.1-8 Let $A =$\{lab work done\}, $B =$\{referral to a specialist\},

$P(A) = 0.41$, $P(B) = 0.53$, $P([A \cup B]') = 0.21$.

$$\begin{aligned} P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ 0.79 &= 0.41 + 0.53 - P(A \cap B) \\ P(A \cap B) &= 0.41 + 0.53 - 0.79 = 0.15. \end{aligned}$$

1.1-10
$$\begin{aligned} A \cup B \cup C &= A \cup (B \cup C) \\ P(A \cup B \cup C) &= P(A) + P(B \cup C) - P[A \cap (B \cup C)] \\ &= P(A) + P(B) + P(C) - P(B \cap C) - P[(A \cap B) \cup (A \cap C)] \\ &= P(A) + P(B) + P(C) - P(B \cap C) - P(A \cap B) - P(A \cap C) \\ &\quad + P(A \cap B \cap C). \end{aligned}$$

1.1-12 **(a)** 1/3; **(b)** 2/3; **(c)** 0; **(d)** 1/2.

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

1.1-14 $P(A) = \dfrac{2[r - r(\sqrt{3}/2)]}{2r} = 1 - \dfrac{\sqrt{3}}{2}.$

1.1-16 Note that the respective probabilities are $p_0,\ p_1 = p_0/4,\ p_2 = p_0/4^2, \cdots$.

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{p_0}{4^k} &= 1 \\ \frac{p_0}{1 - 1/4} &= 1 \\ p_0 &= \frac{3}{4} \end{aligned}$$

$$1 - p_0 - p_1 = 1 - \frac{15}{16} = \frac{1}{16}.$$

1.2 Methods of Enumeration

1.2-2 **(a)** $(4)(5)(2) = 40$; **(b)** $(2)(2)(2) = 8$.

1.2-4 **(a)** $4\binom{6}{3} = 80$;

(b) $4(2^6) = 256$;

(c) $\dfrac{(4-1+3)!}{(4-1)!3!} = 20.$

1.2-6 $S =\{$ DDD, DDFD, DFDD, FDDD, DDFFD, DFDFD, FDDFD, DFFDD, FDFDD, FFDDD, FFF, FFDF, FDFF, DFFF FFDDF, FDFDF, DFFDF, FDDFF, DFDFF, DDFFF $\}$ so there are 20 possibilities. Note that the winning player (2 choices) must win the last set and two of the previous sets, so the number of outcomes is

$$2\left[\binom{2}{2} + \binom{3}{2} + \binom{4}{2}\right] = 20.$$

1.2-8 $3 \cdot 3 \cdot 2^{12} = 36{,}864.$

1.2-10
$$\begin{aligned} \binom{n-1}{r} + \binom{n-1}{r-1} &= \frac{(n-1)!}{r!(n-1-r)!} + \frac{(n-1)!}{(r-1)!(n-r)!} \\ &= \frac{(n-r)(n-1)! + r(n-1)!}{r!(n-r)!} = \frac{n!}{r!(n-r)!} = \binom{n}{r}. \end{aligned}$$

1.2-12
$$0 = (1-1)^n = \sum_{r=0}^{n} \binom{n}{r}(-1)^r(1)^{n-r} = \sum_{r=0}^{n} (-1)^r \binom{n}{r}.$$

$$2^n = (1+1)^n = \sum_{r=0}^{n} \binom{n}{r}(1)^r(1)^{n-r} = \sum_{r=0}^{n} \binom{n}{r}.$$

1.2-14 $\dbinom{5-1+29}{29} = \dfrac{33!}{29!4!} = 40{,}920.$

1.2-16 **(a)** $\dfrac{\dbinom{19}{3}\dbinom{52-19}{6}}{\dbinom{52}{9}} = \dfrac{102{,}486}{351{,}325} = 0.2917;$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

(b) $$\frac{\binom{19}{3}\binom{10}{2}\binom{7}{1}\binom{3}{0}\binom{5}{1}\binom{2}{0}\binom{6}{2}}{\binom{52}{9}} = \frac{7{,}695}{1{,}236{,}664} = 0.00622.$$

1.2-18 **(a)** $P(A) = \sum_{n=1}^{5}(1/2)^n = 1 - (1/2)^5$;

(b) $P(B) = \sum_{n=1}^{10}(1/2)^n = 1 - (1/2)^{10}$;

(c) $P(A \cup B) = P(B) = 1 - (1/2)^{10}$;

(d) $P(A \cap B) = P(A) = 1 - (1/2)^5$;

(e) $P(C) = P(B) - P(A) = (1/2)^5 - (1/2)^{10}$;

(f) $P(B') = 1 - P(B) = (1/2)^{10}$.

1.3 Conditional Probability

1.3-2 **(a)** $\frac{1041}{1456}$;

(b) $\frac{392}{633}$;

(c) $\frac{649}{823}$.

(d) The proportion of women who favor a gun law is greater than the proportion of men who favor a gun law.

1.3-4 **(a)** $P(\text{HH}) = \frac{13}{52} \cdot \frac{12}{51} = \frac{1}{17}$;

(b) $P(\text{HC}) = \frac{13}{52} \cdot \frac{13}{51} = \frac{13}{204}$;

(c) $P(\text{Non-Ace Heart, Ace}) + P(\text{Ace of Hearts, Non-Heart Ace})$

$$= \frac{12}{52} \cdot \frac{4}{51} + \frac{1}{52} \cdot \frac{3}{51} = \frac{51}{52 \cdot 51} = \frac{1}{52}.$$

1.3-6 Let $H = \{\text{died from heart disease}\}$; $P = \{\text{at least one parent had heart disease}\}$.

$$P(H \mid P') = \frac{N(H \cap P')}{N(P')} = \frac{110}{648}.$$

1.3-8 **(a)** $\frac{3}{20} \cdot \frac{2}{19} \cdot \frac{1}{18} = \frac{1}{1140}$;

(b) $\frac{\binom{3}{2}\binom{17}{1}}{\binom{20}{3}} \cdot \frac{1}{17} = \frac{1}{380}$;

(c) $\sum_{k=1}^{9} \frac{\binom{3}{2}\binom{17}{2k-2}}{\binom{20}{2k}} \cdot \frac{1}{20-2k} = \frac{35}{76} = 0.4605$;

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

(d) Draw second. The probability of winning is $1 - 0.4605 = 0.5395$.

1.3-10 **(a)** $P(A) = \frac{52}{52} \cdot \frac{51}{52} \cdot \frac{50}{52} \cdot \frac{49}{52} \cdot \frac{48}{52} \cdot \frac{47}{52} = \frac{8{,}808{,}975}{11{,}881{,}376} = 0.74141$;

(b) $P(A') = 1 - P(A) = 0.25859$.

1.3-12 **(a)** It doesn't matter because $P(B_1) = \frac{1}{18},\ P(B_5) = \frac{1}{18},\ P(B_{18}) = \frac{1}{18}$;

(b) $P(B) = \frac{2}{18} = \frac{1}{9}$ on each draw.

1.3-14 **(a)** $5 \cdot 4 \cdot 3 = 60$;

(b) $5 \cdot 5 \cdot 5 = 125$.

1.3-16 $\frac{3}{5} \cdot \frac{5}{8} + \frac{2}{5} \cdot \frac{4}{8} = \frac{23}{40}$.

1.4 Independent Events

1.4-2 **(a)**
$$\begin{aligned} P(A \cap B) &= P(A)P(B) = (0.3)(0.6) = 0.18; \\ P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ &= 0.3 + 0.6 - 0.18 \\ &= 0.72; \end{aligned}$$

(b) $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0}{0.6} = 0.$

1.4-4 Proof of **(b)**:
$$\begin{aligned} P(A' \cap B) &= P(B)P(A'|B) \\ &= P(B)[1 - P(A|B)] \\ &= P(B)[1 - P(A)] \\ &= P(B)P(A'). \end{aligned}$$

Proof of **(c)**:
$$\begin{aligned} P(A' \cap B') &= P[(A \cup B)'] \\ &= 1 - P(A \cup B) \\ &= 1 - P(A) - P(B) + P(A \cap B) \\ &= 1 - P(A) - P(B) + P(A)P(B) \\ &= [1 - P(A)][1 - P(B)] \\ &= P(A')P(B'). \end{aligned}$$

1.4-6
$$\begin{aligned} P[A \cap (B \cap C)] &= P[A \cap B \cap C] \\ &= P(A)P(B)P(C) \\ &= P(A)P(B \cap C). \end{aligned}$$

$$\begin{aligned} P[A \cap (B \cup C)] &= P[(A \cap B) \cup (A \cap C)] \\ &= P(A \cap B) + P(A \cap C) - P(A \cap B \cap C) \\ &= P(A)P(B) + P(A)P(C) - P(A)P(B)P(C) \\ &= P(A)[P(B) + P(C) - P(B \cap C)] \\ &= P(A)P(B \cup C). \end{aligned}$$

$$\begin{aligned} P[A' \cap (B \cap C')] &= P(A' \cap C' \cap B) \\ &= P(B)[P(A' \cap C') \,|\, B] \\ &= P(B)[1 - P(A \cup C \,|\, B)] \\ &= P(B)[1 - P(A \cup C)] \\ &= P(B)P[(A \cup C)'] \\ &= P(B)P(A' \cap C') \\ &= P(B)P(A')P(C') \\ &= P(A')P(B)P(C') \\ &= P(A')P(B \cap C'). \end{aligned}$$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

$$\begin{aligned} P[A' \cap B' \cap C'] &= P[(A \cup B \cup C)'] \\ &= 1 - P(A \cup B \cup C) \\ &= 1 - P(A) - P(B) - P(C) + P(A)P(B) + P(A)P(C) + \\ &\quad P(B)P(C) - PA)P(B)P(C) \\ &= [1 - P(A)][1 - P(B)][1 - P(C)] \\ &= P(A')P(B')P(C'). \end{aligned}$$

1.4-8 $\frac{1}{6} \cdot \frac{2}{6} \cdot \frac{3}{6} + \frac{1}{6} \cdot \frac{4}{6} \cdot \frac{3}{6} + \frac{5}{6} \cdot \frac{2}{6} \cdot \frac{3}{6} = \frac{2}{9}$.

1.4-10 **(a)** $\frac{3}{4} \cdot \frac{3}{4} = \frac{9}{16}$;

(b) $\frac{1}{4} \cdot \frac{3}{4} + \frac{3}{4} \cdot \frac{2}{4} = \frac{9}{16}$;

(c) $\frac{2}{4} \cdot \frac{1}{4} + \frac{2}{4} \cdot \frac{4}{4} = \frac{10}{16}$.

1.4-12 **(a)** $\left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2$;

(b) $\left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2$;

(c) $\left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2$;

(d) $\frac{5!}{3!\,2!} \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2$.

1.4-14 **(a)** $1 - (0.4)^3 = 1 - 0.064 = 0.936$;

(b) $1 - (0.4)^8 = 1 - 0.00065536 = 0.99934464$.

1.4-16 **(a)** $\sum_{k=0}^{\infty} \frac{1}{5} \left(\frac{4}{5}\right)^{2k} = \frac{5}{9}$;

(b) $\frac{1}{5} + \frac{4}{5} \cdot \frac{3}{4} \cdot \frac{1}{3} + \frac{4}{5} \cdot \frac{3}{4} \cdot \frac{2}{3} \cdot \frac{1}{2} \cdot \frac{1}{1} = \frac{3}{5}$.

1.4-18 **(a)** 7; **(b)** $(1/2)^7$; **(c)** 63; **(d)** No! $(1/2)^{63} = 1/9{,}223{,}372{,}036{,}854{,}775{,}808$.

1.4-20 No. The equations that must hold are

$$(1 - p_1)(1 - p_2) = p_1(1 - p_2) + p_2(1 - p_1) = p_1 p_2.$$

There are no real solutions.

1.5 Bayes' Theorem

1.5-2 **(a)** $\begin{aligned} P(G) &= P(A \cap G) + P(B \cap G) \\ &= P(A)P(G \mid A) + P(B)P(G \mid B) \\ &= (0.40)(0.85) + (0.60)(0.75) = 0.79; \end{aligned}$

(b) $P(A \mid G) = \frac{P(A \cap G)}{P(G)}$

$$= \frac{(0.40)(0.85)}{0.79} = 0.43.$$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

1.5-4 Let event B denote an accident and let A_1 be the event that age of the driver is 16–25. Then

$$\begin{aligned} P(A_1 \mid B) &= \frac{(0.1)(0.05)}{(0.1)(0.05)+(0.55)(0.02)+(0.20)(0.03)+(0.15)(0.04)} \\ &= \frac{50}{50+110+60+60} = \frac{50}{280} = 0.179. \end{aligned}$$

1.5-6 Let B be the event that the policyholder dies. Let A_1, A_2, A_3 be the events that the deceased is standard, preferred and ultra-preferred, respectively. Then

$$\begin{aligned} P(A_1 \mid B) &= \frac{(0.60)(0.01)}{(0.60)(0.01)+(0.30)(0.008)+(0.10)(0.007)} \\ &= \frac{60}{60+24+7} = \frac{60}{91} = 0.659; \\ P(A_2 \mid B) &= \frac{24}{91} = 0.264; \\ P(A_3 \mid B) &= \frac{7}{91} = 0.077. \end{aligned}$$

1.5-8 Let A be the event that the tablet is under warranty.

$$\begin{aligned} P(B_1 \mid A) &= \frac{(0.40)(0.10)}{(0.40)(0.10)+(0.30)(0.05)+(0.20)(0.03)+(0.10)(0.02)} \\ &= \frac{40}{40+15+6+2} = \frac{40}{63} = 0.635; \\ P(B_2 \mid A) &= \frac{15}{63} = 0.238; \\ P(B_3 \mid A) &= \frac{6}{63} = 0.095; \\ P(B_4 \mid A) &= \frac{2}{63} = 0.032. \end{aligned}$$

1.5-10 **(a)** $P(D^+) = (0.02)(0.92)+(0.98)(0.05) = 0.0184+0.0490 = 0.0674$;

(b) $P(A^- \mid D^+) = \dfrac{0.0490}{0.0674} = 0.727$; $P(A^+ \mid D^+) = \dfrac{0.0184}{0.0674} = 0.273$;

(c) $P(A^- \mid D^-) = \dfrac{(0.98)(0.95)}{(0.02)(0.08)+(0.98)(0.95)} = \dfrac{9310}{16+9310} = 0.998$;
$P(A^+ \mid D^-) = 0.002$;

(d) Yes, particularly those in part (b).

1.5-12 Let $D = \{\text{defective roll}\}$. Then

$$\begin{aligned} P(I \mid D) &= \frac{P(I \cap D)}{P(D)} \\ &= \frac{P(I)\cdot P(D \mid I)}{P(I)\cdot P(D \mid I)+P(II)\cdot P(D \mid II)} \\ &= \frac{(0.60)(0.03)}{(0.60)(0.03)+(0.40)(0.01)} \\ &= \frac{0.018}{0.018+0.004} = \frac{0.018}{0.022} = 0.818. \end{aligned}$$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

Chapter 2

Discrete Distributions

2.1 Random Variables of the Discrete Type

2.1-2 **(a)**

$$f(x) = \begin{cases} 0.6, & x = 1, \\ 0.3, & x = 5, \\ 0.1, & x = 10; \end{cases}$$

(b)




Figure 2.1–2: A probability histogram

2.1-4 **(a)**

$$\begin{aligned} \sum_{x=1}^{9} \log_{10}\left(\frac{x+1}{x}\right) &= \sum_{x=1}^{9} [\log_{10}(x+1) - \log_{10} x] \\ &= \log_{10} 2 - \log_{10} 1 + \log_{10} 3 - \log_{10} 2 + \cdots + \log_{10} 10 - \log_{10} 9 \\ &= \log_{10} 10 = 1; \end{aligned}$$

(b)

$$F(x) = \begin{cases} 0, & x < 1, \\ \log_{10} n, & n-1 \le x < n, \quad n = 2, 3, \ldots, 9, \\ 1, & 9 \le x. \end{cases}$$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

2.1-6 **(a)** $f(x) = \dfrac{1}{10}, \qquad x = 0, 1, 2, \cdots, 9;$

(b) $\mathcal{N}(\{0\})/150 = 11/150 = 0.073; \quad \mathcal{N}(\{5\})/150 = 13/150 = 0.087;$
$\mathcal{N}(\{1\})/150 = 14/150 = 0.093; \quad \mathcal{N}(\{6\})/150 = 22/150 = 0.147;$
$\mathcal{N}(\{2\})/150 = 13/150 = 0.087; \quad \mathcal{N}(\{7\})/150 = 16/150 = 0.107;$
$\mathcal{N}(\{3\})/150 = 12/150 = 0.080; \quad \mathcal{N}(\{8\})/150 = 18/150 = 0.120;$
$\mathcal{N}(\{4\})/150 = 16/150 = 0.107; \quad \mathcal{N}(\{9\})/150 = 15/150 = 0.100.$

(c)




Figure 2.1–6: Michigan daily lottery digits

2.1-8 **(a)** $f(x) = \dfrac{6 - |7 - x|}{36}, \quad x = 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12;$

(b)




Figure 2.1–8: Probability histogram for the sum of a pair of dice

Copyright © 2020 Pearson Education, Inc.
Probability and Statistical Inference 10th edition Hogg Solutions Manual

2.1-10 **(a)** The space of W is $S = \{0, 1, 2, 3, 4, 5, 6, 7\}$.

$$P(W = 0) = P(X = 0, Y = 0) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8}, \text{ assuming independence.}$$
$$P(W = 1) = P(X = 0, Y = 1) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$
$$P(W = 2) = P(X = 2, Y = 0) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$
$$P(W = 3) = P(X = 2, Y = 1) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$
$$P(W = 4) = P(X = 0, Y = 4) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$
$$P(W = 5) = P(X = 0, Y = 5) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$
$$P(W = 6) = P(X = 2, Y = 4) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$
$$P(W = 7) = P(X = 2, Y = 5) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8}.$$

That is, $f(w) = P(W = w) = \frac{1}{8}, \quad w \in S.$

(b)




Figure 2.1–10: Probability histogram of sum of two special dice

2.1-12 Let x equal the number of orange balls and $144 - x$ the number of blue balls. Then

$$\begin{aligned} \frac{x}{144} \cdot \frac{x-1}{143} + \frac{144-x}{144} \cdot \frac{143-x}{143} &= \frac{x}{144} \cdot \frac{144-x}{143} + \frac{144-x}{144} \cdot \frac{x}{143} \\ x^2 - x + 144 \cdot 143 - 144x - 143x + x^2 &= 2 \cdot 144x - 2 \cdot x^2 \\ x^2 - 144x + 5{,}148 &= 0 \\ (x - 78)(x - 66) &= 0 \end{aligned}$$

Thus there are 78 orange balls and 66 blue balls.

2.2 Mathematical Expectation

2.2-2 $E(X) = (-1)\left(\frac{4}{9}\right) + (0)\left(\frac{1}{9}\right) + (1)\left(\frac{4}{9}\right) = 0;$

$$E(X^2) = (-1)^2\left(\frac{4}{9}\right) + (0)^2\left(\frac{1}{9}\right) + (1)^2\left(\frac{4}{9}\right) = \frac{8}{9};$$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

$$E(3X^2 - 2X + 4) = 3\left(\frac{8}{9}\right) - 2(0) + 4 = \frac{20}{3}.$$

2.2-4
$$\begin{aligned} 1 &= \sum_{x=0}^{6} f(x) = \frac{9}{10} + c\left(\frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6}\right) \\ c &= \frac{2}{49}; \\ E(\text{Payment}) &= \frac{2}{49}\left(1 \cdot \frac{1}{2} + 2 \cdot \frac{1}{3} + 3 \cdot \frac{1}{4} + 4 \cdot \frac{1}{5} + 5 \cdot \frac{1}{6}\right) = \frac{71}{490} \text{ units.} \end{aligned}$$

2.2-6 Note that $\displaystyle\sum_{x=1}^{\infty} \frac{6}{\pi^2 x^2} = \frac{6}{\pi^2}\sum_{x=1}^{\infty} \frac{1}{x^2} = \frac{6}{\pi^2}\frac{\pi^2}{6} = 1$, so this is a pdf.

$$E(X) = \sum_{x=1}^{\infty} x\,\frac{6}{\pi^2 x^2} = \frac{6}{\pi^2}\sum_{x=1}^{\infty} \frac{1}{x} = +\infty$$

and it is well known that the sum of this harmonic series is not finite.

2.2-8 $E(|X - c|) = \dfrac{1}{7}\displaystyle\sum_{x \in S} |x - c|$, where $S = \{1, 2, 3, 5, 15, 25, 50\}$.

When $c = 5$,

$$E(|X - 5|) = \frac{1}{7}\left[(5-1) + (5-2) + (5-3) + (5-5) + (15-5) + (25-5) + (50-5)\right].$$

If c is either increased or decreased by 1, this expectation is increased by $1/7$. Thus $c = 5$, the median, minimizes this expectation while $b = E(X) = 101/7$ minimizes $E[(X - b)^2]$.

2.2-10 $(1) \cdot \dfrac{15}{36} + (-1) \cdot \dfrac{21}{36} = \dfrac{-6}{36} = \dfrac{-1}{6}$;

$(1) \cdot \dfrac{15}{36} + (-1) \cdot \dfrac{21}{36} = \dfrac{-6}{36} = \dfrac{-1}{6}$;

$(4) \cdot \dfrac{6}{36} + (-1) \cdot \dfrac{30}{36} = \dfrac{-6}{36} = \dfrac{-1}{6}$.

2.2-12 **(a)** The average class size is $\dfrac{(16)(25) + (3)(100) + (1)(300)}{20} = 50$;

(b)
$$f(x) = \begin{cases} 0.4, & x = 25, \\ 0.3, & x = 100, \\ 0.3, & x = 300; \end{cases}$$

(c) $E(X) = 25(0.4) + 100(0.3) + 300(0.3) = 130$.

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

2.3 Special Mathematical Expectations

2.3-2 **(a)**
$$\begin{aligned}\mu &= E(X)\\ &= \sum_{x=1}^{3} x\,\frac{3!}{x!\,(3-x)!}\left(\frac{1}{4}\right)^x\left(\frac{3}{4}\right)^{3-x}\\ &= 3\left(\frac{1}{4}\right)\sum_{k=0}^{2}\frac{2!}{k!\,(2-k)!}\left(\frac{1}{4}\right)^k\left(\frac{3}{4}\right)^{2-k}\\ &= 3\left(\frac{1}{4}\right)\left(\frac{1}{4}+\frac{3}{4}\right)^2=\frac{3}{4};\end{aligned}$$

$$\begin{aligned}E[X(X-1)] &= \sum_{x=2}^{3} x(x-1)\,\frac{3!}{x!\,(3-x)!}\left(\frac{1}{4}\right)^x\left(\frac{3}{4}\right)^{3-x}\\ &= 2(3)\left(\frac{1}{4}\right)^2\frac{3}{4}+6\left(\frac{1}{4}\right)^3\\ &= 6\left(\frac{1}{4}\right)^2=2\left(\frac{1}{4}\right)\left(\frac{3}{4}\right);\\ \sigma^2 &= E[X(X-1)]+E(X)-\mu^2\\ &= (2)\left(\frac{3}{4}\right)\left(\frac{1}{4}\right)+\left(\frac{3}{4}\right)-\left(\frac{3}{4}\right)^2\\ &= (2)\left(\frac{3}{4}\right)\left(\frac{1}{4}\right)+\left(\frac{3}{4}\right)\left(\frac{1}{4}\right)=3\left(\frac{1}{4}\right)\left(\frac{3}{4}\right);\end{aligned}$$

(b)
$$\begin{aligned}\mu &= E(X)\\ &= \sum_{x=1}^{4} x\,\frac{4!}{x!\,(4-x)!}\left(\frac{1}{2}\right)^x\left(\frac{1}{2}\right)^{4-x}\\ &= 4\left(\frac{1}{2}\right)\sum_{k=0}^{3}\frac{3!}{k!\,(3-k)!}\left(\frac{1}{2}\right)^k\left(\frac{1}{2}\right)^{3-k}\\ &= 4\left(\frac{1}{2}\right)\left(\frac{1}{2}+\frac{1}{2}\right)^3=2;\end{aligned}$$

$$\begin{aligned}E[X(X-1)] &= \sum_{x=2}^{4} x(x-1)\,\frac{4!}{x!\,(4-x)!}\left(\frac{1}{2}\right)^x\left(\frac{1}{2}\right)^{4-x}\\ &= 2(6)\left(\frac{1}{2}\right)^4+(6)(4)\left(\frac{1}{2}\right)^4+(12)\left(\frac{1}{2}\right)^4\\ &= 48\left(\frac{1}{2}\right)^4=12\left(\frac{1}{2}\right)^2;\\ \sigma^2 &= (12)\left(\frac{1}{2}\right)^2+\frac{4}{2}-\left(\frac{4}{2}\right)^2=1.\end{aligned}$$

2.3-4 $E[(X-\mu)/\sigma]=(1/\sigma)[E(X)-\mu]=(1/\sigma)(\mu-\mu)=0;$

$$E\{[(X-\mu)/\sigma]^2\}=(1/\sigma^2)E[(X-\mu)^2]=(1/\sigma^2)(\sigma^2)=1.$$

2.3-6 $f(1)=\frac{3}{8}$, $f(2)=\frac{2}{8}$, $f(3)=\frac{3}{8}$,

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

$$\mu = 1 \cdot \frac{3}{8} + 2 \cdot \frac{2}{8} + 3 \cdot \frac{3}{8} = 2,$$

$$\sigma^2 = 1^2 \cdot \frac{3}{8} + 2^2 \cdot \frac{2}{8} + 3^2 \cdot \frac{3}{8} - 2^2 = \frac{3}{4}.$$

2.3-8
$$\begin{aligned} E(X) &= \sum_{x=1}^{4} x \cdot \frac{2x-1}{16} \\ &= \frac{50}{16} = 3.125; \\ E(X^2) &= \sum_{x=1}^{4} x^2 \cdot \frac{2x-1}{16} \\ &= \frac{85}{8}; \\ \text{Var}(X) &= \frac{85}{8} - \left(\frac{25}{8}\right)^2 = \frac{55}{64} = 0.8594; \\ \sigma &= \frac{\sqrt{55}}{8} = 0.9270. \end{aligned}$$

2.3-10 $\mu = E(X) = (-5)(1/16) + (-1)(5/8) + (3)(5/16) = 0$ so
$E[(X-\mu)^3] = E(X^3) = (-5)^3(1/16)+(-1)^3(5/8)+(3^3)(5/16) = (-125-10+135)/16 = 0.$
Thus the index of skewness likewise equals 0. The distribution is not symmetric, however.

2.3-12 **(a)** $f(x) = \left(\frac{364}{365}\right)^{x-1}\left(\frac{1}{365}\right), \qquad x = 1, 2, 3, \ldots;$

(b)
$$\begin{aligned} \mu &= \frac{1}{\frac{1}{365}} = 365, \\ \sigma^2 &= \frac{\frac{364}{365}}{\left(\frac{1}{365}\right)^2} = 132{,}860, \\ \sigma &= 364.500; \end{aligned}$$

(c)
$$\begin{aligned} P(X > 400) &= \left(\frac{364}{365}\right)^{400} = 0.3337, \\ P(X < 300) &= 1 - \left(\frac{364}{365}\right)^{299} = 0.5597. \end{aligned}$$

2.3-14 $P(X \geq 100) = P(X > 99) = (0.99)^{99} = 0.3697.$

2.3-16 **(a)** $f(x) = (1/2)^{x-1}, \quad x = 2, 3, 4, \ldots;$

(b)
$$\begin{aligned} M(t) &= E[e^{tx}] = \sum_{x=2}^{\infty} e^{tx}(1/2)^{x-1} \\ &= 2\sum_{x=2}^{\infty}(e^t/2)^x \\ &= \frac{2(e^t/2)^2}{1 - e^t/2} = \frac{e^{2t}}{2 - e^t}, \qquad t < \ln 2; \end{aligned}$$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

(c) $M'(t) = \dfrac{4e^{2t} - e^{3t}}{(2-e^t)^2}$

$\mu = M'(0) = 3;$

$M''(t) = \dfrac{(2-e^t)^2(8e^{2t} - 3e^{3t}) - (4e^{2t} - e^{3t})2 * (2-e^t)(-e^t)}{(2-e^t)^4}$

$\sigma^2 = M''(0) - \mu^2 = 11 - 9 = 2;$

(d) **(i)** $P(X \leq 3) = 3/4$; **(ii)** $P(X \geq 5) = 1/8$; **(iii)** $P(X = 3) = 1/4$.

2.3-18 $P(X > k + j \,|\, X > k) = \dfrac{P(X > k+j)}{P(X > k)}$

$= \dfrac{q^{k+j}}{q^k} = q^j = P(X > j).$

2.3-20 **(a)** Construct a square in the first quadrant with lengths of sides equal to one. Within this square construct squares with one vertex at the origin and side lengths equal to $1 - 1/2^x$;

(b) $f(x) = \left(1 - \dfrac{1}{2^x}\right)^2 - \left(1 - \dfrac{1}{2^{x-1}}\right)^2 = \dfrac{(2^x-1)^2}{2^{2x}} - \dfrac{(2^{x-1}-1)^2}{2^{2x-2}}$

$= \dfrac{(2^x-1)^2}{2^{2x}} - \dfrac{2^2(2^{x-1}-1)^2}{2^{2x}} = \dfrac{(2^x-1)^2}{2^{2x}} - \dfrac{(2^x-2)^2}{2^{2x}}$

$= \dfrac{2^{2x} - 2\cdot 2^x + 1 - 2^{2x} + 2^2 \cdot 2^x - 4}{2^{2x}} = \dfrac{2^{x+1} - 3}{2^{2x}}, \qquad x = 1, 2, 3, \ldots;$

(c) $\displaystyle\sum_{x=1}^{\infty}\left[\frac{2^{x+1}}{2^{2x}} - \frac{3}{2^{2x}}\right] = \sum_{x=1}^{\infty}\left[\frac{2\cdot 2^x}{2^{2x}} - \frac{3}{4^x}\right] = \frac{4/4}{1-1/2} - \frac{3/4}{1-1/4} = 2 - 1 = 1;$

(d) $M(t) = E(e^{tX}) = \displaystyle\sum_{x=1}^{\infty} e^{tx}\cdot\frac{2^{x+1}-3}{2^{2x}} = \sum_{x=1}^{\infty}\left[\frac{2(2e^t)^x}{4^x} - \frac{3(e^{tx})}{4^x}\right] = \frac{2e^t}{2-e^t} - \frac{3e^t}{4-e^t};$

(e) $M'(t) = \dfrac{4e^t - 2e^{2t} + 2e^{2t}}{(2-e^t)^2} - \dfrac{12e^t - 3e^{2t} + 3e^{2t}}{(4-e^t)^2}$

$= \dfrac{4e^t}{(2-e^t)^2} - \dfrac{12e^t}{(4-e^t)^2}$

$\mu = M'(0) = 4 - \dfrac{12}{9} = \dfrac{8}{3};$

(f) $M''(t) = \dfrac{(2-e^t)^2\cdot 4e^t - 4e^t - 4e^t\cdot 2(2-e^t)(-e^t)}{(2-e^t)^4} -$

$\dfrac{(4-e^t)^2\cdot 12e^t - 12e^t - 12e^t\cdot 2(4-e^t)(-e^t)}{(4-e^t)^4}$

$\sigma^2 = M''(0) - \mu^2 = \dfrac{4+8}{1} - \dfrac{108+72}{81} - \left(\dfrac{8}{3}\right)^2 = \dfrac{8}{3}.$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

2.4 The Binomial Distribution

2.4-2 $f(-1) = \frac{11}{18}, \qquad f(1) = \frac{7}{18};$

$$\mu = (-1)\frac{11}{18} + (1)\frac{7}{18} = -\frac{4}{18};$$

$$\sigma^2 = \left(-1 + \frac{4}{18}\right)^2\left(\frac{11}{18}\right) + \left(1 + \frac{4}{18}\right)^2\left(\frac{7}{18}\right) = \frac{77}{81}.$$

The variance can also be computed by noting that $X^2 = 1$ so

$$\sigma^2 = 1 - \mu^2 = 1 - \left(-\frac{4}{18}\right)^2 = \frac{77}{81}.$$

2.4-4 **(a)** X is $b(7,\ 0.15)$;

(b) (i) $P(X \geq 2) = 1 - P(X \leq 1) = 1 - 0.7166 = 0.2834;$

(ii) $P(X = 1) = P(X \leq 1) - P(X \leq 0) = 0.7166 - 0.3206 = 0.3960;$

(iii) $P(X \leq 3) = 0.9879.$

2.4-6 **(a)** X is $b(15, 0.75)$; $\quad 15 - X$ is $b(15, 0.25)$;

(b) $P(X \geq 10) = P(15 - X \leq 5) = 0.8516;$

(c) $P(X \leq 10) = P(15 - X \geq 5) = 1 - P(15 - X \leq 4) = 1 - 0.6865 = 0.3135;$

(d) $P(X = 10) = P(X \geq 10) - P(X \geq 11)$

$= P(15 - X \leq 5) - P(15 - X \leq 4) = 0.8516 - 0.6865 = 0.1651;$

(e) $\mu = (15)(0.75) = 11.25, \quad \sigma^2 = (15)(0.75)(0.25) = 2.8125; \quad \sigma = \sqrt{2.8125} = 1.67705.$

2.4-8 **(a)** $1 - 0.01^4 = 0.99999999;$ **(b)** $0.99^4 = 0.960596.$

2.4-10 **(a)** X is $b(8,\ 0.90)$;

(b) (i) $P(X = 8) = P(8 - X = 0) = 0.4305;$

(ii) $P(X \leq 6) = P(8 - X \geq 2)$

$= 1 - P(8 - X \leq 1) = 1 - 0.8131 = 0.1869;$

(iii) $P(X \geq 6) = P(8 - X \leq 2) = 0.9619.$

2.4-12 **(a)**

$$f(x) = \begin{cases} 125/216, & x = -1, \\ 75/216, & x = 1, \\ 15/216, & x = 2, \\ 1/216, & x = 3; \end{cases}$$

(b) $\mu = (-1)\cdot\frac{125}{216} + (1)\cdot\frac{75}{216} + (2)\cdot\frac{15}{216} + (3)\cdot\frac{1}{216} = -\frac{17}{216};$

$\sigma^2 = E(X^2) - \mu^2 = \frac{269}{216} - \left(-\frac{17}{216}\right)^2 = 1.2392;$

$\sigma = 1.11;$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

(c) See Figure 2.4-12.




Figure 2.4–12: Losses in chuck-a-luck

2.4-14 Let X equal the number of winning tickets when n tickets are purchased. Then

$$\begin{aligned} P(X \geq 1) &= 1 - P(X = 0) \\ &= 1 - \left(\frac{9}{10}\right)^n. \end{aligned}$$

(a)
$$\begin{aligned} 1 - (0.9)^n &= 0.50 \\ (0.9)^n &= 0.50 \\ n \ln 0.9 &= \ln 0.5 \\ n &= \frac{\ln 0.5}{\ln 0.9} = 6.58 \end{aligned}$$

so $n = 7$;

(b)
$$\begin{aligned} 1 - (0.9)^n &= 0.95 \\ (0.9)^n &= 0.05 \\ n &= \frac{\ln 0.05}{\ln 0.9} = 28.43 \end{aligned}$$

so $n = 29$.

2.4-16 It is given that X is $b(10,\ 0.10)$. We are to find M so that $P(1000X \leq M) \geq 0.99)$ or $P(X \leq M/1000) \geq 0.99$. From Appendix Table II, $P(X \leq 4) = 0.9984 > 0.99$. Thus $M/1000 = 4$ or $M = 4000$ dollars.

2.4-18 Let X equal the number of infected individuals. X is $b(5,\ 0.05)$. The expected number of tests is

$1\,P(X = 0) + 6\,P(X > 0) = 1\,(0.7738) + 6\,(1 - 0.7738) = 2.131.$

2.4-20 **(a)** **(i)** $b(5,\ 0.7)$; **(ii)** $\mu = 3.5, \sigma^2 = 1.05$; **(iii)** 0.1607;
(b) **(i)** geometric, $p = 0.3$; **(ii)** $\mu = 10/3, \sigma^2 = 70/9$; **(iii)** 0.51;
(c) **(i)** Bernoulli, $p = 0.55$; **(ii)** $\mu = 0.55, \sigma^2 = 0.2475$; **(iii)** 0.55;

Copyright © 2020 Pearson Education, Inc.
Probability and Statistical Inference 10th edition Hogg Solutions Manual

(d) **(ii)** $\mu = 2.1, \sigma^2 = 0.89$; **(iii)** 0.7;

(e) **(i)** discrete uniform on $1, 2, \ldots, 10$; **(ii)** 5.5, 8.25; **(iii)** 0.2.

2.5 The Hypergeometric Distribution

2.5-2 **(a)** $\dfrac{\binom{4}{4}\binom{4}{0}}{\binom{8}{4}} = \dfrac{1}{70}$;

(b) $2\dfrac{\binom{4}{4}\binom{4}{0}}{\binom{8}{4}} = \dfrac{2}{70}$.

2.5-4 $P(X \geq 1) = 1 - P(X = 0) = 1 - \dfrac{\binom{3}{0}\binom{17}{5}}{\binom{20}{5}} = 1 - \dfrac{91}{228} = \dfrac{137}{228} = 0.60.$

2.5-6 We have $N = N_1 + N_2$. Thus

$$\begin{aligned} E[X(X-1)] &= \sum_{x=0}^{n} x(x-1)f(x) \\ &= \frac{\displaystyle\sum_{x=2}^{n} x(x-1)\,\frac{N_1!}{x!\,(N_1-x)!}\cdot\frac{N_2!}{(n-x)!\,(N_2-n+x)!}}{\binom{N}{n}} \\ &= N_1(N_1-1)\,\frac{\displaystyle\sum_{x=2}^{n}\frac{(N_1-2)!}{(x-2)!\,(N_1-x)!}\cdot\frac{N_2!}{(n-x)!\,(N_2-n+x)!}}{\binom{N}{n}}. \end{aligned}$$

In the summation, let $k = x - 2$, and in the denominator, note that

$$\binom{N}{n} = \frac{N!}{n!(N-n)!} = \frac{N(N-1)}{n(n-1)}\binom{N-2}{n-2}.$$

Thus

$$\begin{aligned} E[X(X-1)] &= \frac{N_1(N_1-1)}{\dfrac{N(N-1)}{n(n-1)}}\sum_{k=0}^{n-2}\frac{\binom{N_1-2}{k}\binom{N_2}{n-2-k}}{\binom{N-2}{n-2}} \\ &= \frac{N_1(N_1-1)(n)(n-1)}{N(N-1)}. \end{aligned}$$

2.5-8 **(a)** $\dfrac{\binom{5}{5}\binom{65}{0}}{\binom{70}{5}}\cdot\dfrac{1}{25} = \dfrac{1}{302{,}575{,}350}$;

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

(b) $\dfrac{\binom{5}{5}\binom{65}{0}}{\binom{70}{5}} \cdot \dfrac{24}{25} = \dfrac{24}{302{,}575{,}350};$

(c) $\dfrac{\binom{5}{4}\binom{65}{1}}{\binom{70}{5}} \cdot \dfrac{1}{25} = \dfrac{325}{302{,}575{,}350};$

(d) $\dfrac{\binom{5}{4}\binom{65}{1}}{\binom{70}{5}} \cdot \dfrac{24}{25} = \dfrac{7{,}800}{302{,}575{,}350};$

(e) $\dfrac{\binom{5}{0}\binom{65}{5}}{\binom{70}{5}} \cdot \dfrac{1}{25} = \dfrac{8{,}259{,}888}{302{,}575{,}350}.$

2.5-10 **(a)** $P(2,1,6,10)$ means that 2 is in position 1 so 1 cannot be selected. Thus

$$P(2,1,6,10) = \frac{\binom{1}{0}\binom{1}{1}\binom{8}{5}}{\binom{10}{6}} = \frac{56}{210} = \frac{4}{15};$$

(b) $P(i,r,k,n) = \dfrac{\binom{i-1}{r-1}\binom{1}{1}\binom{n-i}{k-r}}{\binom{n}{k}}.$

2.6 The Negative Binomial Distribution

2.6-2 $\binom{10-1}{5-1}\left(\frac{1}{2}\right)^5\left(\frac{1}{2}\right)^5 = \frac{126}{1024} = \frac{63}{512}.$

2.6-4 Let "being missed" be a success and let X equal the number of trials until the first success. Then $p = 0.01$.

$P(X \le 50) = 1 - 0.99^{50} = 1 - 0.605 = 0.395.$

2.6-6 **(a)**
$$\begin{aligned} R(t) &= \ln(1-p+pe^t), \\ R'(t) &= \left[\frac{pe^t}{1-p+pe^t}\right]_{t=0} = p, \\ R''(t) &= \left[\frac{(1-p+pe^t)(pe^t)-(pe^t)(pe^t)}{(1-p+pe^t)^2}\right]_{t=0} = p(1-p); \end{aligned}$$

(b)
$$\begin{aligned} R(t) &= n\ln(1-p+pe^t), \\ R'(t) &= \left[\frac{npe^t}{1-p+pe^t}\right]_{t=0} = np, \\ R''(t) &= n\left[\frac{(1-p+pe^t)(pe^t)-(pe^t)(pe^t)}{(1-p+pe^t)^2}\right]_{t=0} = np(1-p); \end{aligned}$$

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual

(c) $$\begin{aligned} R(t) &= \ln p + t - \ln[1-(1-p)e^t], \\ R'(t) &= \left[1 + \frac{(1-p)e^t}{1-(1-p)e^t}\right]_{t=0} = 1 + \frac{1-p}{p} = \frac{1}{p}, \\ R''(t) &= \left[(-1)\{1-(1-p)e^t\}^2\{-(1-p)e^t\}\right]_{t=0} = \frac{1-p}{p^2}; \end{aligned}$$

(d) $$\begin{aligned} R(t) &= r\left[\ln p + t - \ln\{1-(1-p)e^t\}\right], \\ R'(t) &= r\left[\frac{1}{1-(1-p)e^t}\right]_{t=0} = \frac{r}{p}, \\ R''(t) &= r\left[(-1)\{1-(1-p)e^t\}^{-2}\{-(1-p)e^t\}\right]_{t=0} = \frac{r(1-p)}{p^2}. \end{aligned}$$

2.6-8 $(0.7)(0.7)(0.3) = 0.147$.

2.6-10 **(a)** Let X equal the number of boxes that must be purchased. Then

$$E(X) = \sum_{x=1}^{10} \frac{1}{(11-x)/10} = 29.29;$$

(b) $\dfrac{100 \cdot E(X)}{365} \approx 8$ years.

2.7 The Poisson Distribution

2.7-2 $\lambda = \mu = \sigma^2 = 3$ so $P(X = 2) = 0.423 - 0.199 = 0.224$.

2.7-4 $$\begin{aligned} 3\frac{\lambda^1 e^{-\lambda}}{1!} &= \frac{\lambda^2 e^{-\lambda}}{2!} \\ e^{-\lambda}\lambda(\lambda - 6) &= 0 \\ \lambda &= 6 \end{aligned}$$

Thus $P(X = 4) = 0.285 - 0.151 = 0.134$.

2.7-6 $$\begin{aligned} M(t) &= e^{\lambda(e^t-1)}, \\ M'(t) &= e^{\lambda(e^t-1)}\lambda e^t, \\ M''(t) &= e^{\lambda(e^t-1)}\lambda^2 e^{2t} + e^{\lambda(e^t-1)}\lambda e^t, \\ M'''(t) &= e^{\lambda(e^t-1)}\lambda^3 e^{3t} + e^{\lambda(e^t-1)}\lambda^2 \cdot 2e^{2t} + e^{\lambda(e^t-1)}\lambda^2 e^{2t} + e^{\lambda(e^t-1)}\lambda e^t, \\ E(X^3) &= M'''(0) = \lambda^3 + 3\lambda^2 + \lambda, \\ \gamma &= [\lambda^3 + 3\lambda^2 + \lambda - \lambda^3 - 3\lambda(\lambda + \lambda^2) + 3\lambda^2 \cdot \lambda]/\lambda^{3/2} = 1/\sqrt{\lambda}. \end{aligned}$$

Thus the index of skewness is positive. As λ increases, the index of skewness deceases to zero.

2.7-8 $np = 1000(0.005) = 5$,

(a) $P(X \leq 1) \approx 0.040$;

(b) $P(X = 4, 5, 6) = P(X \leq 6) - P(X \leq 3) \approx 0.762 - 0.265 = 0.497$.

2.7-10 $\sigma = \sqrt{9} = 3$,

$P(3 < X < 15) = P(X \leq 14) - P(X \leq 3) = 0.959 - 0.021 = 0.938$.

2.7-12 Since $E(X) = 0.2$, the expected loss is $(0.2)(\$100{,}000) = \$2{,}000$.

Copyright © 2020 Pearson Education, Inc.

Probability and Statistical Inference 10th edition Hogg Solutions Manual