# *Algebra and Trigonometry 7th edition Blitzer Solutions Manual*

SKU: 9780136922179-SOLUTIONS

# INSTRUCTOR'S SOLUTIONS MANUAL

DANIEL S. MILLER

*Niagara County Community College*

# ALGEBRA AND TRIGONOMETRY


## SEVENTH EDITION

Robert Blitzer

*Miami Dade College*

Pearson

Algebra and Trigonometry 7th edition Blitzer Solutions Manual




The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expressed or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson from electronic files supplied by the author.

Copyright © 2022, 2018, 2014 by Pearson Education, Inc. 221 River Street, Hoboken, NJ 07030. All rights reserved.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.




ISBN-13: 978-0-13-739397-8
ISBN-10: 0-13-739397-0

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

TABLE OF CONTENTS for INSTRUCTOR SOLUTIONS

## *ALGEBRA AND TRIGONOMETRY 7E*


| | | |
|---|---|---|
| Chapter P | Fundamental Concepts of Algebra | 1 |
| Chapter 1 | Equations and Inequalities | 75 |
| Chapter 2 | Functions and Graphs | 215 |
| Chapter 3 | Polynomial and Rational Functions | 359 |
| Chapter 4 | Exponential and Logarithmic Functions | 515 |
| Chapter 5 | Trigonometric Functions | 593 |
| Chapter 6 | Analytic Trigonometry | 835 |
| Chapter 7 | Additional Topics in Trigonometry | 969 |
| Chapter 8 | Systems of Equations and Inequalities | 1117 |
| Chapter 9 | Matrices and Determinants | 1259 |
| Chapter 10 | Conic Sections and Analytic Geometry | 1377 |
| Chapter 11 | Sequences, Induction, and Probability | 1497 |




Algebra and Trigonometry 7th edition Blitzer Solutions Manual

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

# Chapter P
# Fundamental Concepts of Algebra

**Section P.1**

**Check Point Exercises**

**1.** $8+6(x-3)^2 = 8+6(13-3)^2$
$= 8+6(10)^2$
$= 8+6(100)$
$= 8+600$
$= 608$

**2. a.** Since 2016 is 16 years after 2000, substitute 16 for $x$.
$T = -x^2+361x+3193$
$= -(16)^2+361(16)+3193$
$= 8713$
The average cost of tuition and fees at public U.S. colleges for the school year ending in 2016 was \$8713.

**b.** The formula underestimates the actual answer by \$65.

**3.** The elements common to $\{3, 4, 5, 6, 7\}$ and $\{3, 7, 8, 9\}$ are 3 and 7.
$\{3,4,5,6,7\} \cap \{3,7,8,9\} = \{3,7\}$

**4.** The union is the set containing all the elements of either set.
$\{3,4,5,6,7\} \cup \{3,7,8,9\} = \{3,4,5,6,7,8,9\}$

**5.** $\left\{-9,\ -1.3,\ 0,\ 0.\overline{3},\ \frac{\pi}{2},\ \sqrt{9},\ \sqrt{10}\right\}$

**a.** Natural numbers: $\sqrt{9}$ because $\sqrt{9}=3$

**b.** Whole numbers: $0,\ \sqrt{9}$

**c.** Integers: $-9,\ 0,\ \sqrt{9}$

**d.** Rational numbers: $-9,\ -1.3,\ 0,\ 0.\overline{3},\ \sqrt{9}$

**e.** Irrational numbers: $\frac{\pi}{2},\ \sqrt{10}$

**f.** Real numbers: $-9,\ -1.3,\ 0,\ 0.\overline{3},\ \frac{\pi}{2},\ \sqrt{9},\ \sqrt{10}$

**6. a.** $|1-\sqrt{2}|$
Because $\sqrt{2} \approx 1.4$, the number inside the absolute value bars is negative. The absolute value of $x$ when $x < 0$ is $-x$. Thus,
$|1-\sqrt{2}| = -(1-\sqrt{2}) = \sqrt{2}-1$

**b.** $|\pi-3|$
Because $\pi \approx 3.14$, the number inside the absolute value bars is positive. The absolute value of a positive number is the number itself. Thus,
$|\pi-3| = \pi-3.$

**c.** $\frac{|x|}{x}$
Because $x > 0$, $|x| = x$.
Thus, $\frac{|x|}{x} = \frac{x}{x} = 1$

**7.** $|-4-(5)| = |-9| = 9$
The distance between –4 and 5 is 9.

**8.** $7(4x^2+3x)+2(5x^2+x)$
$= 7(4x^2+3x)+2(5x^2+x)$
$= 28x^2+21x+10x^2+2x$
$= 38x^2+23x$

**9.** $6+4[7-(x-2)]$
$= 6+4[7-x+2)]$
$= 6+4[9-x]$
$= 6+36-4x$
$= 42-4x$

**Concept and Vocabulary Check P.1**

**C1.** expression

**C2.** $b$ to the $n$th power; base; exponent

**C3.** formula; modeling; models

**C4.** intersection; $A \cap B$

**C5.** union; $A \cup B$

Copyright © 2022 Pearson Education, Inc


Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**C6.** natural

**C7.** whole

**C8.** integers

**C9.** rational

**C10.** irrational

**C11.** rational; irrational

**C12.** absolute value; $x$, $-x$

**C13.** $b+a$; $ba$

**C14.** $a+(b+c)$; $(ab)c$

**C15.** $ab+ac$

**C16.** 0; inverse; 0; identity

**C17.** inverse; 1; identity

**C18.** simplified

**C19.** $a$

**Exercise Set P.1**

**1.** $7+5(10)=7+50=57$

**2.** $8+6(5)=8+30=38$

**3.** $6(3)-8=18-8=10$

**4.** $8(3)-4=24-4=20$

**5.** $8^2+3(8)=64+24=88$

**6.** $6^2+5(6)=36+30=66$

**7.** $7^2-6(7)+3=49-42+3=7+3=10$

**8.** $8^2-7(8)+4=64-56+4=8+4=12$

**9.** $4+5(9-7)^3=4+5(2)^3$
$=4+5(8)=4+40=44$

**10.** $6+5(8-6)^3=6+5(2)^3$
$=6+5(8)$
$=6+40=46$

**11.** $8^2-3(8-2)=64-3(6)$
$=64-18=46$

**12.** $8^2-4(8-3)=64-4(5)=64-20=44$

**13.** $\dfrac{5(x+2)}{2x-14}=\dfrac{5(10+2)}{2(10)-14}$
$=\dfrac{5(12)}{6}$
$=5\cdot 2$
$=10$

**14.** $\dfrac{7(x-3)}{2x-16}=\dfrac{7(9-3)}{2(9)-16}=\dfrac{7(6)}{2}=7\cdot 3=21$

**15.** $\dfrac{2x+3y}{x+1}; x=-2, y=4$
$=\dfrac{2(-2)+3(4)}{-2+1}=\dfrac{-4+12}{-1}=\dfrac{8}{-1}=-8$

**16.** $\dfrac{2x+y}{xy-2x}$; $x=-2$ and $y=4$
$\dfrac{2(-2)+4}{(-2)(4)-2(-2)}=\dfrac{-4+4}{-8+4}=\dfrac{0}{4}=0$

**17.** $C=\dfrac{5}{9}(50-32)=\dfrac{5}{9}(18)=10$
$50^\circ$F is equivalent to $10^\circ$C.

**18.** $C=\dfrac{5}{9}(F-32)=\dfrac{5}{9}(86-32)=\dfrac{5}{9}(54)=30$
$86^\circ$F is equivalent to $30^\circ$C.

**19.** $h=4+60t-16t^2=4+60(2)-16(2)^2$
$=4+120-16(4)=4+120-64$
$=124-64=60$
Two seconds after it is kicked, the ball's height is 60 feet.


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**20.** $h = 4 + 60t - 16t^2$
$= 4 + 60(3) - 16(3)^2$
$= 4 + 180 - 16(9)$
$= 4 + 180 - 144$
$= 184 - 144 = 40$
Three seconds after it is kicked, the ball's height is 40 feet.

**21.** $\{1,2,3,4\} \cap \{2,4,5\} = \{2,4\}$

**22.** $\{1,3,7\} \cap \{2,3,8\} = \{3\}$

**23.** $\{s,e,t\} \cap \{t,e,s\} = \{s,e,t\}$

**24.** $\{r,e,a,l\} \cap \{l,e,a,r\} = \{r,e,a,l\}$

**25.** $\{1,3,5,7\} \cap \{2,4,6,8,10\} = \{\ \}$
The empty set is also denoted by $\varnothing$.

**26.** $\{1,3,5,7\} \cap \{-5,-3,-1\} = \{\ \}$ or $\varnothing$

**27.** $\{a,b,c,d\} \cap \varnothing = \varnothing$

**28.** $\{w,y,z\} \cap \varnothing = \varnothing$

**29.** $\{1,2,3,4\} \cup \{2,4,5\} = \{1,2,3,4,5\}$

**30.** $\{1,3,7,8\} \cup \{2,3,8\} = \{1,2,3,7,8\}$

**31.** $\{1,3,5,7\} \cup \{2,4,6,8,10\}$
$= \{1,2,3,4,5,6,7,8,10\}$

**32.** $\{0,1,3,5\} \cup \{2,4,6\} = \{0,1,2,3,4,5,6\}$

**33.** $\{a,e,i,o,u\} \cup \varnothing = \{a,e,i,o,u\}$

**34.** $\{e,m,p,t,y\} \cup \varnothing = \{e,m,p,t,y\}$

**35.** **a.** $\sqrt{100}$

**b.** $0, \sqrt{100}$

**c.** $-9, 0, \sqrt{100}$

**d.** $-9, -\frac{4}{5}, 0, 0.25, 9.2, \sqrt{100}$

**e.** $\sqrt{3}$

**f.** $-9, -\frac{4}{5}, 0, 0.25, \sqrt{3}, 9.2, \sqrt{100}$

**36.** **a.** $\sqrt{49}$

**b.** $0, \sqrt{49}$

**c.** $-7, 0, \sqrt{49}$

**d.** $-7, -0.\overline{6}, 0, \sqrt{49}$

**e.** $\sqrt{50}$

**f.** $-7, -0.\overline{6}, 0, \sqrt{49}, \sqrt{50}$

**37.** **a.** $\sqrt{64}$

**b.** $0, \sqrt{64}$

**c.** $-11, 0, \sqrt{64}$

**d.** $-11, -\frac{5}{6}, 0, 0.75, \sqrt{64}$

**e.** $\sqrt{5}, \pi$

**f.** $-11, -\frac{5}{6}, 0, 0.75, \sqrt{5}, \pi, \sqrt{64}$

**38.** **a.** $\sqrt{4}$

**b.** $0, \sqrt{4}$

**c.** $-5, 0, \sqrt{4}$

**d.** $-5, -0.\overline{3}, 0, \sqrt{4}$

**e.** $\sqrt{2}$

**f.** $-5, -0.\overline{3}, 0, \sqrt{2}, \sqrt{4}$

**39.** 0

**40.** Answers will vary. An example is $\frac{1}{2}$.

**41.** Answers will vary. An example is 2.

**42.** Answers will vary. An example is $-2$.

**43.** true; $-13$ is to the left of $-2$ on the number line.

**44.** false; $-6$ is to the left of 2 on the number line.

**45.** true; 4 is to the right of $-7$ on the number line.

Copyright © 2022 Pearson Education, Inc.


Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**46.** true; –13 is to the left of –5 on the number line.

**47.** true; $-\pi = -\pi$

**48.** true; –3 is to the right of –13 on the number line.

**49.** true; 0 is to the right of –6 on the number line.

**50.** true; 0 is to the right of –13 on the number line.

**51.** $|300| = 300$

**52.** $|-203| = 203$

**53.** $|12-\pi| = 12-\pi$

**54.** $|7-\pi| = 7-\pi$

**55.** $\left|\sqrt{2}-5\right| = 5-\sqrt{2}$

**56.** $\left|\sqrt{5}-13\right| = 13-\sqrt{5}$

**57.** $\dfrac{-3}{|-3|} = \dfrac{-3}{3} = -1$

**58.** $\dfrac{-7}{|-7|} = \dfrac{-7}{7} = -1$

**59.** $\big||-3|-|-7|\big| = |3-7| = |-4| = 4$

**60.** $\big||-5|-|-13|\big| = |5-13| = |-8| = 8$

**61.** $|x+y| = |2+(-5)| = |-3| = 3$

**62.** $|x-y| = |2-(-5)| = |7| = 7$

**63.** $|x|+|y| = |2|+|-5| = 2+5 = 7$

**64.** $|x|-|y| = |2|-|-5| = 2-5 = -3$

**65.** $\dfrac{y}{|y|} = \dfrac{-5}{|-5|} = \dfrac{-5}{5} = -1$

**66.** $\dfrac{|x|}{x}+\dfrac{|y|}{y} = \dfrac{|2|}{2}+\dfrac{|-5|}{-5} = \dfrac{2}{2}+\dfrac{5}{-5} = 1+(-1) = 0$

**67.** The distance is $|2-17| = |-15| = 15$.

**68.** The distance is $|4-15| = |-11| = 11$.

**69.** The distance is $|-2-5| = |-7| = 7$.

**70.** The distance is $|-6-8| = |-14| = 14$.

**71.** The distance is $|-19-(-4)| = |-19+4| = |-15| = 15$.

**72.** The distance is $|-26-(-3)| = |-26+3| = |-23| = 23$.

**73.** The distance is
$|-3.6-(-1.4)| = |-3.6+1.4| = |-2.2| = 2.2$.

**74.** The distance is
$|-5.4-(-1.2)| = |-5.4+1.2| = |-4.2| = 4.2$.

**75.** $6 + (-4) = (-4) + 6$;
commutative property of addition

**76.** $11\cdot(7+4) = 11\cdot 7+11\cdot 4$;
distributive property of multiplication over addition

**77.** $6 + (2 + 7) = (6 + 2) + 7$;
associative property of addition

**78.** $6\cdot(2\cdot 3) = 6\cdot(3\cdot 2)$;
commutative property of multiplication

**79.** $(2 + 3) + (4 + 5) = (4 + 5) + (2 + 3)$;
commutative property of addition

**80.** $7\cdot(11\cdot 8) = (11\cdot 8)\cdot 7$;
commutative property of multiplication

**81.** $2(-8 + 6) = -16 + 12$;
distributive property of multiplication over addition

**82.** $-8(3+11) = -24+(-88)$;
distributive property of multiplication over addition

**83.** $\dfrac{1}{x+3}(x+3) = 1; x \neq -3$,
inverse property of multiplication

**84.** $(x+4)+\big[-(x+4)\big] = 0$;
inverse property of addition

**85.** $\begin{aligned} 5(3x+4)-4 &= 5\cdot 3x+5\cdot 4-4 \\ &= 15x+20-4 \\ &= 15x+16 \end{aligned}$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**86.** $2(5x+4)-3=2\cdot 5x+2\cdot 4-3$
$=10x+8-3$
$=10x+5$

**87.** $5(3x-2)+12x=5\cdot 3x-5\cdot 2+12x$
$=15x-10+12x$
$=27x-10$

**88.** $2(5x-1)+14x=2\cdot 5x-2\cdot 1+14x$
$=10x-2+14x$
$=24x-2$

**89.** $7(3y-5)+2(4y+3)$
$=7\cdot 3y-7\cdot 5+2\cdot 4y+2\cdot 3$
$=21y-35+8y+6$
$=29y-29$

**90.** $4(2y-6)+3(5y+10)$
$=4\cdot 2y-4\cdot 6+3\cdot 5y+3\cdot 10$
$=8y-24+15y+30$
$=23y+6$

**91.** $5(3y-2)-(7y+2)=15y-10-7y-2$
$=8y-12$

**92.** $4(5y-3)-(6y+3)=20y-12-6y-3$
$=14y-15$

**93.** $7-4\left[3-(4y-5)\right]=7-4[3-4y+5]$
$=7-4[8-4y]$
$=7-32+16y$
$=16y-25$

**94.** $6-5[8-(2y-4)]=6-5[8-2y+4]$
$=6-5[12-2y]$
$=6-60+10y$
$=10y-54$

**95.** $18x^2+4-\left[6\left(x^2-2\right)+5\right]$
$=18x^2+4-\left[6x^2-12+5\right]$
$=18x^2+4-\left[6x^2-7\right]$
$=18x^2+4-6x^2+7$
$=18x^2-6x^2+4+7$
$=(18-6)x^2+11=12x^2+11$

**96.** $14x^2+5-\left[7\left(x^2-2\right)+4\right]$
$=14x^2+5-\left[7x^2-14+4\right]$
$=14x^2+5-\left[7x^2-10\right]$
$=14x^2+5-7x^2+10$
$=14x^2-7x^2+5+10$
$=(14-7)x^2+15$
$=7x^2+15$

**97.** $-(-14x)=14x$

**98.** $-(-17y)=17y$

**99.** $-(2x-3y-6)=-2x+3y+6$

**100.** $-(5x-13y-1)=-5x+13y+1$

**101.** $\frac{1}{3}(3x)+[(4y)+(-4y)]=x+0=x$

**102.** $\frac{1}{2}(2y)+[(-7x)+7x]=y+0=y$

**103.** $|-6|\ \square\ |-3|$
$6\ \square\ 3$
$6>3$
Since $6>3$, $|-6|>|-3|$.

**104.** $|-20|\ \square\ |-50|$
$20\ \square\ 50$
$20<50$
Since $20<50$, $|-20|<|-50|$.

**105.** $\left|\frac{3}{5}\right|\ \square\ |-0.6|$
$|0.6|\ \square\ |-0.6|$
$0.6\ \square\ 0.6$
$0.6\ =\ 0.6$
Since $0.6=0.6$, $\left|\frac{3}{5}\right|=|-0.6|$.

Copyright © 2022 Pearson Education, Inc.


Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**106.** $\left|\frac{5}{2}\right| \square |-2.5|$

$|2.5| \square |-2.5|$

$2.5 \square 2.5$

$2.5 = 2.5$

Since $2.5 = 2.5$, $\left|\frac{5}{2}\right| = |-2.5|$.

**107.** $\frac{30}{40} - \frac{3}{4} \square \frac{14}{15} \cdot \frac{15}{14}$

$\frac{30}{40} - \frac{30}{40} \square \frac{\cancel{14}}{\cancel{15}} \cdot \frac{\cancel{15}}{\cancel{14}}$

$0 \square 1$

$0 < 1$

Since $0 < 1$, $\frac{30}{40} - \frac{3}{4} < \frac{14}{15} \cdot \frac{15}{14}$.

**108.** $\frac{17}{18} \cdot \frac{18}{17} \square \frac{50}{60} - \frac{5}{6}$

$\frac{\cancel{17}}{\cancel{18}} \cdot \frac{\cancel{18}}{\cancel{17}} \square \frac{50}{60} - \frac{50}{60}$

$1 \square 0$

$1 > 0$

Since $1 > 0$, $\frac{17}{18} \cdot \frac{18}{17} > \frac{50}{60} - \frac{5}{6}$.

**109.** $\frac{8}{13} \div \frac{8}{13} \square |-1|$

$\frac{8}{13} \cdot \frac{13}{8} \square 1$

$1 \square 1$

$1 = 1$

Since $1 = 1$, $\frac{8}{13} \div \frac{8}{13} = |-1|$.

**110.** $|-2| \square \frac{4}{17} \div \frac{4}{17}$

$2 \square \frac{4}{17} \cdot \frac{17}{4}$

$2 \square 1$

$2 > 1$

Since $2 > 1$, $|-2| > \frac{4}{17} \div \frac{4}{17}$.

**111.** $8^2 - 16 \div 2^2 \cdot 4 - 3 = 64 - 16 \div 4 \cdot 4 - 3$

$= 64 - 4 \cdot 4 - 3$

$= 64 - 16 - 3$

$= 48 - 3$

$= 45$

**112.** $10^2 - 100 \div 5^2 \cdot 2 - 3 = 100 - 100 \div 25 \cdot 2 - 3$

$= 100 - 4 \cdot 2 - 3$

$= 100 - 8 - 3$

$= 92 - 3$

$= 89$

**113.** $\frac{5 \cdot 2 - 3^2}{[3^2 - (-2)]^2} = \frac{5 \cdot 2 - 9}{[9 - (-2)]^2}$

$= \frac{10 - 9}{[9 + 2]^2}$

$= \frac{10 - 9}{11^2}$

$= \frac{1}{121}$

**114.** $\frac{10 \div 2 + 3 \cdot 4}{(12 - 3 \cdot 2)^2} = \frac{5 + 12}{(12 - 6)^2}$

$= \frac{17}{6^2}$

$= \frac{17}{36}$

**115.** $8 - 3[-2(2 - 5) - 4(8 - 6)] = 8 - 3[-2(-3) - 4(2)]$

$= 8 - 3[6 - 8]$

$= 8 - 3[-2]$

$= 8 + 6$

$= 14$

**116.** $8 - 3[-2(5 - 7) - 5(4 - 2)] = 8 - 3[-2(-2) - 5(2)]$

$= 8 - 3[4 - 10]$

$= 8 - 3[-6]$

$= 8 + 18$

$= 26$

**117.** $\frac{2(-2) - 4(-3)}{5 - 8} = \frac{-4 + 12}{-3}$

$= \frac{8}{-3}$

$= -\frac{8}{3}$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**118.** $\dfrac{6(-4)-5(-3)}{9-10}=\dfrac{-24+15}{-1}$

$=\dfrac{-9}{-1}$

$=9$

**119.** $\dfrac{(5-6)^2-2|3-7|}{89-3\cdot 5^2}=\dfrac{(-1)^2-2|-4|}{89-3\cdot 25}$

$=\dfrac{1-2(4)}{89-75}$

$=\dfrac{1-8}{14}$

$=\dfrac{-7}{14}$

$=-\dfrac{1}{2}$

**120.** $\dfrac{12\div 3\cdot 5\left|2^2+3^2\right|}{7+3-6^2}=\dfrac{12\div 3\cdot 5|4+9|}{7+3-36}$

$=\dfrac{4\cdot 5|13|}{10-36}$

$=\dfrac{20(13)}{-26}$

$=\dfrac{260}{-26}$

$=-10$

**121.** $x-(x+4)=x-x-4=-4$

**122.** $x-(8-x)=x-8+x=2x-8$

**123.** $6(-5x)=-30x$

**124.** $10(-4x)=-40x$

**125.** $5x-2x=3x$

**126.** $6x-(-2x)=6x+2x=8x$

**127.** $8x-(3x+6)=8x-3x-6=5x-6$

**128.** $8-3(x+6)=8-3x-18=-3x-10$

**129.** **a.** $H=\dfrac{7}{10}(220-a)$

$H=\dfrac{7}{10}(220-20)$

$=\dfrac{7}{10}(200)$

$=140$

The lower limit of the heart rate for a 20-year-old with this exercise goal is 140 beats per minute.

**b.** $H=\dfrac{4}{5}(220-a)$

$H=\dfrac{4}{5}(220-20)$

$=\dfrac{4}{5}(200)$

$=160$

The upper limit of the heart rate for a 20-year-old with this exercise goal is 160 beats per minute.

**130.** **a.** $H=\dfrac{1}{2}(220-a)$

$H=\dfrac{1}{2}(220-30)$

$=\dfrac{1}{2}(190)$

$=95$

The lower limit of the heart rate for a 30-year-old with this exercise goal is 95 beats per minute.

**b.** $H=\dfrac{3}{5}(220-a)$

$H=\dfrac{3}{5}(220-30)$

$=\dfrac{3}{5}(190)$

$=114$

The upper limit of the heart rate for a 30-year-old with this exercise goal is 114 beats per minute.

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**131. a.** Model 1

$T = 975x + 13,547$

$= 975(18) + 13,547$

$= 31,097$

Model 1 estimates the cost to have been \$31,097 in 2018, which overestimates the actual cost by \$366.

Model 2

$T = 32x^2 + 331x + 15,479$

$= 32(18)^2 + 331(18) + 15,479$

$= 31,805$

Model 2 estimates the cost to have been \$31,805 in 2018, which overestimates the actual cost by \$1074.

**b.** Model 2

$T = 32x^2 + 331x + 15,479$

$= 32(30)^2 + 331(30) + 15,479$

$= 54,209$

Model 2 estimates the cost will be \$54,209 in 2030.

**132. a.** Model 1

$T = 975x + 13,547$

$= 975(10) + 13,547$

$= 23,297$

Model 1 estimates the cost to have been \$23,297 in 2010, which overestimates the actual cost by \$1028.

Model 2

$T = 32x^2 + 331x + 15,479$

$= 32(10)^2 + 331(10) + 15,479$

$= 21,989$

Model 2 estimates the cost to have been \$21,989 in 2010, which underestimates the actual cost by \$280.

**b.** Model 2

$T = 32x^2 + 331x + 15,479$

$= 32(25)^2 + 331(25) + 15,479$

$= 43,754$

Model 2 estimates the cost will be \$43,754 in 2025.

**133. a.** $0.015x + 0.0175(6000 - x)$

$= 0.015x + 105 - 0.0175x$

$= 105 - 0.0025x$

**b.** $0.015x + 0.0175(6000 - x)$

$= 0.015(4400) + 0.0175(6000 - 4400)$

$= 0.015(4400) + 0.0175(1600)$

$= 94$

$105 - 0.0025x$

$= 105 - 0.0025(4400)$

$= 94$

Both expressions determine the total interest is \$94

**c.** $105 - 0.0025x$

$= 105 - 0.0025(3000)$

$= 97.5$

The total interest is \$97.50

**134. a.** $0.06t + 0.5(50 - t)$

$= 0.06t + 25 - 0.5t$

$= 25 - 0.44t$

**b.** $0.06t + 0.5(50 - t)$

$= 0.06t + 0.5(50 - 20)$

$= 0.06(20) + 0.5(30)$

$= 16.2$

$25 - 0.44t$

$= 25 - 0.44(20)$

$= 16.2$

Both expressions determine the total distance is 16.2 miles.

**c.** $25 - 0.44t$

$= 25 - 0.44(25)$

$= 14$

The total distance is 14 miles

**135. – 143.** Answers will vary.

**144.** does not make sense; Explanations will vary. Sample explanation: Models do not always accurately predict future values.

**145.** does not make sense; Explanations will vary. Sample explanation: To use the model, substitute 0 for $x$.


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**146.** makes sense

**147.** does not make sense; Explanations will vary. Sample explanation: The commutative property changes order and the associative property changes groupings.

**148.** false; Changes to make the statement true will vary. A sample change is: Some rational numbers are not integers.

**149.** false; Changes to make the statement true will vary. A sample change is: All whole numbers are integers.

**150.** true

**151.** false; Changes to make the statement true will vary. A sample change is: Some irrational numbers are negative.

**152.** false; Changes to make the statement true will vary. A sample change is: The term $x$ has a coefficient of 1.

**153.** false; Changes to make the statement true will vary. A sample change is: $5+3(x-4)=5+3x-12=3x-7.$

**154.** false; Changes to make the statement true will vary. A sample change is: $-x-x=-2x.$

**155.** true

**156.** $\sqrt{2}\approx 1.4$

$1.4<1.5$

$\sqrt{2}<1.5$

**157.** $-\pi > -3.5$

**158.** $-\dfrac{3.14}{2}=-1.57$

$-\dfrac{\pi}{2}\approx -1.571$

$-1.57 > -1.571$

$-\dfrac{3.14}{2} > -\dfrac{\pi}{2}$

**159.** **a.** $b^4\cdot b^3=(b\cdot b\cdot b\cdot b)(b\cdot b\cdot b)=b^7$

**b.** $b^5\cdot b^5=(b\cdot b\cdot b\cdot b\cdot b)(b\cdot b\cdot b\cdot b\cdot b)=b^{10}$

**c.** add the exponents

**160.** **a.** $\dfrac{b^7}{b^3}=\dfrac{b\cdot b\cdot b\cdot b\cdot b\cdot b\cdot b}{b\cdot b\cdot b}=b^4$

**b.** $\dfrac{b^8}{b^2}=\dfrac{b\cdot b\cdot b\cdot b\cdot b\cdot b\cdot b\cdot b}{b\cdot b}=b^6$

**c.** subtract the exponents

**161.** $6.2\times 10^3=6.2\times 10\times 10\times 10=6200$

It moves the decimal point 3 places to the right.

**Section P.2**

**Check Point Exercises**

**1.** **a.** $3^3\cdot 3^2=3^{3+2}=3^5$ or 243

**b.** $\left(4x^3y^4\right)\left(10x^2y^6\right)=4\cdot 10\cdot x^3\cdot x^2\cdot y^4\cdot y^6$

$=40x^{3+2}\cdot y^{4+6}$

$=40x^5y^{10}$

**2.** **a.** $\dfrac{(-3)^6}{(-3)^3}=(-3)^3=-27$

**b.** $\dfrac{27x^{14}y^8}{3x^3y^5}=\dfrac{27}{3}\cdot\dfrac{x^{14}}{x^3}\cdot\dfrac{y^8}{y^5}=9x^{14-3}y^{8-5}=9x^{11}y^3$

**3.** **a.** $5^{-2}=\dfrac{1}{5^2}=\dfrac{1}{25}$

**b.** $(-3)^{-3}=\dfrac{1}{(-3)^3}=\dfrac{1}{-27}=-\dfrac{1}{27}$

**c.** $\dfrac{1}{4^{-2}}=\dfrac{1}{\frac{1}{4^2}}=1\cdot\dfrac{4^2}{1}=4^2=16$

**d.** $3x^{-6}y^4=3\cdot\dfrac{1}{x^6}\cdot y^4=\dfrac{3y^4}{x^6}$

**4.** **a.** $\left(3^3\right)^2=3^{3\cdot 2}=3^6$ or 729

**b.** $\left(y^7\right)^{-2}=y^{7(-2)}=y^{-14}=\dfrac{1}{y^{14}}$

**c.** $\left(b^{-3}\right)^{-4}=b^{-3(-4)}=b^{12}$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**5.** $(-4x)^3 = (-4)^3(x)^3 = -64x^3$

**6. a.** $\left(-\frac{2}{y}\right)^5 = \frac{(-2)^5}{y^5} = \frac{-32}{y^5}$

**b.** $\left(\frac{x^5}{3}\right)^3 = \frac{\left(x^5\right)^3}{3^3} = \frac{x^{15}}{27}$

**7. a.** $\left(2x^3y^6\right)^4 = (2)^4\left(x^3\right)^4\left(y^6\right)^4 = 16x^{12}y^{24}$

**b.** $\left(-6x^2y^5\right)\left(3xy^3\right) = (-6)\cdot 3\cdot x^2\cdot x\cdot y^5\cdot y^3$
$= -18x^3y^8$

**c.** $\frac{100x^{12}y^2}{20x^{16}y^{-4}} = \left(\frac{100}{20}\right)\left(\frac{x^{12}}{x^{16}}\right)\left(\frac{y^2}{y^{-4}}\right)$
$= 5x^{12-16}y^{2-(-4)}$
$= 5x^{-4}y^6$
$= \frac{5y^6}{x^4}$

**d.** $\left(\frac{5x}{y^4}\right)^{-2} = \frac{(5)^{-2}(x)^{-2}}{\left(y^4\right)^{-2}}$
$= \frac{(5)^{-2}(x)^{-2}}{\left(y^4\right)^{-2}}$
$= \frac{5^{-2}x^{-2}}{y^{-8}}$
$= \frac{y^8}{5^2x^2}$
$= \frac{y^8}{25x^2}$

**8. a.** $-2.6\times10^9 = -2,600,000,000$

**b.** $3.017\times10^{-6} = 0.000003017$

**9. a.** $5,210,000,000 = 5.21\times10^9$

**b.** $-0.00000006893 = -6.893\times10^{-8}$

**10.** $410\times10^7 = \left(4.1\times10^2\right)\times10^7$
$= 4.1\times\left(10^2\times10^7\right)$
$= 4.1\times10^9$

**11 a.** $\left(7.1\times10^5\right)\left(5\times10^{-7}\right)$
$= 7.1\cdot5\times10^5\cdot10^{-7}$
$= 35.5\times10^{-2}$
$= \left(3.55\times10^1\right)\times10^{-2}$
$= 3.55\times\left(10^1\times10^{-2}\right)$
$= 3.55\times10^{-1}$

**b.** $\frac{1.2\times10^6}{3\times10^{-3}} = \frac{1.2}{3}\cdot\frac{10^6}{10^{-3}}$
$= 0.4\times10^{6-(-3)}$
$= 0.4\times10^9$
$= 4\times10^8$

**12.** $\frac{13,500,000,000,000}{309,000,000} = \frac{1.35\times10^{13}}{3.09\times10^8} = \frac{1.35}{3.09}\cdot\frac{10^{13}}{10^8}$
$= 0.44\times10^5$
$= 44,000$

Each citizen would have to pay $44,000 in 2010. This would be an increase of $33,000, per citizen, between 2010 and 2020.

**Concept and Vocabulary Check P.2**

**C1.** $b^{m+n}$; add

**C2.** $b^{m-n}$; subtract

**C3.** 1

**C4.** $\frac{1}{b^n}$

**C5.** false

**C6.** $b^n$

**C7.** true

**C8.** a number greater than or equal to 1 and less than 10; integer


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**C9.** true

**C10.** false

**Exercise Set P.2**

**1.** $5^2 \cdot 2 = (5 \cdot 5) \cdot 2 = 25 \cdot 2 = 50$

**2.** $6^2 \cdot 2 = (6 \cdot 6) \cdot 2 = 36 \cdot 2 = 72$

**3.** $(-2)^6 = (-2)(-2)(-2)(-2)(-2)(-2) = 64$

**4.** $(-2)^4 = (-2)(-2)(-2)(-2) = 16$

**5.** $-2^6 = -2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 = -64$

**6.** $-2^4 = -2 \cdot 2 \cdot 2 \cdot 2 = -16$

**7.** $(-3)^0 = 1$

**8.** $(-9)^0 = 1$

**9.** $-3^0 = -1$

**10.** $-9^0 = -1$

**11.** $4^{-3} = \frac{1}{4^3} = \frac{1}{4 \cdot 4 \cdot 4} = \frac{1}{64}$

**12.** $2^{-6} = \frac{1}{2^6} = \frac{1}{2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2} = \frac{1}{64}$

**13.** $2^2 \cdot 2^3 = 2^{2+3} = 2^5 = 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 = 32$

**14.** $3^3 \cdot 3^2 = 3^{3+2} = 3^5 = 3 \cdot 3 \cdot 3 \cdot 3 \cdot 3 = 243$

**15.** $(2^2)^3 = 2^{2 \cdot 3} = 2^6 = 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 = 64$

**16.** $(3^3)^2 = 3^{3 \cdot 2} = 3^6 = 3 \cdot 3 \cdot 3 \cdot 3 \cdot 3 \cdot 3 = 729$

**17.** $\frac{2^8}{2^4} = 2^{8-4} = 2^4 = 2 \cdot 2 \cdot 2 \cdot 2 = 16$

**18.** $\frac{3^8}{3^4} = 3^{8-4} = 3^4 = 3 \cdot 3 \cdot 3 \cdot 3 = 81$

**19.** $3^{-3} \cdot 3 = 3^{-3+1} = 3^{-2} = \frac{1}{3^2} = \frac{1}{3 \cdot 3} = \frac{1}{9}$

**20.** $2^{-3} \cdot 2 = 2^{-3+1} = 2^{-2} = \frac{1}{2^2} = \frac{1}{2 \cdot 2} = \frac{1}{4}$

**21.** $\frac{2^3}{2^7} = 2^{3-7} = 2^{-4} = \frac{1}{2^4} = \frac{1}{2 \cdot 2 \cdot 2 \cdot 2} = \frac{1}{16}$

**22.** $\frac{3^4}{3^7} = 3^{4-7} = 3^{-3} = \frac{1}{3^3} = \frac{1}{3 \cdot 3 \cdot 3} = \frac{1}{27}$

**23.** $x^{-2}y = \frac{1}{x^2} \cdot y = \frac{y}{x^2}$

**24.** $xy^{-3} = x \cdot \frac{1}{y^3} = \frac{x}{y^3}$

**25.** $x^0 y^5 = 1 \cdot y^5 = y^5$

**26.** $x^7 \cdot y^0 = x^7 \cdot 1 = x^7$

**27.** $x^3 \cdot x^7 = x^{3+7} = x^{10}$

**28.** $x^{11} \cdot x^5 = x^{11+5} = x^{16}$

**29.** $x^{-5} \cdot x^{10} = x^{-5+10} = x^5$

**30.** $x^{-6} \cdot x^{12} = x^{-6+12} = x^6$

**31.** $(x^3)^7 = x^{3 \cdot 7} = x^{21}$

**32.** $(x^{11})^5 = x^{11 \cdot 5} = x^{55}$

**33.** $(x^{-5})^3 = x^{-5 \cdot 3} = x^{-15} = \frac{1}{x^{15}}$

**34.** $(x^{-6})^4 = x^{-6 \cdot 4} = x^{-24} = \frac{1}{x^{24}}$

**35.** $\frac{x^{14}}{x^7} = x^{14-7} = x^7$

**36.** $\frac{x^{30}}{x^{10}} = x^{30-10} = x^{20}$

**37.** $\frac{x^{14}}{x^{-7}} = x^{14-(-7)} = x^{14+7} = x^{21}$

**38.** $\frac{x^{30}}{x^{-10}} = x^{30-(-10)} = x^{30+10} = x^{40}$

**39.** $(8x^3)^2 = 8^2 (x^3)^2 = 8^2 x^{3 \cdot 2} = 64x^6$

**40.** $(6x^4)^2 = (6)^2 (x^4)^2 = 6^2 x^{4 \cdot 2} = 36x^8$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**41.** $\left(-\frac{4}{x}\right)^3 = \frac{(-4)^3}{x^3} = -\frac{64}{x^3}$

**42.** $\left(-\frac{6}{y}\right)^3 = \frac{(-6)^3}{y^3} = -\frac{216}{y^3}$

**43.** $(-3x^2y^5)^2 = (-3)^2(x^2)^2 \cdot (y^5)^2$
$= 9x^{2\cdot 2}y^{5\cdot 2}$
$= 9x^4y^{10}$

**44.** $(-3x^4y^6)^3 = (-3)^3(x^4)^3(y^6)^3$
$= -27x^{4\cdot 3}y^{6\cdot 3}$
$= -27x^{12}y^{18}$

**45.** $(3x^4)(2x^7) = 3 \cdot 2x^4 \cdot x^7 = 6x^{4+7} = 6x^{11}$

**46.** $(11x^5)(9x^{12}) = 11 \cdot 9x^5x^{12} = 99x^{5+12} = 99x^{17}$

**47.** $(-9x^3y)(-2x^6y^4) = (-9)(-2)x^3x^6yy^4$
$= 18x^{3+6}y^{1+4}$
$= 18x^9y^5$

**48.** $(-5x^4y)(-6x^7y^{11}) = (-5)(-6)x^4x^7yy^{11}$
$= 30x^{4+7}y^{1+11}$
$= 30x^{11}y^{12}$

**49.** $\frac{8x^{20}}{2x^4} = \left(\frac{8}{2}\right)\left(\frac{x^{20}}{x^4}\right) = 4x^{20-4} = 4x^{16}$

**50.** $\frac{20x^{24}}{10x^6} = \left(\frac{20}{10}\right)\left(\frac{x^{24}}{x^6}\right) = 2x^{24-6} = 2x^{18}$

**51.** $\frac{25a^{13} \cdot b^4}{-5a^2 \cdot b^3} = \left(\frac{25}{-5}\right)\left(\frac{a^{13}}{a^2}\right)\left(\frac{b^4}{b^3}\right)$
$= -5a^{13-2}b^{4-3}$
$= -5a^{11}b$

**52.** $\frac{35a^{14}b^6}{-7a^7b^3} = \left(\frac{35}{-7}\right)\left(\frac{a^{14}}{a^7}\right)\left(\frac{b^6}{b^3}\right)$
$= -5a^{14-7}b^{6-3}$
$= -5a^7b^3$

**53.** $\frac{14b^7}{7b^{14}} = \left(\frac{14}{7}\right)\left(\frac{b^7}{b^{14}}\right) = 2 \cdot b^{7-14} = 2b^{-7} = \frac{2}{b^7}$

**54.** $\frac{20b^{10}}{10b^{20}} = \left(\frac{20}{10}\right)\left(\frac{b^{10}}{b^{20}}\right)$
$= 2b^{10-20}$
$= 2b^{-10}$
$= \frac{2}{b^{10}}$

**55.** $(4x^3)^{-2} = (4^{-2})(x^3)^{-2}$
$= 4^{-2}x^{-6}$
$= \frac{1}{4^2x^6}$
$= \frac{1}{16x^6}$

**56.** $(10x^2)^{-3} = 10^{-3}x^{2\cdot(-3)}$
$= 10^{-3}x^{-6}$
$= \frac{1}{10^3x^6}$
$= \frac{1}{1000x^6}$

**57.** $\frac{24x^3 \cdot y^5}{32x^7y^{-9}} = \frac{3}{4}x^{3-7}y^{5-(-9)}$
$= \frac{3}{4}x^{-4}y^{14}$
$= \frac{3y^{14}}{4x^4}$

**58.** $\frac{10x^4y^9}{30x^{12}y^{-3}} = \frac{1}{3}x^{4-12}y^{9-(-3)}$
$= \frac{1}{3}x^{-8}y^{12}$
$= \frac{y^{12}}{3x^8}$

**59.** $\left(\frac{5x^3}{y}\right)^{-2} = \frac{5^{-2}x^{-6}}{y^{-2}} = \frac{y^2}{25x^6}$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**60.** $\left(\frac{3x^4}{y}\right)^{-3} = \left(\frac{y}{3x^4}\right)^3$

$= \frac{y^3}{3^3 x^{4\cdot 3}}$

$= \frac{y^3}{27x^{12}}$

**61.** $\left(\frac{-15a^4b^2}{5a^{10}b^{-3}}\right)^3 = \left(\frac{-3b^{2-(-3)}}{a^{10-4}}\right)^3$

$= \left(\frac{-3b^5}{a^6}\right)^3$

$= \frac{-27b^{15}}{a^{18}}$

**62.** $\left(\frac{-30a^{14}b^8}{10a^{17}b^{-2}}\right)^3 = \left(\frac{-3b^{8-(-2)}}{a^{17-14}}\right)^3$

$= \left(\frac{-3b^{10}}{a^3}\right)^3$

$= \frac{-27b^{30}}{a^9}$

**63.** $\left(\frac{3a^{-5}b^2}{12a^3b^{-4}}\right)^0 = 1$

**64.** $\left(\frac{4a^{-5}b^3}{12a^3b^{-5}}\right)^0 = 1$

**65.** $3.8\times10^2 = 380$

**66.** $9.2\times10^2 = 920$

**67.** $6\times10^{-4} = 0.0006$

**68.** $7\times10^{-5} = 0.00007$

**69.** $-7.16\times10^6 = -7{,}160{,}000$

**70.** $-8.17\times10^6 = -8{,}170{,}000$

**71.** $7.9\times10^{-1} = 0.79$

**72.** $6.8\times10^{-1} = 0.68$

**73.** $-4.15\times10^{-3} = -0.00415$

**74.** $-3.14\times10^{-3} = -0.00314$

**75.** $-6.00001\times10^{10} = -60{,}000{,}100{,}000$

**76.** $-7.00001\times10^{10} = -70{,}000{,}100{,}000$

**77.** $32{,}000 = 3.2\times10^4$

**78.** $64{,}000 = 6.4\times10^4$

**79.** $638{,}000{,}000{,}000{,}000{,}000$

$= 6.38\times10^{17}$

**80.** $579{,}000{,}000{,}000{,}000{,}000 = 5.79\times10^{17}$

**81.** $-5716 = -5.716\times10^3$

**82.** $-3829 = -3.829\times10^3$

**83.** $0.0027 = 2.7\times10^{-3}$

**84.** $0.0083 = 8.3\times10^{-3}$

**85.** $-0.00000000504 = -5.04\times10^{-9}$

**86.** $-0.00000000405 = -4.05\times10^{-9}$

**87.** $\left(3\times10^4\right)\left(2.1\times10^3\right) = (3\times2.1)\left(10^4\times10^3\right)$

$= 6.3\times10^{4+3} = 6.3\times10^7$

**88.** $\left(2\times10^4\right)\left(4.1\times10^3\right) = 8.2\times10^7$

**89.** $\left(1.6\times10^{15}\right)\left(4\times10^{-11}\right) = (1.6\times4)\left(10^{15}\times10^{-11}\right)$

$= 6.4\times10^{15+(-11)}$

$= 6.4\times10^4$

**90.** $\left(1.4\times10^{15}\right)\left(3\times10^{-11}\right) = (1.4\times3)\left(10^{15}\times10^{-11}\right)$

$= 4.2\times10^{15+(-11)}$

$= 4.2\times10^4$

**91.** $\left(6.1\times10^{-8}\right)\left(2\times10^{-4}\right) = (6.1\times2)\left(10^{-8}\times10^{-4}\right)$

$= 12.2\times10^{-8+(-4)}$

$= 12.2\times10^{-12}$

$= 1.22\times10^{-11}$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**92.** $\left(5.1\times10^{-8}\right)\left(3\times10^{-4}\right)=15.3\times10^{-12}$
$=1.53\times10^{-11}$

**93.** $\left(4.3\times10^{8}\right)\left(6.2\times10^{4}\right)$
$=\left(4.3\times6.2\right)\left(10^{8}\times10^{4}\right)$
$=26.66\times10^{8+4}$
$=26.66\times10^{12}$
$=2.666\times10^{13}\approx2.67\times10^{13}$

**94.** $\left(8.2\times10^{8}\right)\left(4.6\times10^{4}\right)$
$=37.72\times10^{8+4}=37.72\times10^{12}$
$=3.772\times10^{13}\approx3.77\times10^{13}$

**95.** $\dfrac{8.4\times10^{8}}{4\times10^{5}}=\dfrac{8.4}{4}\times\dfrac{10^{8}}{10^{5}}$
$=2.1\times10^{8-5}=2.1\times10^{3}$

**96.** $\dfrac{6.9\times10^{8}}{3\times10^{5}}=2.3\times10^{8-5}=2.3\times10^{3}$

**97.** $\dfrac{3.6\times10^{4}}{9\times10^{-2}}=\dfrac{3.6}{9}\times\dfrac{10^{4}}{10^{-2}}$
$=0.4\times10^{4-(-2)}$
$=0.4\times10^{6}=4\times10^{5}$

**98.** $\dfrac{1.2\times10^{4}}{2\times10^{-2}}=0.6\times10^{4-(-2)}=0.6\times10^{6}$
$=\left(6\times10^{-1}\right)\times10^{6}=6\times10^{5}$

**99.** $\dfrac{4.8\times10^{-2}}{2.4\times10^{6}}=\dfrac{4.8}{2.4}\times\dfrac{10^{-2}}{10^{6}}$
$=2\times10^{-2-6}=2\times10^{-8}$

**100.** $\dfrac{7.5\times10^{-2}}{2.5\times10^{6}}=3\times10^{-2-6}=3\times10^{-8}$

**101.** $\dfrac{2.4\times10^{-2}}{4.8\times10^{-6}}=\dfrac{2.4}{4.8}\times\dfrac{10^{-2}}{10^{-6}}$
$=0.5\times10^{-2-(-6)}$
$=0.5\times10^{4}=5\times10^{3}$

**102.** $\dfrac{1.5\times10^{-2}}{5\times10^{-6}}=0.5\times10^{-2-(-6)}$
$=0.5\times10^{4}=5\times10^{3}$

**103.** $\dfrac{480,000,000,000}{0.00012}=\dfrac{4.8\times10^{11}}{1.2\times10^{-4}}$
$=\dfrac{4.8}{1.2}\times\dfrac{10^{11}}{10^{-4}}$
$=4\times10^{11-(-4)}$
$=4\times10^{15}$

**104.** $\dfrac{282,000,000,000}{0.00141}=\dfrac{2.82\times10^{11}}{1.41\times10^{-3}}$
$=2\times10^{11-(-3)}$
$=2\times10^{14}$

**105.** $\dfrac{0.00072\times0.003}{0.00024}$
$=\dfrac{\left(7.2\times10^{-4}\right)\left(3\times10^{-3}\right)}{2.4\times10^{-4}}$
$=\dfrac{7.2\times3}{2.4}\times\dfrac{10^{-4}\cdot10^{-3}}{10^{-4}}=9\times10^{-3}$

**106.** $\dfrac{66000\times0.001}{0.003\times0.002}=\dfrac{\left(6.6\times10^{4}\right)\left(1\times10^{-3}\right)}{\left(3\times10^{-3}\right)\left(2\times10^{-3}\right)}$
$=\dfrac{6.6\times10^{1}}{6\times10^{-6}}=1.1\times10^{1-(-6)}$
$=1.1\times10^{7}$

**107.** $\dfrac{\left(x^{-2}y\right)^{-3}}{\left(x^{2}y^{-1}\right)^{3}}=\dfrac{x^{6}y^{-3}}{x^{6}y^{-3}}$
$=x^{6-6}y^{-3-(-3)}=x^{0}y^{0}=1$

**108.** $\dfrac{\left(xy^{-2}\right)^{-2}}{\left(x^{-2}y\right)^{-3}}=\dfrac{x^{-2}y^{4}}{x^{6}y^{-3}}$
$=x^{-2-6}y^{4-(-3)}=x^{-8}y^{7}=\dfrac{y^{7}}{x^{8}}$

**109.** $\left(2x^{-3}yz^{-6}\right)(2x)^{-5}=2x^{-3}yz^{-6}\cdot2^{-5}x^{\cdot-5}$
$=2^{-4}x^{-8}yz^{-6}=\dfrac{y}{2^{4}x^{8}z^{6}}=\dfrac{y}{16x^{8}z^{6}}$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**110.** $\left(3x^{-4}yz^{-7}\right)(3x)^{-3} = 3x^{-4}yz^{-7}\cdot 3^{-3}x^{\cdot -3}$

$= 3^{-2}x^{-7}yz^{-7} = \dfrac{y}{3^2x^7z^7} = \dfrac{y}{9x^7z^7}$

**111.** $\left(\dfrac{x^3y^4z^5}{x^{-3}y^{-4}z^{-5}}\right)^{-2} = \left(x^6y^8z^{10}\right)^{-2}$

$= x^{-12}y^{-16}z^{-20} = \dfrac{1}{x^{12}y^{16}z^{20}}$

**112.** $\left(\dfrac{x^4y^5z^6}{x^{-4}y^{-5}z^{-6}}\right)^{-4} = \left(x^8y^{10}z^{12}\right)^{-4}$

$= x^{-32}y^{-40}z^{-48} = \dfrac{1}{x^{32}y^{40}z^{48}}$

**113.** $\dfrac{\left(2^{-1}x^{-2}y^{-1}\right)^{-2}\left(2x^{-4}y^3\right)^{-2}\left(16x^{-3}y^3\right)^0}{\left(2x^{-3}y^{-5}\right)^2}$

$= \dfrac{\left(2^2x^2y^2\right)\left(2^{-2}x^8y^{-6}\right)(1)}{\left(2^2x^{-6}y^{-10}\right)}$

$= \dfrac{x^{18}y^6}{4}$

**114.** $\dfrac{\left(2^{-1}x^{-3}y^{-1}\right)^{-2}\left(2x^{-6}y^4\right)^{-2}\left(9x^3y^{-3}\right)^0}{\left(2x^{-4}y^{-6}\right)^2}$

$= \dfrac{\left(2^2x^6y^2\right)\left(2^{-2}x^{12}y^{-8}\right)(1)}{\left(2^2x^{-8}y^{-12}\right)}$

$= \dfrac{x^{26}y^6}{4}$

**115.** **a.** $5.19\times10^{11}$

**b.** $4.8\times10^7$

**c.** $\dfrac{5.19\times10^{11}}{4.8\times10^7} = \dfrac{5.19}{4.8}\times\dfrac{10^{11}}{10^7}$

$\approx 1.0813\times10^4$

$\approx 10{,}813$

$10,813 yearly per person

**d.** $\dfrac{10{,}813}{12} \approx 901$

$901 monthly per person

**116.** **a.** $3.3\times10^{10}$

**b.** $2.57\times10^7$

**c.** $\dfrac{3.3\times10^{10}}{2.57\times10^7} = \dfrac{3.3}{2.57}\times\dfrac{10^{10}}{10^7}$

$\approx 1.284\times10^3$

$\approx 1284$

$1284 yearly per person

**d.** $\dfrac{1284}{12} = 107$

$107 monthly per person

**117.** Medicaid

$\dfrac{1.98\times10^{11}}{5.34\times10^7} = \dfrac{1.98}{5.34}\times\dfrac{10^{11}}{10^7}$

$\approx 0.3708\times10^4$

$\approx 3708$

Medicare

$\dfrac{2.94\times10^{11}}{4.23\times10^7} = \dfrac{2.94}{4.23}\times\dfrac{10^{11}}{10^7}$

$\approx 0.6950\times10^4$

$\approx 6950$

Medicare provides the greater yearly per person benefit by, $6950  $3708, or $3242.

**118.** **a.** $2.55\times10^{13}$

**b.** $2.54\times10^{11}$

**c.** $\dfrac{2.55\times10^{13}}{2.54\times10^{11}} = \dfrac{2.55}{2.54}\times\dfrac{10^{13}}{10^{11}}$

$\approx 1.00\times10^2$

$\approx 100$

approximately 100 years

**119.** **a.** $2.55\times10^{13}$

**b.** $6\times10^4$

**c.** $\dfrac{2.55\times10^{13}}{6\times10^4} = \dfrac{2.55}{6}\times\dfrac{10^{13}}{10^4}$

$= 0.425\times10^9$

$= 425{,}000{,}000$

425,000,000 Americans

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**120.** **a.** $9.84\times10^{11}$

**b.** $3.2\times10^{7}$

**c.** $\dfrac{9.84\times10^{11}}{3.2\times10^{7}}=\dfrac{9.84}{3.2}\times\dfrac{10^{11}}{10^{7}}$

$=3.075\times10^{4}$

$=30,750$

30,750 years

**121. – 128.** Answers will vary.

**129.** does not make sense; Explanations will vary.
Sample explanation: $36(x^3)^9=36x^{27}$ not $36x^{12}$.

**130.** makes sense

**131.** does not make sense; Explanations will vary.
Sample explanation: $4.6\times10^{12}$ represents over 4 trillion. The entire world population is measured in billions ($10^9$).

**132.** makes sense

**133.** false; Changes to make the statement true will vary.
A sample change is: $4^{-2}>4^{-3}$.

**134.** true

**135.** false; Changes to make the statement true will vary.
A sample change is: $(-2)^4\neq2^{-4}$ because $16\neq\dfrac{1}{16}$.

**136.** false; Changes to make the statement true will vary.
A sample change is: $5^2\cdot5^{-2}=2^5\cdot2^{-5}$.

**137.** false; Changes to make the statement true will vary.
A sample change is: $534.7\neq5347$.

**138.** false; Changes to make the statement true will vary.
A sample change is:
$\dfrac{8\times10^{30}}{2\times10^{-5}}=4\times10^{30-(-5)}=4\times10^{35}$.

**139.** false; Changes to make the statement true will vary.
A sample change is:
$(7\times10^5)+(2\times10^{-3})=700,000.002$.

**140.** true

**141.** The doctor has gathered:

$2^{-1}+2^{-2}=\dfrac{1}{2}+\dfrac{1}{2^2}=\dfrac{2}{4}+\dfrac{1}{4}=\dfrac{3}{4}$

So, $1-\dfrac{3}{4}=\dfrac{1}{4}$ is remaining.

**142.** $b^A=MN, b^C=M, b^D=N$

$b^A=b^Cb^D$

$A=C+D$

**143.** $\dfrac{70\text{ bts}}{\cancel{\text{min}}}\cdot\dfrac{60\cancel{\text{ min}}}{\cancel{\text{hr}}}\cdot\dfrac{24\cancel{\text{ hrs}}}{\cancel{\text{day}}}\cdot\dfrac{365\cancel{\text{ days}}}{\cancel{\text{yr}}}\cdot80\cancel{\text{ yrs}}$

$=70\cdot60\cdot24\cdot365\cdot80$ beats

$=2943360000$ beats

$=2.94336\times10^9$ beats

$\approx2.94\times10^9$ beats

The heartbeats approximately $2.94\times10^9$ times over a lifetime of 80 years.

**144.** Answers will vary.

**145.** **a.** $\sqrt{16}\cdot\sqrt{4}=4\cdot2=8$

**b.** $\sqrt{16\cdot4}=\sqrt{64}=8$

**c.** $\sqrt{16}\cdot\sqrt{4}=\sqrt{16\cdot4}$

**146.** **a.** $\sqrt{300}\approx17.32$

**b.** $10\sqrt{3}\approx17.32$

**c.** $\sqrt{300}=10\sqrt{3}$

**147.** **a.** $21x+10x=31x$

**b.** $21\sqrt{2}+10\sqrt{2}=31\sqrt{2}$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**Section P.3**

**Check Point Exercises**

**1. a.** $\sqrt{81} = 9$

**b.** $-\sqrt{9} = -3$

**c.** $\sqrt{\frac{1}{25}} = \frac{1}{5}$

**d.** $\sqrt{36+64} = \sqrt{100} = 10$

**e.** $\sqrt{36} + \sqrt{64} = 6 + 8 = 14$

**2. a.** $\sqrt{75} = \sqrt{25 \cdot 3} = \sqrt{25}\sqrt{3} = 5\sqrt{3}$

**b.** $\sqrt{5x} \cdot \sqrt{10x} = \sqrt{5x \cdot 10x}$
$= \sqrt{50x^2}$
$= \sqrt{25 \cdot 2x^2}$
$= \sqrt{25x^2} \cdot \sqrt{2}$
$= 5x\sqrt{2}$

**3. a.** $\sqrt{\frac{25}{16}} = \frac{\sqrt{25}}{\sqrt{16}} = \frac{5}{4}$

**b.** $\frac{\sqrt{150x^3}}{\sqrt{2x}} = \sqrt{\frac{150x^3}{2x}}$
$= \sqrt{75x^2}$
$= \sqrt{25x^2} \cdot \sqrt{3}$
$= 5x\sqrt{3}$

**4. a.** $8\sqrt{13} + 9\sqrt{13} = (8+9)\sqrt{3}$
$= 17\sqrt{13}$

**b.** $\sqrt{17x} - 20\sqrt{17x}$
$= 1\sqrt{17x} - 20\sqrt{17x}$
$= (1-20)\sqrt{17x}$
$= -19\sqrt{17x}$

**5. a.** $5\sqrt{27} + \sqrt{12}$
$= 5\sqrt{9 \cdot 3} + \sqrt{4 \cdot 3}$
$= 5 \cdot 3\sqrt{3} + 2\sqrt{3}$
$= 15\sqrt{3} + 2\sqrt{3}$
$= (15+2)\sqrt{3}$
$= 17\sqrt{3}$

**b.** $6\sqrt{18x} - 4\sqrt{8x}$
$= 6\sqrt{9 \cdot 2x} - 4\sqrt{4 \cdot 2x}$
$= 6 \cdot 3\sqrt{2x} - 4 \cdot 2\sqrt{2x}$
$= 18\sqrt{2x} - 8\sqrt{2x}$
$= (18-8)\sqrt{2x}$
$= 10\sqrt{2x}$

**6. a.** If we multiply numerator and denominator by $\sqrt{3}$, the denominator becomes $\sqrt{3} \cdot \sqrt{3} = \sqrt{9} = 3$. Therefore, multiply by 1, choosing $\frac{\sqrt{3}}{\sqrt{3}}$ for 1.

$$\frac{5}{\sqrt{3}} = \frac{5}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{5\sqrt{3}}{\sqrt{9}} = \frac{5\sqrt{3}}{3}$$

**b.** The *smallest* number that will produce a perfect square in the denominator of $\frac{6}{\sqrt{12}}$ is $\sqrt{3}$ because $\sqrt{12} \cdot \sqrt{3} = \sqrt{36} = 6$. So multiply by 1, choosing $\frac{\sqrt{3}}{\sqrt{3}}$ for 1.

$$\frac{6}{\sqrt{12}} = \frac{6}{\sqrt{12}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{6\sqrt{3}}{\sqrt{36}} = \frac{6\sqrt{3}}{6} = \sqrt{3}$$

**7.** Multiply by $\frac{4-\sqrt{5}}{4-\sqrt{5}}$.

$$\frac{8}{4+\sqrt{5}} = \frac{8}{4+\sqrt{5}} \cdot \frac{4-\sqrt{5}}{4-\sqrt{5}}$$
$$= \frac{8(4-\sqrt{5})}{4^2 - (\sqrt{5})^2}$$
$$= \frac{8(4-\sqrt{5})}{16-5}$$
$$= \frac{8(4-\sqrt{5})}{11} \text{ or } \frac{32-8\sqrt{5}}{11}$$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**8.** **a.** $\sqrt[3]{40} = \sqrt[3]{8 \cdot 5} = \sqrt[3]{8} \cdot \sqrt[3]{5} = 2\sqrt[3]{5}$

**b.** $\sqrt[5]{8} \cdot \sqrt[5]{8} = \sqrt[5]{64} = \sqrt[5]{32} \cdot \sqrt[5]{2} = 2\sqrt[5]{2}$

**c.** $\sqrt[3]{\frac{125}{27}} = \frac{\sqrt[3]{125}}{\sqrt[3]{27}} = \frac{5}{3}$

**9.**
$$\begin{aligned} & 3\sqrt[3]{81} - 4\sqrt[3]{3} \\ &= 3\sqrt[3]{27 \cdot 3} - 4\sqrt[3]{3} \\ &= 3 \cdot 3\sqrt[3]{3} - 4\sqrt[3]{3} \\ &= 9\sqrt[3]{3} - 4\sqrt[3]{3} \\ &= (9-4)\sqrt[3]{3} \\ &= 5\sqrt[3]{3} \end{aligned}$$

**10.** **a.** $25^{\frac{1}{2}} = \sqrt{25} = 5$

**b.** $8^{\frac{1}{3}} = \sqrt[3]{8} = 2$

**c.** $-81^{\frac{1}{4}} = -\sqrt[4]{81} = -3$

**d.** $(-8)^{\frac{1}{3}} = \sqrt[3]{-8} = -2$

**e.** $27^{-\frac{1}{3}} = \frac{1}{27^{\frac{1}{3}}} = \frac{1}{\sqrt[3]{27}} = \frac{1}{3}$

**11.** **a.** $27^{\frac{4}{3}} = \left(\sqrt[3]{27}\right)^4 = (3)^4 = 81$

**b.** $4^{\frac{3}{2}} = \left(\sqrt{4}\right)^3 = (2)^3 = 8$

**c.** $32^{-\frac{2}{5}} = \frac{1}{32^{\frac{2}{5}}} = \frac{1}{\left(\sqrt[5]{32}\right)^2} = \frac{1}{2^2} = \frac{1}{4}$

**12.** **a.**
$$\begin{aligned} & \left(2x^{4/3}\right)\left(5x^{8/3}\right) \\ &= 2 \cdot 5x^{4/3} \cdot x^{8/3} \\ &= 10x^{(4/3)+(8/3)} \\ &= 10x^{12/3} \\ &= 10x^4 \end{aligned}$$

**b.**
$$\begin{aligned} \frac{20x^4}{5x^{3/2}} &= \left(\frac{20}{5}\right)\left(\frac{x^4}{x^{3/2}}\right) \\ &= 4x^{4-(3/2)} \\ &= 4x^{(8/2)-(3/2)} \\ &= 4x^{5/2} \end{aligned}$$

**13.** $\sqrt[6]{x^3} = x^{3/6} = x^{1/2} = \sqrt{x}$

**Concept and Vocabulary Check P.3**

**C1.** principal

**C2.** $8^2$

**C3.** $|a|$

**C4.** $\sqrt{a} \cdot \sqrt{b}$

**C5.** $\frac{\sqrt{a}}{\sqrt{b}}$

**C6.** $18\sqrt{3}$

**C7.** 5; $6\sqrt{3}$

**C8.** $7 - \sqrt{3}$

**C9.** $\sqrt{10} + \sqrt{2}$

**C10.** index; radicand

**C11.** $(-2)^5$

**C12.** $a$; $|a|$

**C13.** $\sqrt[n]{a}$

**C14.** 2; 8

**Exercise Set P.3**

**1.** $\sqrt{36} = \sqrt{6^2} = 6$

**2.** $\sqrt{25} = \sqrt{5^2} = 5$

**3.** $-\sqrt{36} = -\sqrt{6^2} = -6$

**4.** $-\sqrt{25} = -\sqrt{5^2} = -5$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**5.** $\sqrt{-36}$, The square root of a negative number is not real.

**6.** $\sqrt{-25}$, The square root of a negative number is not real.

**7.** $\sqrt{25-16} = \sqrt{9} = 3$

**8.** $\sqrt{144+25} = \sqrt{169} = 13$

**9.** $\sqrt{25} - \sqrt{16} = 5 - 4 = 1$

**10.** $\sqrt{144} + \sqrt{25} = 12 + 5 = 17$

**11.** $\sqrt{(-13)^2} = \sqrt{169} = 13$

**12.** $\sqrt{(-17)^2} = \sqrt{289} = 17$

**13.** $\sqrt{50} = \sqrt{25 \cdot 2} = \sqrt{25}\sqrt{2} = 5\sqrt{2}$

**14.** $\sqrt{27} = \sqrt{9 \cdot 3} = \sqrt{9}\sqrt{3} = 3\sqrt{3}$

**15.** $\begin{aligned}\sqrt{45x^2} &= \sqrt{9x^2 \cdot 5} \\ &= \sqrt{9x^2}\sqrt{5} \\ &= \sqrt{9}\sqrt{x^2}\sqrt{5} \\ &= 3|x|\sqrt{5}\end{aligned}$

**16.** $\begin{aligned}\sqrt{125x^2} &= \sqrt{25x^2 \cdot 5} \\ &= \sqrt{25x^2}\sqrt{5} \\ &= \sqrt{25}\sqrt{x^2}\sqrt{5} \\ &= 5|x|\sqrt{5}\end{aligned}$

**17.** $\begin{aligned}\sqrt{2x} \cdot \sqrt{6x} &= \sqrt{2x \cdot 6x} \\ &= \sqrt{12x^2} \\ &= \sqrt{4x^2} \cdot \sqrt{3} \\ &= 2x\sqrt{3}\end{aligned}$

**18.** $\begin{aligned}\sqrt{10x} \cdot \sqrt{8x} &= \sqrt{10x \cdot 8x} \\ &= \sqrt{80x^2} \\ &= \sqrt{16x^2} \cdot \sqrt{5} \\ &= 4x\sqrt{5}\end{aligned}$

**19.** $\sqrt{x^3} = \sqrt{x^2} \cdot \sqrt{x} = x\sqrt{x}$

**20.** $\sqrt{y^3} = \sqrt{y^2} \cdot \sqrt{y} = y\sqrt{y}$

**21.** $\begin{aligned}\sqrt{2x^2} \cdot \sqrt{6x} &= \sqrt{2x^2 \cdot 6x} \\ &= \sqrt{12x^3} \\ &= \sqrt{4x^2} \cdot \sqrt{3x} \\ &= 2x\sqrt{3x}\end{aligned}$

**22.** $\begin{aligned}\sqrt{6x} \cdot \sqrt{3x^2} &= \sqrt{6x \cdot 3x^2} \\ &= \sqrt{18x^3} \\ &= \sqrt{9x^2} \cdot \sqrt{2x} \\ &= 3x\sqrt{2x}\end{aligned}$

**23.** $\sqrt{\frac{1}{81}} = \frac{\sqrt{1}}{\sqrt{81}} = \frac{1}{9}$

**24.** $\sqrt{\frac{1}{49}} = \frac{\sqrt{1}}{\sqrt{49}} = \frac{1}{7}$

**25.** $\sqrt{\frac{49}{16}} = \frac{\sqrt{49}}{\sqrt{16}} = \frac{7}{4}$

**26.** $\sqrt{\frac{121}{9}} = \frac{\sqrt{121}}{\sqrt{9}} = \frac{11}{3}$

**27.** $\frac{\sqrt{48x^3}}{\sqrt{3x}} = \sqrt{\frac{48x^3}{3x}} = \sqrt{16x^2} = 4x$

**28.** $\frac{\sqrt{72x^3}}{\sqrt{8x}} = \sqrt{\frac{72x^3}{8x}} = \sqrt{9x^2} = 3x$

**29.** $\begin{aligned}\frac{\sqrt{150x^4}}{\sqrt{3x}} &= \sqrt{\frac{150x^4}{3x}} \\ &= \sqrt{50x^3} \\ &= \sqrt{25x^2} \cdot \sqrt{2x} \\ &= 5x\sqrt{2x}\end{aligned}$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**30.** $\dfrac{\sqrt{24x^4}}{\sqrt{3x}} = \sqrt{\dfrac{24x^4}{3x}}$
$= \sqrt{8x^3}$
$= \sqrt{4x^2}\cdot\sqrt{2x}$
$= 2x\sqrt{2x}$

**31.** $\dfrac{\sqrt{200x^3}}{\sqrt{10x^{-1}}} = \sqrt{\dfrac{200x^3}{10x^{-1}}}$
$= \sqrt{20x^{3-(-1)}}$
$= \sqrt{20x^4}$
$= \sqrt{4\cdot 5x^4}$
$= 2x^2\sqrt{5}$

**32.** $\dfrac{\sqrt{500x^3}}{\sqrt{10x^{-1}}} = \sqrt{\dfrac{500x^3}{10x^{-1}}} = \sqrt{50x^{3-(-1)}}$
$= \sqrt{50x^4} = \sqrt{25\cdot 2x^4} = 5x^2\sqrt{2}$

**33.** $7\sqrt{3}+6\sqrt{3} = (7+6)\sqrt{3} = 13\sqrt{3}$

**34.** $8\sqrt{5}+11\sqrt{5} = (8+11)\sqrt{5} = 19\sqrt{5}$

**35.** $6\sqrt{17x}-8\sqrt{17x} = (6-8)\sqrt{17x} = -2\sqrt{17x}$

**36.** $4\sqrt{13x}-6\sqrt{13x} = (4-6)\sqrt{13x} = -2\sqrt{13x}$

**37.** $\sqrt{8}+3\sqrt{2} = \sqrt{4\cdot 2}+3\sqrt{2}$
$= 2\sqrt{2}+3\sqrt{2}$
$= (2+3)\sqrt{2}$
$= 5\sqrt{2}$

**38.** $\sqrt{20}+6\sqrt{5} = \sqrt{4\cdot 5}+6\sqrt{5}$
$= 2\sqrt{5}+6\sqrt{5}$
$= (2+6)\sqrt{5}$
$= 8\sqrt{5}$

**39.** $\sqrt{50x}-\sqrt{8x} = \sqrt{25\cdot 2x}-\sqrt{4\cdot 2x}$
$= 5\sqrt{2x}-2\sqrt{2x}$
$= (5-2)\sqrt{2x}$
$= 3\sqrt{2x}$

**40.** $\sqrt{63x}-\sqrt{28x} = \sqrt{9\cdot 7x}-\sqrt{4\cdot 7x}$
$= 3\sqrt{7x}-2\sqrt{7x}$
$= (3-2)\sqrt{7x}$
$= \sqrt{7x}$

**41.** $3\sqrt{18}+5\sqrt{50} = 3\sqrt{9\cdot 2}+5\sqrt{25\cdot 2}$
$= 3\cdot 3\sqrt{2}+5\cdot 5\sqrt{2}$
$= 9\sqrt{2}+25\sqrt{2}$
$= (9+25)\sqrt{2}$
$= 34\sqrt{2}$

**42.** $4\sqrt{12}-2\sqrt{75} = 4\sqrt{4\cdot 3}-2\sqrt{25\cdot 3}$
$= 4\cdot 2\sqrt{3}-2\cdot 5\sqrt{3}$
$= 8\sqrt{3}-10\sqrt{3}$
$= (8-10)\sqrt{3}$
$= -2\sqrt{3}$

**43.** $3\sqrt{8}-\sqrt{32}+3\sqrt{72}-\sqrt{75}$
$= 3\sqrt{4\cdot 2}-\sqrt{16\cdot 2}+3\sqrt{36\cdot 2}-\sqrt{25\cdot 3}$
$= 3\cdot 2\sqrt{2}-4\sqrt{2}+3\cdot 6\sqrt{2}-5\sqrt{3}$
$= 6\sqrt{2}-4\sqrt{2}+18\sqrt{2}-5\sqrt{3}$
$= 20\sqrt{2}-5\sqrt{3}$

**44.** $3\sqrt{54}-2\sqrt{24}-\sqrt{96}+4\sqrt{63}$
$= 3\sqrt{9\cdot 6}-2\sqrt{4\cdot 6}-\sqrt{16\cdot 6}+4\sqrt{9\cdot 7}$
$= 3\cdot 3\sqrt{6}-2\cdot 2\sqrt{6}-4\sqrt{6}+4\cdot 3\sqrt{7}$
$= 9\sqrt{6}-4\sqrt{6}-4\sqrt{6}+12\sqrt{7}$
$= \sqrt{6}+12\sqrt{7}$

**45.** $\dfrac{1}{\sqrt{7}} = \dfrac{1}{\sqrt{7}}\cdot\dfrac{\sqrt{7}}{\sqrt{7}} = \dfrac{\sqrt{7}}{7}$

**46.** $\dfrac{2}{\sqrt{10}} = \dfrac{2}{\sqrt{10}}\cdot\dfrac{\sqrt{10}}{\sqrt{10}} = \dfrac{2\sqrt{10}}{10} = \dfrac{\sqrt{10}}{5}$

**47.** $\dfrac{\sqrt{2}}{\sqrt{5}} = \dfrac{\sqrt{2}}{\sqrt{5}}\cdot\dfrac{\sqrt{5}}{\sqrt{5}} = \dfrac{\sqrt{10}}{5}$

**48.** $\dfrac{\sqrt{7}}{\sqrt{3}} = \dfrac{\sqrt{7}}{\sqrt{3}}\cdot\dfrac{\sqrt{3}}{\sqrt{3}} = \dfrac{\sqrt{21}}{3}$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**49.** $$\frac{13}{3+\sqrt{11}} = \frac{13}{3+\sqrt{11}} \cdot \frac{3-\sqrt{11}}{3-\sqrt{11}}$$
$$= \frac{13(3-\sqrt{11})}{3^2-(\sqrt{11})^2}$$
$$= \frac{13(3-\sqrt{11})}{9-11}$$
$$= \frac{13(3-\sqrt{11})}{-2}$$

**50.** $$\frac{3}{3+\sqrt{7}} = \frac{3}{3+\sqrt{7}} \cdot \frac{3-\sqrt{7}}{3-\sqrt{7}}$$
$$= \frac{3(3-\sqrt{7})}{3^2-(\sqrt{7})^2}$$
$$= \frac{3(3-\sqrt{7})}{9-7}$$
$$= \frac{3(3-\sqrt{7})}{2}$$

**51.** $$\frac{7}{\sqrt{5}-2} = \frac{7}{\sqrt{5}-2} \cdot \frac{\sqrt{5}+2}{\sqrt{5}+2}$$
$$= \frac{7(\sqrt{5}+2)}{(\sqrt{5})^2-2^2}$$
$$= \frac{7(\sqrt{5}+2)}{5-4}$$
$$= 7(\sqrt{5}+2)$$

**52.** $$\frac{5}{\sqrt{3}-1} = \frac{5}{\sqrt{3}-1} \cdot \frac{\sqrt{3}+1}{\sqrt{3}+1}$$
$$= \frac{5(\sqrt{3}+1)}{(\sqrt{3})^2-1^2}$$
$$= \frac{5(\sqrt{3}+1)}{3-1}$$
$$= \frac{5(\sqrt{3}+1)}{2}$$

**53.** $$\frac{6}{\sqrt{5}+\sqrt{3}} = \frac{6}{\sqrt{5}+\sqrt{3}} \cdot \frac{\sqrt{5}-\sqrt{3}}{\sqrt{5}-\sqrt{3}}$$
$$= \frac{6(\sqrt{5}-\sqrt{3})}{(\sqrt{5})^2-(\sqrt{3})^2}$$
$$= \frac{6(\sqrt{5}-\sqrt{3})}{5-3}$$
$$= \frac{6(\sqrt{5}-\sqrt{3})}{2}$$
$$= 3(\sqrt{5}-\sqrt{3})$$

**54.** $$\frac{11}{\sqrt{7}-\sqrt{3}} = \frac{11}{\sqrt{7}-\sqrt{3}} \cdot \frac{\sqrt{7}+\sqrt{3}}{\sqrt{7}+\sqrt{3}}$$
$$= \frac{11(\sqrt{7}+\sqrt{3})}{(\sqrt{7})^2-(\sqrt{3})^2}$$
$$= \frac{11(\sqrt{7}+\sqrt{3})}{7-3}$$
$$= \frac{11(\sqrt{7}+\sqrt{3})}{4}$$

**55.** $\sqrt[3]{125} = \sqrt[3]{5^3} = 5$

**56.** $\sqrt[3]{8} = \sqrt[3]{2^3} = 2$

**57.** $\sqrt[3]{-8} = \sqrt[3]{(-2)^3} = -2$

**58.** $\sqrt[3]{-125} = \sqrt[3]{(-5)^3} = -5$

**59.** $\sqrt[4]{-16}$ is not a real number.

**60.** $\sqrt[4]{-81}$ is not a real number.

**61.** $\sqrt[4]{(-3)^4} = |-3| = 3$

**62.** $\sqrt[4]{(-2)^4} = |-2| = 2$

**63.** $\sqrt[5]{(-3)^5} = -3$

**64.** $\sqrt[5]{(-2)^5} = -2$

**65.** $\sqrt[5]{-\frac{1}{32}} = \sqrt[5]{-\frac{1}{2^5}} = -\frac{1}{2}$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**66.** $\sqrt[6]{\frac{1}{64}} = \frac{\sqrt[6]{1}}{\sqrt[6]{2^6}} = \frac{1}{2}$

**67.** $\sqrt[3]{32} = \sqrt[3]{8 \cdot 4} = \sqrt[3]{8}\sqrt[3]{4} = 2 \cdot \sqrt[3]{4}$

**68.** $\sqrt[3]{150}$ cannot be simplified further.

**69.** $\sqrt[3]{x^4} = \sqrt[3]{x^3 \cdot x} = x \cdot \sqrt[3]{x}$

**70.** $\sqrt[3]{x^5} = \sqrt[3]{x^3 x^2} = x\sqrt[3]{x^2}$

**71.** $\sqrt[3]{9} \cdot \sqrt[3]{6} = \sqrt[3]{54} = \sqrt[3]{27 \cdot 2} = \sqrt[3]{27}\sqrt[3]{2} = 3\sqrt[3]{2}$

**72.** $\sqrt[3]{12} \cdot \sqrt[3]{4} = \sqrt[3]{48} = \sqrt[3]{8 \cdot 6} = 2\sqrt[3]{6}$

**73.** $\frac{\sqrt[5]{64x^6}}{\sqrt[5]{2x}} = \sqrt[5]{\frac{64x^6}{2x}} = \sqrt[5]{32x^5} = 2x$

**74.** $\frac{\sqrt[4]{162x^5}}{\sqrt[4]{2x}} = \sqrt[4]{\frac{162x^5}{2x}} = \sqrt[4]{81x^4} = 3x$

**75.** $4\sqrt[5]{2} + 3\sqrt[5]{2} = 7\sqrt[5]{2}$

**76.** $6\sqrt[5]{3} + 2\sqrt[5]{3} = 8\sqrt[5]{3}$

**77.** $5\sqrt[3]{16} + \sqrt[3]{54} = 5\sqrt[3]{8 \cdot 2} + \sqrt[3]{27 \cdot 2}$
$= 5 \cdot 2\sqrt[3]{2} + 3\sqrt[3]{2}$
$= 10\sqrt[3]{2} + 3\sqrt[3]{2}$
$= 13\sqrt[3]{2}$

**78.** $3\sqrt[3]{24} + \sqrt[3]{81} = \sqrt[3]{8 \cdot 3} + \sqrt[3]{27 \cdot 3}$
$= 3 \cdot 2\sqrt[3]{3} + 3\sqrt[3]{3}$
$= 6\sqrt[3]{3} + 3\sqrt[3]{3}$
$= 9\sqrt[3]{3}$

**79.** $\sqrt[3]{54xy^3} - y\sqrt[3]{128x}$
$= \sqrt[3]{27 \cdot 2xy^3} - y\sqrt[3]{64 \cdot 2x}$
$= 3y\sqrt[3]{2x} - 4y\sqrt[3]{2x}$
$= -y\sqrt[3]{2x}$

**80.** $\sqrt[3]{24xy^3} - y\sqrt[3]{81x}$
$= \sqrt[3]{8 \cdot 3xy^3} - y\sqrt[3]{27 \cdot 3x}$
$= 2y\sqrt[3]{3x} - 3y\sqrt[3]{3x}$
$= -y\sqrt[3]{3x}$

**81.** $\sqrt{2} + \sqrt[3]{8} = \sqrt{2} + 2$

**82.** $\sqrt{3} + \sqrt[3]{15}$ will not simplify.

**83.** $36^{1/2} = \sqrt{36} = 6$

**84.** $121^{1/2} = \sqrt{121} = 11$

**85.** $8^{1/3} = \sqrt[3]{8} = 2$

**86.** $27^{1/3} = \sqrt[3]{27} = 3$

**87.** $125^{2/3} = \left(\sqrt[3]{125}\right)^2 = 5^2 = 25$

**88.** $8^{2/3} = \left(\sqrt[3]{8}\right)^2 = 4$

**89.** $32^{-4/5} = \frac{1}{32^{4/5}} = \frac{1}{2^4} = \frac{1}{16}$

**90.** $16^{-5/2} = \frac{1}{16^{5/2}} = \frac{1}{(\sqrt{16})^5} = \frac{1}{4^5} = \frac{1}{1024}$

**91.** $\left(7x^{1/3}\right)\left(2x^{1/4}\right) = 7 \cdot 2x^{1/3} \cdot x^{1/4}$
$= 14 \cdot x^{1/3+1/4}$
$= 14x^{7/12}$

**92.** $(3x^{2/3})(4x^{3/4}) = 3 \cdot 4x^{2/3} \cdot x^{3/4}$
$= 12 \cdot x^{2/3+3/4}$
$= 12x^{17/12}$

**93.** $\frac{20x^{1/2}}{5x^{1/4}} = \left(\frac{20}{5}\right)\left(\frac{x^{1/2}}{x^{1/4}}\right)$
$= 4 \cdot x^{1/2-1/4}$
$= 4x^{1/4}$

**94.** $\frac{72x^{3/4}}{9x^{1/3}} = \left(\frac{72}{9}\right)\left(\frac{x^{3/4}}{x^{1/3}}\right) = 8 \cdot x^{3/4-1/3} = 8x^{5/12}$

**95.** $\left(x^{2/3}\right)^3 = x^{2/3 \cdot 3} = x^2$

**96.** $(x^{4/5})^5 = x^{4/5 \cdot 5} = x^4$

**97.** $(25x^4y^6)^{1/2} = 25^{1/2}x^{4 \cdot 1/2}y^{6 \cdot 1/2} = 5x^2|y|^3$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**98.** $(125x^9y^6)^{1/3} = 125^{1/3}x^{9/3}y^{6/3} = 5x^3y^2$

**99.** $$\frac{\left(3y^{\frac{1}{4}}\right)^3}{y^{\frac{1}{12}}} = \frac{27y^{\frac{3}{4}}}{y^{\frac{1}{12}}} = 27y^{\frac{3}{4}-\frac{1}{12}}$$
$$= 27y^{\frac{8}{12}} = 27y^{\frac{2}{3}}$$

**100.** $$\frac{\left(2y^{1/5}\right)^4}{y^{3/10}} = \frac{2^4\left(y^{1/5}\right)^4}{y^{3/10}}$$
$$= \frac{16y^{4/5}}{y^{3/10}} = 16y^{4/5-3/10} = 16y^{1/2}$$

**101.** $\sqrt[4]{5^2} = 5^{2/4} = 5^{1/2} = \sqrt{5}$

**102.** $\sqrt[4]{7^2} = 7^{2/4} = 7^{1/2} = \sqrt{7}$

**103.** $\sqrt[3]{x^6} = x^{6/3} = x^2$

**104.** $\sqrt[4]{x^{12}} = x^{12/4} = |x|^3$

**105.** $\sqrt[6]{x^4} = \sqrt[6/2]{x^{4/2}} = \sqrt[3]{x^2}$

**106.** $\sqrt[9]{x^6} = \sqrt[9/3]{x^{6/3}} = \sqrt[3]{x^2}$

**107.** $\sqrt[9]{x^6y^3} = x^{\frac{6}{9}}y^{\frac{3}{9}} = x^{\frac{2}{3}}y^{\frac{1}{3}} = \sqrt[3]{x^2y}$

**108.** $\sqrt[12]{x^4y^8} = |x|^{\frac{4}{12}}|y|^{\frac{8}{12}} = |x|^{\frac{1}{3}}|y|^{\frac{2}{3}} = \sqrt[3]{|x|y^2}$

**109.** $\sqrt[3]{\sqrt[4]{16}+\sqrt{625}} = \sqrt[3]{2+25} = \sqrt[3]{27} = 3$

**110.** $\sqrt[3]{\sqrt{\sqrt{169}+\sqrt{9}}+\sqrt{\sqrt[3]{1000}+\sqrt[3]{216}}}$
$= \sqrt[3]{\sqrt{13+3}+\sqrt{10+6}}$
$= \sqrt[3]{\sqrt{16}+\sqrt{16}}$
$= \sqrt[3]{4+4} = \sqrt[3]{8}$
$= 2$

**111.** $\left(49x^{-2}y^4\right)^{-1/2}\left(xy^{1/2}\right)$
$= (49)^{-1/2}\left(x^{-2}\right)^{-1/2}\left(y^4\right)^{-1/2}\left(xy^{1/2}\right)$
$= \frac{1}{49^{1/2}}x^{(-2)(-1/2)}y^{(4)(-1/2)}\left(xy^{1/2}\right)$
$= \frac{1}{7}x^1y^{-2}\cdot xy^{1/2} = \frac{1}{7}x^{1+1}y^{-2+(1/2)}$
$= \frac{1}{7}x^2y^{-3/2} = \frac{x^2}{7y^{3/2}}$

**112.** $\left(8x^{-6}y^3\right)^{1/3}\left(x^{5/6}y^{-1/3}\right)^6$
$= 8^{1/3}x^{(-6)(1/3)}y^{(3)(1/3)}x^{(5/6)(6)}y^{(-1/3)(6)}$
$= 2x^{-2}y^1x^5y^{-2} = 2x^{-2+5}y^{1+(-2)}$
$= 2x^3y^{-1} = \frac{2x^3}{y}$

**113.** $\left(\frac{x^{-5/4}y^{1/3}}{x^{-3/4}}\right)^{-6} = \left(x^{(-5/4)-(-3/4)}y^{1/3}\right)^{-6}$
$= \left(x^{-2/4}y^{1/3}\right)^{-6} = x^{(-2/4)(-6)}y^{(1/3)(-6)}$
$= x^3y^{-2} = \frac{x^3}{y^2}$

**114.** $\left(\frac{x^{1/2}y^{-7/4}}{y^{-5/4}}\right)^{-4} = \left(x^{1/2}y^{(-7/4)-(-5/4)}\right)^{-4}$
$= \left(x^{1/2}y^{-2/4}\right)^{-4} = x^{(1/2)(-4)}y^{(-2/4)(-4)}$
$= x^{-2}y^2 = \frac{y^2}{x^2}$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**115. a.** $P = \dfrac{x\left(13+\sqrt{x}\right)}{5\sqrt{x}}$

$= \dfrac{25\left(13+\sqrt{25}\right)}{5\sqrt{25}}$

$= \dfrac{25(13+5)}{5(5)}$

$= 18$

18% of 25-year-olds must pay more taxes.

**b.** $P = \dfrac{x\left(13+\sqrt{x}\right)}{5\sqrt{x}}$

$= \dfrac{13x+x\sqrt{x}}{5\sqrt{x}} \cdot \dfrac{\sqrt{x}}{\sqrt{x}}$

$= \dfrac{13x\sqrt{x}+x^2}{5x}$

$= \dfrac{13\sqrt{x}+x}{5}$

**c.** $P = \dfrac{13\sqrt{x}+x}{5}$

$= \dfrac{13\sqrt{25}+25}{5}$

$= \dfrac{13(5)+25}{5}$

$= 18$

Yes we get the same amount, 18%.
No, this does not prove we have correctly rationalized the denominator. Explanations will vary.

**116. a.** For 2020: $E = 5\sqrt{x}+34.1$

$= 5\sqrt{10}+34.1$

For 2050: $E = 5\sqrt{x}+34.1$

$= 5\sqrt{40}+34.1$

$= 5\cdot 2\sqrt{10}+34.1$

$= 10\sqrt{10}+34.1$

Difference:

$(10\sqrt{10}+34.1)-(5\sqrt{10}+34.1)$

$= 10\sqrt{10}+34.1-5\sqrt{10}-34.1$

$= 10\sqrt{10}-5\sqrt{10}+34.1-34.1$

$= 5\sqrt{10}$

The difference is $5\sqrt{10}$.

**b.** $5\sqrt{10} \approx 15.8$

This underestimates the difference projected by the graph of $65.8-47.3=18.5$ by 2.7. This represents a difference of 2.7 million people.

**117.** $\dfrac{2}{\sqrt{5}-1} \cdot \dfrac{\sqrt{5}+1}{\sqrt{5}+1} = \dfrac{2(\sqrt{5}+1)}{5-1}$

$= \dfrac{2(\sqrt{5}+1)}{4}$

$= \dfrac{\sqrt{5}+1}{2}$

$\approx 1.62$

About 1.62 to 1.

**118.** $R_a = R_f\sqrt{1-\left(\dfrac{v}{c}\right)^2}$

$= R_f\sqrt{1-\left(\dfrac{0.9c}{c}\right)^2}$

$= R_f\sqrt{1-(0.9)^2}$

$= R_f\sqrt{0.19}$

$\approx 0.44R_f$

$R_a = 0.44R_f$

$44 = 0.44R_f$

$\dfrac{44}{0.44} = \dfrac{0.44R_f}{0.44}$

$100 = R_f$

If you are gone for 44 weeks, then 100 weeks will have passed for your friend.


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**119.** Perimeter:

$$\begin{aligned} P &= 2l + 2w \\ &= 2 \cdot \sqrt{125} + 2 \cdot 2\sqrt{20} \\ &= 2 \cdot \sqrt{25 \cdot 5} + 4\sqrt{4 \cdot 5} \\ &= 2 \cdot 5\sqrt{5} + 4 \cdot 2\sqrt{5} \\ &= 10\sqrt{5} + 8\sqrt{5} \\ &= 18\sqrt{5} \text{ feet} \end{aligned}$$

Area:

$$\begin{aligned} A &= lw \\ &= \sqrt{125} \cdot 2\sqrt{20} \\ &= 2\sqrt{125 \cdot 20} \\ &= 2\sqrt{2500} \\ &= 2 \cdot 50 \\ &= 100 \text{ square feet} \end{aligned}$$

**120.** Perimeter:

$$\begin{aligned} P &= 2l + 2w \\ &= 2 \cdot 4\sqrt{20} + 2 \cdot \sqrt{80} \\ &= 8\sqrt{4 \cdot 5} + 2\sqrt{16 \cdot 5} \\ &= 8 \cdot 2\sqrt{5} + 2 \cdot 4\sqrt{5} \\ &= 16\sqrt{5} + 8\sqrt{5} \\ &= 24\sqrt{5} \text{ feet} \end{aligned}$$

Area:

$$\begin{aligned} A &= lw \\ &= 4\sqrt{20} \cdot \sqrt{80} \\ &= 4\sqrt{20 \cdot 80} \\ &= 4\sqrt{1600} \\ &= 4 \cdot 40 \\ &= 160 \text{ square feet} \end{aligned}$$

**121. – 128.** Answers will vary.

**129.** makes sense

**130.** makes sense

**131.** does not make sense; Explanations will vary. Sample explanation: $2\sqrt{20} + 4\sqrt{75}$ simplifies to $4\sqrt{5} + 20\sqrt{3}$ and thus the radical terms are not common.

**132.** does not make sense; Explanations will vary. Sample explanation: Finding the $n$th root first often gives smaller numbers on the middle step.

**133.** false; Changes to make the statement true will vary. A sample change is: $7^{1/2} \cdot 7^{1/2} = 7^1 = 7$.

**134.** false; Changes to make the statement true will vary. A sample change is: $(8)^{-\frac{1}{3}} = \dfrac{1}{(8)^{\frac{1}{3}}} = \dfrac{1}{\sqrt[3]{8}} = \dfrac{1}{2}$.

**135.** false; Changes to make the statement true will vary. The cube root of –8 is the real number –2.

**136.** false; Changes to make the statement true will vary. A sample change is: $\dfrac{\sqrt{20}}{8} = \dfrac{\sqrt{5}}{4}$.

**137.** $\left(5 + \sqrt{\boxed{3}}\right)\left(5 - \sqrt{\boxed{3}}\right) = 22$

$$\begin{aligned} 25 - \boxed{3} &= 22 \\ \boxed{3} &= 3 \end{aligned}$$

**138.** $\sqrt{\boxed{25}x^{\boxed{14}}} = 5x^7$

**139.**

$$\begin{aligned} &\sqrt{13 + \sqrt{2} + \frac{7}{3 + \sqrt{2}}} \\ &= \sqrt{13 + \sqrt{2} + \frac{7}{3 + \sqrt{2}} \cdot \frac{3 - \sqrt{2}}{3 - \sqrt{2}}} \\ &= \sqrt{13 + \sqrt{2} + \frac{21 - 7\sqrt{2}}{9 - 2}} \\ &= \sqrt{13 + \sqrt{2} + \frac{21 - 7\sqrt{2}}{7}} \\ &= \sqrt{13 + \sqrt{2} + 3 - \sqrt{2}} \\ &= \sqrt{16} \\ &= 4 \end{aligned}$$

**140. a.** $3^{\frac{1}{2}} \boxed{>} 3^{\frac{1}{3}}$

Calculator Check: $1.7321 > 1.4422$

**b.** $\sqrt{7} + \sqrt{18} \boxed{>} \sqrt{7 + 18}$

Calculator Check: $6.8884 > 5$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**141. a.** $\dfrac{ab}{a^2+ab+b^2}+\left(\dfrac{ac-ad-bc+bd}{ac-ad+bc-bd}\div\dfrac{a^3-b^3}{a^3+b^3}\right)=\dfrac{ab}{a^2+ab+b^2}+\left(\dfrac{a(c-d)-b(c-d)}{a(c-d)+b(c-d)}\cdot\dfrac{a^3+b^3}{a^3-b^3}\right)$

$=\dfrac{ab}{a^2+ab+b^2}+\left(\dfrac{\cancel{(c-d)}\cancel{(a-b)}}{\cancel{(c-d)}\cancel{(a+b)}}\cdot\dfrac{\cancel{(a+b)}\left(a^2-ab+b^2\right)}{\cancel{(a-b)}\left(a^2+ab+b^2\right)}\right)=\dfrac{ab}{a^2+ab+b^2}+\dfrac{a^2-ab+b^2}{a^2+ab+b^2}$

$=\dfrac{ab+a^2-ab+b^2}{a^2+ab+b^2}=\dfrac{a^2+b^2}{a^2+ab+b^2}$

Her son is 8 years old.

**b.** Son's portion:

$$\frac{8^{-\frac{4}{3}}+2^{-2}}{16^{-\frac{3}{4}}+2^{-1}}=\frac{\dfrac{1}{\left(\sqrt[3]{8}\right)^4}+\dfrac{1}{2^2}}{\dfrac{1}{\left(\sqrt[4]{16}\right)^3}+\dfrac{1}{2}}$$

$$=\frac{\dfrac{1}{2^4}+\dfrac{1}{4}}{\dfrac{1}{2^3}+\dfrac{1}{2}}$$

$$=\frac{\dfrac{1}{16}+\dfrac{1}{4}}{\dfrac{1}{8}+\dfrac{1}{2}}$$

$$=\frac{\dfrac{5}{16}}{\dfrac{5}{8}}$$

$$=\frac{8}{16}$$

$$=\frac{1}{2}$$

Mom's portion:

$\dfrac{1}{2}\left(1-\dfrac{1}{2}\right)=\dfrac{1}{2}\left(\dfrac{1}{2}\right)=\dfrac{1}{4}$

**142.** $(2x^3y^2)(5x^4y^7)=10x^7y^9$

**143.** $2x^4(8x^4+3x)=2x^4(8x^4)+2x^4(3x)$

$=16x^8+6x^5$

**144.** $2x(x^2+4x+5)+3(x^2+4x+5)$

$=2x^3+8x^2+10x+3x^2+12x+15$

$=2x^3+8x^2+3x^2+10x+12x+15$

$=2x^3+11x^2+22x+15$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**Section P.4**

**Check Point Exercises**

**1. a.** $(-17x^3+4x^2-11x-5)+(16x^3-3x^2+3x-15)$

$=(-17x^3+16x^3)+(4x^2-3x^2)+(-11x+3x)+(-5-15)$

$=-x^3+x^2-8x-20$

**b.** $(13x^2-9x^2-7x+1)-(-7x^3+2x^2-5x+9)$

$=(13x^3-9x^2-7x+1)+(7x^3-2x^2+5x-9)$

$=(13x^3+7x^3)+(-9x^2-2x^2)+(-7x+5x)+(1-9)$

$=20x^3-11x^2-2x-8$

**2.** $(5x-2)(3x^2-5x+4)$

$=5x(3x^2-5x+4)-2(3x^2-5x+4)$

$=5x\cdot 3x^2-5x\cdot 5x+5x\cdot 4-2\cdot 3x^2+2\cdot 5x-2\cdot 4$

$=15x^3-25x^2+20x-6x^2+10x-8$

$=15x^3-31x^2+30x-8$

**3.** $(7x-5)(4x-3)=7x\cdot 4x+7x(-3)+(-5)4x+(-5)(-3)$

$=28x^2-21x-20x+15$

$=28x^2-41x+15$

**4. a.** Use the special-product formula shown.

$(A+B)(A-B)=A^2-B^2$

$(7x+8)(7x-8)=(7x)^2-(8)^2$

$=49x^2-64$

**b.** Use the special-product formula shown.

$(A+B)(A-B)=A^2-B^2$

$(2y^3-5)(2y^3+5)=(2y^3+5)(2y^3-5)$

$=(2y^3)^2-(5)^2$

$=4y^6-25$

**5. a.** Use the special-product formula shown.

$(A+B)^2=A^2+2AB+B^2$

$(x+10)^2=x^2+2(x)(10)+10^2$

$=x^2+20x+100$

**b.** Use the special-product formula shown.

$(A+B)^2=A^2+2AB+B^2$

$(5x+4)^2=(5x)^2+2(5x)(4)+4^2$

$=25x^2+40x+16$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**6.** **a.** Use the special-product formula shown.

$$(A-B)^2 = A^2 - 2AB + B^2$$
$$(x-9)^2 = x^2 - 2(x)(9) + 9^2$$
$$= x^2 - 18x + 81$$

**b.** Use the special-product formula shown.

$$(A-B)^2 = A^2 - 2AB + B^2$$
$$(7x-3)^2 = (7x)^2 - 2(7x)(3) + 3^2$$
$$= 49x^2 - 42x + 9$$

**7.**
$$(x^3 - 4x^2y + 5xy^2 - y^3) - (x^3 - 6x^2y + y^3)$$
$$= (x^3 - 4x^2y + 5xy^2 - y^3) + (-x^3 + 6x^2y - y^3)$$
$$= (x^3 - x^3) + (-4x^2y + 6x^2y) + (5xy^2) + (-y^3 - y^3)$$
$$= 2x^2y + 5xy^2 - 2y^3$$

**8.** **a.**
$$(7x-6y)(3x-y) = (7x)(3x) + (7x)(-y) + (-6y)(3x) + (-6y)(-y)$$
$$= 21x^2 - 7xy - 18xy + 6y^2$$
$$= 21x^2 - 25xy + 6y^2$$

**b.**
$$(2x+4y)^2 = (2x)^2 + 2(2x)(4y) + (4y)^2$$
$$= 4x^2 + 16xy + 16y^2$$

**Concept and Vocabulary Check P.4**

**C1.** whole

**C2.** standard

**C3.** monomial

**C4.** binomial

**C5.** trinomial

**C6.** $n$

**C7.** like;

**C8.** distributive; $4x^3 - 8x^2 + 6$; $7x^3$

**C9.** $5x$; 3; like

**C10.** $3x^2$; $5x$; $21x$; 35

**C11.** $A^2 - B^2$; minus

**C12.** $A^2 + 2AB + B^2$; squared; product of the terms; squared

**C13.** $A^2 - 2AB + B^2$; minus; product of the terms; plus

**C14.** $n+m$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**Exercise Set P.4**

**1.** yes; $2x+3x^2-5=3x^2+2x-5$

**2.** no; The term $3x^{-1}$ does not have a whole number exponent.

**3.** no; The form of a polynomial involves addition and subtraction, not division.

**4.** yes; $x^2-x^3+x^4-5=x^4-x^3+x^2-5$

**5.** $3x^2$ has degree 2
$-5x$ has degree 1
4 has degree 0
$3x^2-5x+4$ has degree 2.

**6.** $-4x^3$ has degree 3
$7x^2$ has degree 2
$-11$ has degree 0
$-4x^3+7x^2-11$ has degree 3.

**7.** $x^2$ has degree 2
$-4x^3$ has degree 3
$9x$ has degree 1
$-12x^4$ has degree 4
63 has degree 0
$x^2-4x^3+9x-12x^4+63$ has degree 4.

**8.** $x^2$ has degree 2
$-8x^3$ has degree 3
$15x^4$ has degree 4
91 has degree 0
$x^2-8x^3+15x^4+91$ has degree 4.

**9.** $(-6x^3+5x^2-8x+9)+(17x^3+2x^2-4x-13)=(-6x^3+17x^3)+(5x^2+2x^2)+(-8x-4x)+(9-13)$
$=11x^3+7x^2-12x-4$
The degree is 3.

**10.** $(-7x^3+6x^2-11x+13)+(19x^3-11x^2+7x-17)=(-7x^3+19x^3)+(6x^2-11x^2)+(-11x+7x)+(13-17)$
$=12x^3-5x^2-4x-4$
The degree is 3.

**11.** $(17x^3-5x^2+4x-3)-(5x^3-9x^2-8x+11)=(17x^3-5x^2+4x-3)+(-5x^3+9x^2+8x-11)$
$=(17x^3-5x^3)+(-5x^2+9x^2)+(4x+8x)+(-3-11)$
$=12x^3+4x^2+12x-14$
The degree is 3.

Copyright © 2022 Pearson Education, Inc.


Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**12.** $(18x^4-2x^3-7x+8)-(9x^4-6x^3-5x+7)=(18x^4-2x^3-7x+8)+(-9x^4+6x^3+5x-7)$
$=(18x^4-9x^4)+(-2x^3+6x^3)+(-7x+5x)+(8-7)$
$=9x^4+4x^3-2x+1$

The degree is 4.

**13.** $(5x^2-7x-8)+(2x^2-3x+7)-(x^2-4x-3)=(5x^2-7x-8)+(2x^2-3x+7)+(-x^2+4x+3)$
$=(5x^2+2x^2-x^2)+(-7x-3x+4x)+(-8+7+3)$
$=6x^2-6x+2$

The degree is 2.

**14.** $(8x^2+7x-5)-(3x^2-4x)-(-6x^3-5x^2+3)=(8x^2+7x-5)+(-3x^2+4x)+(6x^3+5x^2-3)$
$=6x^3+(8x^2-3x^2+5x^2)+(7x+4x)+(-5-3)$
$=6x^3+10x^2+11x-8$

The degree is 3.

**15.** $(x+1)(x^2-x+1)=x(x^2)-x\cdot x+x\cdot 1+1(x^2)-1\cdot x+1\cdot 1$
$=x^3-x^2+x+x^2-x+1$
$=x^3+1$

**16.** $(x+5)\left(x^2-5x+25\right)=x(x^2)-x(5x)+x(25)+5(x^2)-5(5x)+5(25)$
$=x^3-5x^2+25x+5x^2-25x+125)$
$=x^3+125$

**17.** $(2x-3)(x^2-3x+5)=(2x)(x^2)+(2x)(-3x)+(2x)(5)+(-3)(x^2)+(-3)(-3x)+(-3)(5)$
$=2x^3-6x^2+10x-3x^2+9x-15$
$=2x^3-9x^2+19x-15$

**18.** $(2x-1)(x^2-4x+3)=(2x)(x^2)+(2x)(-4x)+(2x)(3)+(-1)(x^2)+(-1)(-4x)+(-1)(3)$
$=2x^3-8x^2+6x-x^2+4x-3$
$=2x^3-9x^2+10x-3$

**19.** $(x+7)(x+3)=x^2+3x+7x+21=x^2+10x+21$

**20.** $(x+8)(x+5)=x^2+5x+8x+40=x^2+13x+40$

**21.** $(x-5)(x+3)=x^2+3x-5x-15=x^2-2x-15$

**22.** $(x-1)(x+2)=x^2+2x-x-2=x^2+x-2$

**23.** $(3x+5)(2x+1)=(3x)(2x)+3x(1)+5(2x)+5=6x^2+3x+10x+5=6x^2+13x+5$

**24.** $(7x+4)(3x+1)=(7x)(3x)+7x(1)+4(3x)+4(1)=21x^2+7x+12x+4=21x^2+19x+4$

**25.** $(2x-3)(5x+3)=(2x)(5x)+(2x)(3)+(-3)(5x)+(-3)(3)=10x^2+6x-15x-9=10x^2-9x-9$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**26.** $(2x-5)(7x+2)=(2x)(7x)+(2x)(2)+(-5)(7x)+(-5)(2)=14x^2+4x-35x-10=14x^2-31x-10$

**27.** $(5x^2-4)(3x^2-7)=(5x^2)(3x^2)+(5x^2)(-7)+(-4)(3x^2)+(-4)(-7)=15x^4-35x^2-12x^2+28=15x^4-47x^2+28$

**28.** $(7x^2-2)(3x^2-5)=(7x^2)(3x^2)+(7x^2)(-5)+(-2)(3x^2)+(-2)(-5)=21x^4-35x^2-6x^2+10=21x^4-41x^2+10$

**29.** $\left(8x^3+3\right)\left(x^2-5\right)=\left(8x^3\right)\left(x^2\right)+\left(8x^3\right)(-5)+(3)\left(x^2\right)+(3)(-5)=8x^5-40x^3+3x^2-15$

**30.** $\left(7x^3+5\right)\left(x^2-2\right)=\left(7x^3\right)\left(x^2\right)+\left(7x^3\right)(-2)+(5)\left(x^2\right)+(5)(-2)=7x^5-14x^3+5x^2-10$

**31.** $(x+3)(x-3)=x^2-3^2=x^2-9$

**32.** $(x+5)(x-5)=x^2-5^2=x^2-25$

**33.** $(3x+2)(3x-2)=(3x)^2-2^2=9x^2-4$

**34.** $(2x+5)(2x-5)=(2x)^2-5^2=4x^2-25$

**35.** $(5-7x)(5+7x)=5^2-(7x)^2=25-49x^2$

**36.** $(4-3x)(4+3x)=4^2-(3x)^2=16-9x^2$

**37.** $(4x^2+5x)(4x^2-5x)=(4x^2)^2-(5x)^2=16x^4-25x^2$

**38.** $(3x^2+4x)(3x^2-4x)=(3x^2)^2-(4x)^2=9x^4-16x^2$

**39.** $\left(1-y^5\right)\left(1+y^5\right)=(1)^2-\left(y^5\right)^2=1-y^{10}$

**40.** $\left(2-y^5\right)\left(2+y^5\right)=(2)^2-\left(y^5\right)^2=4-y^{10}$

**41.** $(x+2)^2=x^2+2\cdot x\cdot 2+2^2=x^2+4x+4$

**42.** $(x+5)^2=x^2+2\cdot x\cdot 5+5^2=x^2+10x+25$

**43.** $(2x+3)^2=(2x)^2+2(2x)(3)+3^2=4x^2+12x+9$

**44.** $(3x+2)^2=(3x)^2+2(3x)(2)+2^2=9x^2+12x+4$

**45.** $(x-3)^2=x^2-2\cdot x\cdot 3+3^2=x^2-6x+9$

**46.** $(x-4)^2=x^2-2\cdot x\cdot 4+4^2=x^2-8x+16$

**47.** $(4x^2-1)^2=(4x^2)^2-2(4x^2)(1)+1^2=16x^4-8x^2+1$

**48.** $(5x^2-3)^2=(5x^2)^2-2(5x^2)(3)+3^2=25x^4-30x^2+9$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**49.** $(7-2x)^2 = 7^2 - 2(7)(2x) + (2x)^2 = 49 - 28x + 4x^2 = 4x^2 - 28x + 49$

**50.** $(9-5x)^2 = 9^2 - 2(9)(5x) + (5x)^2 = 81 - 90x + 25x^2$ or $25x^2 - 90x + 81$

**51.** $(x+1)^3 = x^3 + 3 \cdot x^2 \cdot 1 + 3x \cdot 1^2 + 1^3 = x^3 + 3x^2 + 3x + 1$

**52.** $(x+2)^3 = x^3 + 3 \cdot x^2 \cdot 2 + 3 \cdot x \cdot 2^2 + 2^3 = x^3 + 6x^2 + 12x + 8$

**53.** $(2x+3)^3 = (2x)^3 + 3 \cdot (2x)^2 \cdot 3 + 3(2x) \cdot 3^2 + 3^3 = 8x^3 + 36x^2 + 54x + 27$

**54.** $(3x+4)^3 = (3x)^3 + 3(3x)^2 \cdot 4 + 3(3x) \cdot 4^2 + 4^3 = 27x^3 + 108x^2 + 144x + 64$

**55.** $(x-3)^3 = x^3 - 3 \cdot x^3 \cdot 3 + 3 \cdot x \cdot 3^2 - 3^3 = x^3 - 9x^2 + 27x - 27$

**56.** $(x-1)^3 = x^3 - 3x^2 \cdot 1 + 3x \cdot 1^2 - 1^3 = x^3 - 3x^2 + 3x - 1$

**57.** $(3x-4)^3 = (3x)^3 - 3(3x)^2 \cdot 4 + 3(3x) \cdot 4^2 - 4^3 = 27x^3 - 108x^2 + 144x - 64$

**58.** $(2x-3)^3 = (2x)^3 - 3(2x)^2 \cdot 3 + 3(2x) \cdot 3^2 - 3^3 = 8x^3 - 36x^2 + 54x - 27$

**59.** $(5x^2y - 3xy) + (2x^2y - xy) = (5x^2y + 2x^2y) + (-3xy - xy)$
$= (5+2)x^2y + (-3-1)xy$
$= 7x^2y - 4xy$ is of degree 3.

**60.** $(-2x^2y + xy) + (4x^2y + 7xy) = (-2x^2y + 4x^2y) + (xy + 7xy)$
$= (-2+4)x^2y + (1+7)xy$
$= 2x^2y + 8xy$ is of degree 3.

**61.** $(4x^2y + 8xy + 11) + (-2x^2y + 5xy + 2) = (4x^2y - 2x^2y) + (8xy + 5xy) + (11+2)$
$= (4-2)x^2y + (8+5)xy + 13$
$= 2x^2y + 13xy + 13$ is of degree 3.

**62.** $(7x^4y^2 - 5x^2y^2 + 3xy) + (-18x^4y^2 - 6x^2y^2 - xy) = (7x^4y^2 - 18x^4y^2) + (-5x^2y^2 - 6x^2y^2) + (3xy - xy)$
$= (7-18)x^4y^2 + (-5-6)x^2y^2 + (3-1)xy$
$= -11x^4y^2 - 11x^2y^2 + 2xy$ is of degree 6.

**63.** $(x^3 + 7xy - 5y^2) - (6x^3 - xy + 4y^2) = (x^3 + 7xy - 5y^2)$
$= (x^3 - 6x^3) + (7xy + xy) + (-5y^2 - 4y^2)$
$= (1-6)x^3 + (7+1)xy + (-5-4)y^2$
$= -5x^3 + 8xy - 9y^2$ is of degree 3.


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**64.** $(x^4-7xy-5y^3)-(6x^4-3xy+4y^3)=(x^4-7xy-5y^3)+(-6x^4+3xy-4y^3)$
$=(x^4-6x^4)+(-7xy+3xy)+(-5y^3-4y^3)$
$=(1-6)x^4+(-7+3)xy+(-5-4)y^3$
$=-5x^4-4xy-9y^3$ is of degree 4.

**65.** $(3x^4y^2+5x^3y-3y)-(2x^4y^2-3x^3y-4y+6x)=(3x^4y^2+5x^3y-3y)+(-2x^4y^2+3x^3y+4y-6x)$
$=(3x^4y^2-2x^4y^2)+(5x^3y+3x^3y)+(-3y+4y)-6x$
$=(3-2)x^4y^2+(5+3)x^3y+(-3+4)y-6x$
$=x^4y^2+8x^3y+y-6x$ is of degree 6.

**66.** $(5x^4y^2+6x^3y-7y)-(3x^4y^2-5x^3y-6y+8x)=(5x^4y^2+6x^3y-7y)+(-3x^4y^2+5x^3y+6y-8x)$
$=(5x^4y^2-3x^4y^2)+(6x^3y+5x^3y)+(-7y+6y)-8x$
$=(5-3)x^4y^2+(6+5)x^3y+(-7+6)y-8x$
$=2x^4y^2+11x^3y-y-8x$ is of degree 6.

**67.** $(x+5y)(7x+3y)=x(7x)+x(3y)+(5y)(7x)+(5y)(3y)$
$=7x^2+3xy+35xy+15y^2$
$=7x^2+38xy+15y^2$

**68.** $(x+9y)(6x+7y)=x(6x)+x(7y)+(9y)(6x)+(9y)(7y)$
$=6x^2+7xy+54xy+63y^2$
$=6x^2+61xy+63y^2$

**69.** $(x-3y)(2x+7y)=x(2x)+x(7y)+(-3y)(2x)+(-3y)(7y)$
$=2x^2+7xy-6xy-21y^2$
$=2x^2+xy-21y^2$

**70.** $(3x-y)(2x+5y)=(3x)(2x)+(3x)(5y)+(-y)(2x)+(-y)(5y)$
$=6x^2+15xy-2xy-5y^2$
$=6x^2+13xy-5y^2$

**71.** $(3xy-1)(5xy+2)=(3xy)(5xy)+(3xy)(2)+(-1)(5xy)+(-1)(2)$
$=15x^2y^2+6xy-5xy-2$
$=15x^2y^2+xy-2$

**72.** $(7x^2y+1)(2x^2y-3)=(7x^2y)(2x^2y)+(7x^2y)(-3)+(1)2x^2y+(1)(-3)$
$=14x^4y^2-21x^2y+2x^2y-3$
$=14x^4y^2-19x^2y-3$

**73.** $(7x+5y)^2=(7x)^2+2(7x)(5y)+(5y)^2=49x^2+70xy+25y^2$

**74.** $(9x+7y)^2=(9x)^2+2(9x)(7y)+(7y)^2=81x^2+126xy+49y^2$

Copyright © 2022 Pearson Education, Inc.


Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**75.** $(x^2y^2-3)^2=(x^2y^2)^2-2(x^2y^2)(3)+3^2=x^4y^4-6x^2y^2+9$

**76.** $(x^2y^2-5)^2=(x^2y^2)^2-2(x^2y^2)(5)+5^2=x^4y^4-10x^2y^2+25$

**77.** $$\begin{aligned}(x-y)(x^2+xy+y^2)&=x(x^2)+x(xy)+x(y^2)+(-y)(x^2)+(-y)(xy)+(-y)(y^2)\\&=x^3+x^2y+xy^2-x^2y-xy^2-y^3\\&=x^3-y^3\end{aligned}$$

**78.** $$\begin{aligned}(x+y)(x^2-xy+y^2)&=x(x^2)+x(-xy)+x(y^2)+y(x^2)+y(-xy)+y(y^2)\\&=x^3-x^2y+xy^2+x^2y-xy^2+y^3\\&=x^3+y^3\end{aligned}$$

**79.** $(3x+5y)(3x-5y)=(3x)^2-(5y)^2=9x^2-25y^2$

**80.** $(7x+3y)(7x-3y)=(7x)^2-(3y)^2=49x^2-9y^2$

**81.** $\left(7xy^2-10y\right)\left(7xy^2+10y\right)=\left(7xy^2\right)^2-\left(10y\right)^2=49x^2y^4-100y^2$

**82.** $\left(3xy^2-4y\right)\left(3xy^2+4y\right)=\left(3xy^2\right)^2-\left(4y\right)^2=9x^2y^4-16y^2$

**83.** $$\begin{aligned}(3x+4y)^2-(3x-4y)^2&=\left[(3x)^2+2(3x)(4y)+(4y)^2\right]-\left[(3x)^2-2(3x)(4y)+(4y)^2\right]\\&=\left(9x^2+24xy+16y^2\right)-\left(9x^2-24xy+16y^2\right)\\&=9x^2+24xy+16y^2-9x^2+24xy-16y^2\\&=48xy\end{aligned}$$

**84.** $$\begin{aligned}(5x+2y)^2-(5x-2y)^2&=\left[(5x)^2+2(5x)(2y)+(2y)^2\right]-\left[(5x)^2-2(5x)(2y)+(2y)^2\right]\\&=\left(25x^2+20xy+4y^2\right)-\left(25x^2-20xy+4y^2\right)\\&=25x^2+20xy+4y^2-25x^2+20xy-4y^2\\&=40xy\end{aligned}$$

**85.** $$\begin{aligned}&(5x-7)(3x-2)-(4x-5)(6x-1)\\&=\left[15x^2-10x-21x+14\right]-\left[24x^2-4x-30x+5\right]\\&=\left(15x^2-31x+14\right)-\left(24x^2-34x+5\right)\\&=15x^2-31x+14-24x^2+34x-5\\&=-9x^2+3x+9\end{aligned}$$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**86.** $(3x+5)(2x-9)-(7x-2)(x-1)$

$=\left(6x^2-27x+10x-45\right)-\left(7x^2-7x-2x+2\right)$

$=\left(6x^2-17x-45\right)-\left(7x^2-9x+2\right)$

$=6x^2-17x-45-7x^2+9x-2$

$=-x^2-8x-47$

**87.** $(2x+5)(2x-5)\left(4x^2+25\right)$

$=\left[(2x)^2-5^2\right]\left(4x^2+25\right)$

$=\left(4x^2-25\right)\left(4x^2+25\right)$

$=\left(4x^2\right)^2-(25)^2$

$=16x^4-625$

**88.** $(3x+4)(3x-4)\left(9x^2+16\right)$

$=\left[(3x)^2-4^2\right]\left(9x^2+16\right)$

$=\left(9x^2-16\right)\left(9x^2+16\right)$

$=\left(9x^2\right)^2-(16)^2$

$=81x^4-256$

**89.** $\dfrac{(2x-7)^5}{(2x-7)^3}=(2x-7)^{5-3}$

$=(2x-7)^2$

$=(2x)^2-2(2x)(7)+(7)^2$

$=4x^2-28x+49$

**90.** $\dfrac{(5x-3)^6}{(5x-3)^4}=(5x-3)^{6-4}$

$=(5x-3)^2$

$=(5x)^2-2(5x)(3)+(3)^2$

$=25x^2-30x+9$

**91.** **a.** $S=0.2x^3-1.5x^2+3.4x+25+\left(0.1x^3-1.3x^2+3.3x+5\right)$

$S=0.2x^3-1.5x^2+3.4x+25+0.1x^3-1.3x^2+3.3x+5$

$S=0.3x^3-2.8x^2+6.7x+30$

Copyright © 2022 Pearson Education, Inc.



Algebra and Trigonometry 7th edition Blitzer Solutions Manual

**b.** $S = 0.3x^3 - 2.8x^2 + 6.7x + 30$

$S = 0.3(5)^3 - 2.8(5)^2 + 6.7(5) + 30$

$S = 31$

The model gives a score of 31 for the group in the 45-54 age range which is the same as the score displayed by the bar graph.

**92.** **a.** $S = 0.1x^3 - 1.2x^2 + 2.4x + 17 + \left(0.2x^3 - 1.6x^2 + 4.3x + 13\right)$

$S = 0.1x^3 - 1.2x^2 + 2.4x + 17 + 0.2x^3 - 1.6x^2 + 4.3x + 13$

$S = 0.3x^3 - 2.8x^2 + 6.7x + 30$

**b.** $S = 0.3x^3 - 2.8x^2 + 6.7x + 30$

$S = 0.3(6)^3 - 2.8(6)^2 + 6.7(6) + 30$

$S = 34.2$

The model gives a score of 34.2 for the group in the 55-64 age range. This overestimates the score shown on the bar graph by 0.2.

**93.** $x(8-2x)(10-2x) = x\left(80 - 36x + 4x^2\right)$

$= 80x - 36x^2 + 4x^3$

$= 4x^3 - 36x^2 + 80x$

**94.** $x(8-2x)(5-2x) = x\left(40 - 26x + 4x^2\right)$

$= 40x - 26x^2 + 4x^3$

$= 4x^3 - 26x^2 + 40x$

**95.** $(x+9)(x+3) - (x+5)(x+1)$

$= x^2 + 12x + 27 - \left(x^2 + 6x + 5\right)$

$= x^2 + 12x + 27 - x^2 - 6x - 5$

$= 6x + 22$

**96.** $(x+4)(x+3) - (x+2)(x+1)$

$= x^2 + 7x + 12 - \left(x^2 + 3x + 2\right)$

$= x^2 + 7x + 12 - x^2 - 3x - 2$

$= 4x + 10$

**97. – 102.** Answers will vary.

**103.** makes sense

**104.** does not make sense; Explanations will vary. Sample explanation: FOIL is used to multiply two binomials.

**105.** makes sense

**106.** makes sense, although answers may vary

**107.** false; Changes to make the statement true will vary. A sample change is: $(3x^3 + 2)(3x^3 - 2) = 9x^6 - 4$


Copyright © 2022 Pearson Education, Inc

Algebra and Trigonometry 7th edition Blitzer Solutions Manual