# *Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual*

SKU: 9780139078743-SOLUTIONS

# CHAPTER 2: Analytic Functions

## EXERCISES 2.1: Functions of a Complex Variable

1. a. $w = (3x^2 - 3y^2 + 5x + 1) + i(6xy + 5y + 1)$

   b. $w = \dfrac{x}{x^2+y^2} + i\left(-\dfrac{y}{x^2+y^2}\right)$

   c. $w = \dfrac{1}{z-i} = \dfrac{x}{x^2+(y-1)^2} + i\dfrac{-y+1}{x^2+(y-1)^2}$

   d. $w = \dfrac{2x^2-2y^2+3}{\sqrt{(x-1)^2+y^2}} + i\dfrac{4xy}{\sqrt{(x-1)^2+y^2}}$

   e. $w = e^{3x}\cos 3y + ie^{3x}\sin 3y$

   f. $w = (e^x + e^{-x})\cos y + i(e^x - e^{-x})\sin y$
   $= 2\cosh x \cos y + i2\sinh x \sin y$

2. a. $\mathbf{C}$

   b. $\mathbf{C} \setminus \{0\}$

   c. $\mathbf{C} \setminus \{i, -i\}$

   d. $\mathbf{C} \setminus \{1\}$

   e. $\mathbf{C}$

   f. $\mathbf{C}$

3. a. $\operatorname{Re} w > 5$

   b. $\operatorname{Im} w \geq 0$

   c. $|w| \geq 1$

   d. The intersection of $|w| < 2$ and $-\pi < \operatorname{Arg} w < \pi/2$

4. a. Taking $\theta$ from 0 to $2\pi$, the points $z = re^{i\theta}$ traverse the circle $|z| = r$ exactly once in the counterclockwise direction. For the same values of $\theta$ the points $w = \dfrac{1}{re^{i\theta}} = \dfrac{1}{r}e^{-i\theta}$ traverse the circle $|w| = \dfrac{1}{r}$ exactly once in the clockwise direction, hence the mapping is onto.

   b. For $z = re^{i\theta_0}$ on the ray $\operatorname{Arg} z = \theta_0$, $w = \dfrac{1}{re^{i\theta_0}} = \dfrac{1}{r}e^{-i\theta_0}$ is on the ray $\operatorname{Arg} w = -\theta_0$. Taking values $0 < r < \infty$ shows that this mapping goes onto the ray $\operatorname{Arg} w = -\theta_0$.



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

4 (c) $|z-1| = 1$ $2\pi > \theta \geq 0 \Rightarrow z = 1 + e^{i\theta}$. $F(z) = 1/z. = 1/(1+e^{i\theta})$
$= (1+e^{-i\theta})/\{2(1+\cos\theta)\} = \frac{1}{2} - i(\frac{1}{2})\sin\theta/(1+\cos\theta)$
which is a vertical line at $x = \frac{1}{2}$.

5. a. domain: $\mathbf{C}$
range: $\mathbf{C} \setminus \{0\}$

b. $f(-z) = e^{-z} = \dfrac{1}{e^z} = \dfrac{1}{f(z)}$

c. circle $|w| = e$

d. ray $\operatorname{Arg} w = \pi/4$


e. infinite sector $0 \leq \operatorname{Arg} w \leq \pi/4$

6. a. $J\left(\dfrac{1}{z}\right) = \dfrac{1}{2}\left(\dfrac{1}{z} + \dfrac{1}{1/z}\right) = \dfrac{1}{2}\left(z + \dfrac{1}{z}\right) = J(z)$


b. For $z = e^{i\theta}$ on the unit circle $|z| = 1$, $J(z) = \dfrac{1}{2}\left(e^{i\theta} + \dfrac{1}{e^{i\theta}}\right) = \cos\theta$. For all values of $\theta$, this ranges over the real interval $[-1, 1]$.

c. For $z = re^{i\theta}$ on the circle $|z| = r$, $J(z) = \dfrac{1}{2}\left(re^{i\theta} + \dfrac{1}{re^{i\theta}}\right) = \dfrac{1}{2}\left(r + \dfrac{1}{r}\right)\cos\theta + i\dfrac{1}{2}\left(r - \dfrac{1}{r}\right)\sin\theta$. Setting $u$ and $v$ equal to the real and imaginary parts of this expression, respectively, one gets a pair of parametric equations that are equivalent to the ellipse $\dfrac{u^2}{[\frac{1}{2}(r+\frac{1}{r})]^2} + \dfrac{v^2}{[\frac{1}{2}(r-\frac{1}{r})]^2} = 1$, which has foci at $\pm 1$.

7. a.





b.







Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual


c.





8. a.




b.





c.







Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual


9. a.





b.





10. a. translate by $i$, rotate $\pi/4$





b. reduce by $1/2$, rotate $\pi/4$




c. translate by $i$, reduce by $1/2$





d. reduce by $1/2$, rotate $\pi/4$, translate by $i$







Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual


11. a. translate by $-3$, rotate $-\pi/2$





b. magnify by 2, rotate $-\pi/2$




c. translate by $-3$, magnify by 2




d. magnify by 2, rotate $-\pi/2$, translate by $-3$




12. Let $a = \rho e^{i\phi}$, $F(z) = \rho z$, $G(z) = e^{i\phi} z$, and $H(z) = z + b$. Then $H(G(F(z))) = az + b$.

13. (a) $w = u + iv = z^2 = (1 + iy)^2 = 1 - y^2 + i2y$
$u = 1 - y^2$, $v = 2y \Rightarrow y = v/2 \Rightarrow u = 1 - v^2/4$ a parabola in the w-plane.

(b) $w = u + iv = z^2 = (x + iy)^2 = (x + i/x)^2 = x^2 - 1/x^2 + 2i$
$u = x^2 - 1/x^2$, $v = 2$ a straight line in the w-plane.

(c) $w = u + iv = z^2 = (1 + e^{i\theta})^2 = (1 + 2e^{i\theta} + e^{i2\theta}) = (e^{-i\theta} + 2 + e^{i\theta})e^{i\theta}$
$= (2 + 2\cos\theta)e^{i\theta} = 2(1 + \cos\theta)e^{i\theta}$ a cardioid in the w-plane.

14. (a) $x_1 = 2x/(|z|^2 + 1)$, $x_2 = 2y/(|z|^2 + 1)$, $x_3 = (|z|^2 - 1)/(|z|^2 + 1)$

$w = e^{i\varphi} z = x\cos\varphi - y\sin\varphi + i(x\sin\varphi + y\cos\varphi)$, $|w| = |z|$

$\underline{x}_1 = (x\cos\varphi - y\sin\varphi)/(|z|^2 + 1)$, $\underline{x}_2 = (x\sin\varphi + y\cos\varphi)/(|z|^2 + 1)$, $\underline{x}_3 = x_3$

$\underline{x}_1 = (x_1\cos\varphi - x_2\sin\varphi)$, $\underline{x}_2 = (x_1\sin\varphi + x_2\cos\varphi)$, $\underline{x}_3 = x_3$ which corresponds to a rotation of an angle $\varphi$ about the $x_3$ axis.

(b) $w = -1/z$. $|w| = 1/|z|$. $w = -1/(x+iy) = -x/|z| + iy/|z|$
$\underline{x}_1 = -x_1$, $\underline{x}_2 = x_2$, $\underline{x}_3 = -x_3$ so that $(\underline{x}_1, \underline{x}_2, \underline{x}_3)$ is obtained from $(x_1, x_2, x_3)$ by a $180^\circ$ rotation about the $x_2$ axis.

15. $w = (1+z)/(1 - z) = (1+x + iy)/(1 - x - iy) = (1-|z|^2 + i2y)/(1 - 2x + |z|^2)$
$|w|^2 = (1 + 2x + |z|^2)/(1 - 2x + |z|^2)$.
$(\underline{x}_1, \underline{x}_2, \underline{x}_3) = (-x_3, x_2, x_1)$ so that $(\underline{x}_1, \underline{x}_2, \underline{x}_3)$ is obtained by a $90^\circ$ counterclockwise rotation about the $x_2$ axis.




Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

16. $w = (1 - iz)/(1 + iz) = (1- ix + y)/(1 + ix - y) = (1-|z|^2 + i2x)/(1- 2y + |z|^2)$
$|w|^2 = (1 + 2y + |z|^2)/(1- 2y + |z|^2)$.
$(\underline{x}_1, \underline{x}_2, \underline{x}_3) = (-x_3, -x_1, x_2)$ so that $(\underline{x}_1, \underline{x}_2, \underline{x}_3)$ is obtained as a $90^o$ counterclockwise rotation about the $x_2$ axis followed by a $90^o$ counterclockwise rotation about the $x_3$axis.


17. Any circle or line in the z-plane corresponds to a line or circle on the stenographic projection onto he Riemann sphere. The function w=1/z rotates the Riemann sphere $180^o$ about the $x_1$ axis. Lines and circles on the rotated sphere project to lines and circles in the w-plane. As a result lines and circles in the z-plane map to lines and circles in the w-plane.

## EXERCISES 2.2: Limits and Continuity

1. The first five terms are, respectively, $\frac{i}{2}$, $-\frac{1}{4}$, $-\frac{i}{8}$, $\frac{1}{16}$, and $\frac{i}{32}$. The sequence converges to 0 in a spiral-like fashion.




2. $2i$, $-4$, $-8i$, 16, $32i$; divergent because terms grow in modulus without bound.






Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

3. If $\lim_{n\to\infty} z_n = z_0$, then for any $\varepsilon>0$, there is an integer N such that $|z_n - z_0|<\varepsilon$ for all $n>N$. For the same integer N we have $|x_n - x_0|<=|z_n - z_0|<\varepsilon$ and $|y_n - y_0|<=|z_n - z_0|<\varepsilon$ for all $n>N$. Therefore, $\lim_{n\to\infty} x_n = x_0$ and $\lim_{n\to\infty} y_n = y_0$.

If $\lim_{n\to\infty} x_n = x_0$ and $\lim_{n\to\infty} y_n = y_0$, then for any $\varepsilon_1 >0$ and $\varepsilon_2 >0$ there are integers $N_1$ and $N_2$ such $|x_n - x_0|<\varepsilon_1$ for all $n > N_1$ and $|y_n - y_0|<\varepsilon_2$ for all $n > N_2$. Given any $\varepsilon > 0$; let $\varepsilon_1 = \varepsilon/2$ and $\varepsilon_2 = \varepsilon/2$. Then $|z_n - z_0|<=\varepsilon\ |x_n - x_0| + |y_n - y_0| < \varepsilon_1 + \varepsilon_2 = \varepsilon$ for all $n >$ maximum$(N_1, N_2)$. Thus $\lim_{n\to\infty} z_n = z_0$.

4. If $z_n = x_n + iy_n \to z_0 = x_0 + iy_0$, then $x_n \to x_0$ and $y_n \to y_0$ (see Problem 3). $\underline{z}_n = x_n - iy_n \to x_0 - iy_0 = \underline{z}_0$.

If $\underline{z}_n = x_n - iy_n \to \underline{z}_0 = x_0 - iy_0$, then $x_n \to x_0$ and $y_n \to y_0$ (see Problem 3).. $z_n = x_n + iy_n \to x_0 + iy_0 = z_0$. Thus $z_n \to z_0$ if and only if $\underline{z}_n \to \underline{z}_0$.

5. $\lim_{n\to\infty} |z_n| = 0 \Longrightarrow$ **There exists an integer** $N$ **such that** $||z_n| - 0| = |z_n| < \varepsilon$ **whenever** $n > N. \Longrightarrow |z_n - 0| < \varepsilon$ **whenever** $n > N. \Longrightarrow \lim_{n\to\infty} z_n = 0$, **and conversely.**

6. $z_0^n \to 0$ **as** $n \to \infty$ **by problem 3, since the real-valued sequence** $|z_0^n| \to 0$ **as** $n \to \infty$. **On the other hand, if** $|z_0| > 1$, **then** $|z_0^n| \to \infty$ **as** $n \to \infty$ **so** $z_0^n$ **diverges.**

7. a. **converges to 0**
   b. **does not converge**
   c. **converges to** $\pi$
   d. **converges to** $2 + i$
   e. **converges to 0**
   f. **does not converge**

8. **Given** $\varepsilon > 0$, **choose** $\delta = \varepsilon/6$. **Then whenever** $0 < |z - (1 + i)| < \delta$,

$$|6z - 4 - (2 + 6i)| = 6|z - (1 + i)| < 6(\varepsilon/6) = \varepsilon$$

9. **Given** $\varepsilon > 0$, **choose** $\delta = \dfrac{\varepsilon}{1+\varepsilon}$. **Whenever** $0 < |z - (-i)| < \delta$ **notice that** $|z| > 1 - \delta$ **and**

$$\left|\frac{1}{z} - i\right| = \left|\left(-\frac{i}{z}\right)(i + z)\right| = \frac{1}{|z|}|z - (-i)| < \left(\frac{1}{1-\delta}\right)\delta = \varepsilon$$



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

10. **Given that $f$ and $g$ are continuous at $z_0$,**

$$\lim_{z\to z_0} f(z) \pm g(z) = \lim_{z\to z_0} f(z) \pm \lim_{z\to z_0} g(z) = f(z_0) \pm g(z_0)$$

$$\Longrightarrow f(z) \pm g(z) \text{ is continuous at } z_0.$$

$$\lim_{z\to z_0} f(z)g(z) = \lim_{z\to z_0} f(z) \lim_{z\to z_0} g(z) = f(z_0)g(z_0)$$

$$\Longrightarrow f(z)g(z) \text{ is continuous at } z_0.$$

$$\lim_{z\to z_0} \frac{f(z)}{g(z)} = \frac{\lim_{z\to z_0} f(z)}{\lim_{z\to z_0} g(z)} = \frac{f(z_0)}{g(z_0)}, \text{ provided } g(z_0) \neq 0$$

$$\Longrightarrow \frac{f(z)}{g(z)} \text{ is continuous at } z_0.$$

11. a. $-8i$

b. $-\frac{7}{2}i$

c. $6i$

d. $-1/2$

e. $2z_0$

f. $4\sqrt{2}$

12. Clearly $\operatorname{Arg} z$ is discontinuous at $z = 0$. Let $a > 0$ be any real number and consider the sequence

$$z_n = -a - i/n \quad n = 1, 2, \ldots, \text{ which converges to } -a.$$

For each $n$, $-\pi < \operatorname{Arg} z_n < -\pi/2$, but $\operatorname{Arg}(-a) = \pi$.

13. $\lim_{z\to z_0} f(z)$ exists for all $z \neq -1$; $f$ is continuous for all $z \neq 0, -1$; $f$ has a removable discontinuity at $z = 0$.

14. Let $z_0$ be any complex number. Given $\varepsilon > 0$ choose $\delta = \varepsilon$. Then whenever $|z - z_0| < \delta$,

$$|g(z) - g(z_0)| = |\overline{z} - \overline{z}_0| = |\overline{z - z_0}| = |z - z_0| < \varepsilon.$$

15. Given $\varepsilon > 0$ choose $\delta$ so that $|f(z) - f(z_0)| < \varepsilon$ whenever $|z - z_0| < \delta$. Then, whenever $|z - z_0| < \delta$:

a. $|\overline{f(z)} - \overline{f(z_0)}| = |\overline{f(z) - f(z_0)}| = |f(z) - f(z_0)| < \varepsilon$

b. $|\operatorname{Re} f(z) - \operatorname{Re} f(z_0)| = |\operatorname{Re}(f(z) - f(z_0))| \leq |f(z) - f(z_0)| < \varepsilon$

c. $|\operatorname{Im} f(z) - \operatorname{Im} f(z_0)| = |\operatorname{Im}(f(z) - f(z_0))| \leq |f(z) - f(z_0)| < \varepsilon$

d. $||f(z)| - |f(z_0)|| \leq |f(z) - f(z_0)| < \varepsilon$



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

16. Given $\varepsilon > 0$, choose $\delta_0 > 0$ such that $|f(g(z)) - f(g(z_0))| < \varepsilon$ whenever $|g(z) - g(z_0)| < \delta_0$. Now choose $\delta > 0$ such that $|g(z) - g(z_0)| < \delta_0$ whenever $|z - z_0| < \delta$. Then $|f(g(z)) - f(g(z_0))| < \varepsilon$ whenever $|z - z_0| < \delta$; hence $f(g(z))$ is continuous at $z_0$.

17. No: Observe that although $\frac{1}{n} \to 0$ and $\frac{i}{n} \to 0$ as $n \to \infty$, $f\left(\frac{1}{n}\right) \to 1 + 2i$ and $f\left(\frac{i}{n}\right) \to 2i$; thus $\lim_{z \to 0} f(z)$ does not exist.

18. If $\lim_{z \to z_0} f(z) = w_0$, then given $\varepsilon > 0$ there exists $\delta > 0$ such that $|f(z) - w_0| < \varepsilon$ for all $|z - z_0| < \delta$. Notice that
$|\underline{f(z)} - \underline{w_0}| = |\underline{f(z) - w_0}| = |f(z) - w_0| < \varepsilon$ for all $|z - z_0| < \delta$. So that $\lim_{z \to z_0} \underline{f(z)} = \underline{w_0}$.
$\lim_{x \to x_0, y \to y_0} \mu(x,y) = \lim_{z \to z_0} ((f(z) + \underline{f(z)})/2) = (w_0 + \underline{w_0})/2 = \mu_0$.
$\lim_{x \to x_0, y \to y_0} \upsilon(x,y) = \lim_{z \to z_0} ((f(z) - \underline{f(z)})/2i) = (w_0 - \underline{w_0})/2i = \upsilon_0$.
Thus, $\lim_{x \to x_0, y \to y_0} \mu(x,y) = \mu_0$ and $\lim_{x \to x_0, y \to y_0} \upsilon(x,y) = \upsilon_0$.

Conversely, if $\lim_{x \to x_0, y \to y_0} \mu(x,y) = \mu_0$ and $\lim_{x \to x_0, y \to y_0} \upsilon(x,y) = \upsilon_0$, then (by Theorem 1.) $\mu_0 + i\upsilon_0 = \lim_{x \to x_0, y \to y_0} \mu(x,y) + i \lim_{x \to x_0, y \to y_0} \upsilon(x,y) = \lim_{z \to z_0} ((f(z) + \underline{f(z)})/2) + \lim_{z \to z_0} ((f(z) - \underline{f(z)})/2) = \lim_{z \to z_0} f(z) = w_0$.
Also $\mu_0 - i\upsilon_0 = \lim_{x \to x_0, y \to y_0} \mu(x,y) - i \lim_{x \to x_0, y \to y_0} \upsilon(x,y) = \lim_{z \to z_0} ((f(z) + \underline{f(z)})/2) - \lim_{z \to z_0} ((f(z) - \underline{f(z)})/2) = \lim_{z \to z_0} f(z) = \underline{w_0}$.

Thus, $\lim_{z \to z_0} f(z) = w_0$.

19. $-\frac{1}{2} - i$, since $\lim_{\substack{x \to 1 \\ y \to -1}} \frac{x}{x^2 + 3y} = -\frac{1}{2}$ and $\lim_{\substack{x \to 1 \\ y \to -1}} xy = -1$.

20. For any $z_0$ in the complex plane,

$$\lim_{z \to z_0} e^z = \lim_{\substack{x \to x_0 \\ y \to y_0}} e^x \cos y + i \lim_{\substack{x \to x_0 \\ y \to y_0}} e^x \sin y = e^{x_0} \cos y_0 + i e^{x_0} \sin y_0 = e^{z_0}.$$

21. a. 1
b. 0
c. $-\pi/2 + i$
d. 1

21

22. By contradiction: Suppose $\lim_{z \to z_0} f(z) \neq w_0$. Then there is an $\varepsilon > 0$ for which there exists a sequence $\{z_n\}$ such that $|z_n - z_0| < \frac{1}{n}$ but $|f(z_n) - w_0| > \varepsilon$. For this sequence, $\lim_{n \to \infty} z_n = z_0$ but $\lim_{n \to \infty} f(z_n) \neq w_0$, contrary to hypothesis.



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

23. If $z_n \to \infty$, then for any M>0 there exist an integer N such $|z_n| > M$ for al $n > N$. Consider the chordal distance $\chi(z_n,\infty) = 2/\sqrt{(|z_n|^2 + 1)} < 2/\sqrt{(|z_n|^2)} = 2/|z_n| < 2/M < \varepsilon$ for all $n > N$. Thus $z_n \to \infty$ as $n \to \infty$ is equivalent to $\chi(z_n,\infty) \to 0$ as $n \to \infty$.

24. If $\lim_{z\to z_0} f(z) = \infty$, then for any M>0 there exis $\delta > 0$ such that $|f(z)| > M$ for all $|z-z_0| < \delta$. Consider $\chi(f(z),\infty) = 2/\sqrt{(|f(z)|^2 + 1)} < 2/\sqrt{(|f(z)|^2)} = 2/|f(z)| < 2/M < \varepsilon$ for all $|z-z_0| < \delta$. Thus $\lim_{z\to z_0} f(z) = \infty$, is equivalent to $\lim_{z\to\infty} \chi(f(z),\infty) = 0$.

25. (a) $\infty$ (b) 3 (c) $\infty$ (d) $\infty$ (f) the limit does not exist.

## EXERCISES 2.3: Analyticity

1. Let $\Delta z = z - z_0$ so that $\Delta z \to 0 \iff z \to z_0$. Then

$$\lim_{\Delta z\to 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} = L \iff$$

given $\varepsilon > 0$, there is a $\delta > 0$ such that

$$\left|\frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} - L\right| < \varepsilon \text{ whenever} |\Delta z - 0| < \delta \iff$$

$$\left|\frac{f(z) - f(z_0)}{z - z_0} - L\right| < \varepsilon \text{whenever} |z - z_0| < \delta \iff$$

$$\lim_{z\to z_0} \frac{f(z) - f(z_0)}{z - z_0} = L.$$

2. If $\lambda(z) = \dfrac{f(z) - f(z_0)}{z - z_0} - f'(z_0)$, then $\lambda(z) \to 0$ as $z \to z_0$ and $f(z_0) + f'(z_0)(z - z_0) + \lambda(z)(z - z_0) = f(z)$.

3. $\lim_{z\to z_0} f(z) = \lim_{z\to z_0} [f(z_0) + f'(z_0)(z - z_0) + \lambda(z)(z - z_0)]$
$= f(z_0) + 0 + 0 = f(z_0)$.



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

4. a. $\lim\limits_{\Delta z\to 0}\dfrac{\mathrm{Re}\,(z+\Delta z)-\mathrm{Re}\,(z)}{\Delta z}=\lim\limits_{\Delta z\to 0}\dfrac{\mathrm{Re}\,(\Delta z)}{\Delta z}=\begin{cases}1, & \text{if } \Delta z=\Delta x\\ 0, & \text{if } \Delta z=i\Delta y\end{cases}$

b. $\lim\limits_{\Delta z\to 0}\dfrac{\mathrm{Im}\,(z+\Delta z)-\mathrm{Im}(z)}{\Delta z}=\lim\limits_{\Delta z\to 0}\dfrac{\mathrm{Im}\,(\Delta z)}{\Delta z}=\begin{cases}0, & \text{if } \Delta z=\Delta x\\ -i, & \text{if } \Delta z=i\Delta y\end{cases}$

c. Case 1, $z=0$.

$$\lim_{\Delta z\to 0}\frac{|0+\Delta z|-|0|}{\Delta z}=\lim_{\Delta z\to 0}\frac{\sqrt{(\Delta x)^2+(\Delta y)^2}}{\Delta x+i\Delta y}=\begin{cases}\pm 1, & \text{if } \Delta z=\Delta x\\ -i, & \text{if } \Delta z=\pm i\Delta y\end{cases}$$

Case 2, $z\neq 0$.

$$\begin{aligned}&\lim_{\Delta z\to 0}\frac{|z+\Delta z|-|z|}{\Delta z}\\ &=\lim_{\Delta z\to 0}\frac{\sqrt{(x+\Delta x)^2+(y+\Delta y)^2}-\sqrt{x^2+y^2}}{\Delta x+i\Delta y}\\ &=\lim_{\Delta z\to 0}\frac{(x+\Delta x)^2+(y+\Delta y)^2-(x^2+y^2)}{(\Delta x+i\Delta y)(\sqrt{(x+\Delta x)^2+(y+\Delta y)^2}+\sqrt{x^2+y^2})}\\ &=\lim_{\Delta z\to 0}\frac{2x\Delta x+(\Delta x)^2+2y\Delta y+(\Delta y)^2}{(\Delta x+i\Delta y)(\sqrt{(x+\Delta x)^2+(y+\Delta y)^2}+\sqrt{x^2+y^2})}\\ &=\begin{cases}\dfrac{x}{\sqrt{x^2+y^2}}, & \text{if } \Delta z=\Delta x,\ z\neq 0\\ \dfrac{y}{i\sqrt{x^2+y^2}}, & \text{if } \Delta z=i\Delta y,\ z\neq 0\end{cases}\end{aligned}$$

5. Rule 5: $(f\pm g)'(z_0)=\lim\limits_{\Delta z\to 0}\dfrac{(f\pm g)(z_0+\Delta z)-(f\pm g)(z_0)}{\Delta z}$

$$\begin{aligned}&=\lim_{\Delta z\to 0}\left[\frac{f(z_0+\Delta z)-f(z_0)}{\Delta z}\pm\frac{g(z_0+\Delta z)-g(z_0)}{\Delta z}\right]\\ &=f'(z_0)\pm g'(z_0)\end{aligned}$$

Rule 7: $(fg)'(z_0)=\lim\limits_{\Delta z\to 0}\dfrac{fg(z_0+\Delta z)-fg(z_0)}{\Delta z}$



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

$$= \lim_{\Delta z \to 0} \left\{ \frac{f(z_0+\Delta z)g(z_0+\Delta z) - f(z_0+\Delta z)g(z_0)}{\Delta z} \right.$$

$$\left. + \frac{f(z_0+\Delta z)g(z_0) - f(z_0)g(z_0)}{\Delta z} \right\}$$

$$= \lim_{\Delta z \to 0} \left\{ f(z_0+\Delta z)\frac{[g(z_0+\Delta z) - g(z_0)]}{\Delta z} \right.$$

$$\left. + g(z_0)\frac{[f(z_0+\Delta z) - f(z_0)]}{\Delta z} \right\}$$

$$= f(z_0)g'(z_0) + g(z_0)f'(z_0)$$

6. Let $n > 0$ be an integer.
Then $\dfrac{d}{dz}z^{-n} = \dfrac{d}{dz}\left(\dfrac{1}{z^n}\right) = \dfrac{-nz^{n-1}}{z^{2n}}$ (using Rule 8) $= -nz^{-n-1}$.

7. a. $18z^2 + 16z + i$

b. $-12z(z^2 - 3i)^{-7}$

c. $\dfrac{-iz^4 + (2+27i)z^2 + 2\pi z + 18}{(iz^3 + 2z + \pi)^2}$

d. $\dfrac{-(z+2)^2(5z^2 + (16+i)z - 3 + 8i)}{(z^2 + iz + 1)^5}$

e. $24i(z^3-1)^3(z^2+iz)^{99}(53z^4 + 28iz^3 - 50z - 25i)$

8. Let $z = z_0 + \Delta z$. Then

$$\lim_{z \to z_0} \frac{|f(z) - f(z_0)|}{|z - z_0|} = \left| \lim_{\Delta z \to 0} \frac{f(z_0+\Delta z) - f(z_0)}{\Delta z} \right| = |f'(z_0)|.$$

$$\lim_{z \to z_0} \arg[f(z) - f(z_0)] - \arg(z - z_0) = \lim_{z \to z_0} \arg\left[\frac{f(z) - f(z_0)}{z - z_0}\right]$$

$$\arg\left[\lim_{\Delta z \to 0} \frac{f(z_0+\Delta z) - f(z_0)}{\Delta z}\right] = \arg[f'(z_0)]$$



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

9. a. $2-3i$

   b. $\pm i$

   c. $\dfrac{-1\pm i\sqrt{15}}{2}$

   d. $\dfrac{1}{2}, 1$

10. $\displaystyle\lim_{\Delta z\to 0}\frac{|z_0+\Delta z|^2-|z_0|^2}{\Delta z}$

$$= \lim_{\Delta z\to 0}\frac{(z_0+\Delta z)(\overline{z}_0+\overline{\Delta z})-z_0\overline{z}_0}{\Delta z}$$

$$= \lim_{\Delta z\to 0}\left(\overline{z}_0+\frac{\overline{\Delta z}}{\Delta z}z_0+\overline{\Delta z}\right) = \begin{cases} \overline{z_0}+z_0 & \text{if } \Delta z=\Delta x \\ \overline{z_0}-z_0 & \text{if } \Delta z=i\Delta y \end{cases}$$

If $z_0=0$, then the difference quotient is

$$\lim_{\Delta z\to 0}(0+0+\overline{\Delta z})=0.$$

11. a. nowhere analytic

    b. nowhere analytic

    c. analytic except at $z=5$

    d. everywhere analytic

    e. nowhere analytic

    f. analytic except at $z=0$

    g. nowhere analytic

    h. nowhere analytic

12. The case when $n=1$ is trivial. Assume that the result holds for all positive integers less than or equal to $n$ and define $Q(z)=P(z)(z-z_{n+1})$. Since $Q'(z)=P'(z)(z-z_{n+1})+P(z)$, it follows that

$$\frac{Q'(z)}{Q(z)}=\frac{P'(z)}{P(z)}+\frac{1}{z-z_{n+1}}=\frac{1}{z-z_1}+\frac{1}{z-z_2}+\cdots+\frac{1}{z-z_{n+1}}$$



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

13. a, b, d, f, and g are always true

14. $\lim\limits_{z\to z_0} \dfrac{f(z)}{g(z)} = \lim\limits_{z\to z_0} \dfrac{[f(z)-f(z_0)]/(z-z_0)}{[g(z)-g(z_0)]/(z-z_0)} = \dfrac{f'(z_0)}{g'(z_0)}$

15. $\dfrac{3}{5}$

16. Any point on the line through $z_1$ and $z_2$ has the form $z = -\dfrac{1}{2} + i\sqrt{3}\left(\dfrac{1}{2} - t\right)$, $t$ real (see Section 1.3, Exercise 18). However, $f(z_2) - f(z_1) = 0$ but $f'(w) = 3w^2 \neq 0$ on the line in question.

17. $F'(z_0) = f(z_0)(gh)'(z_0) + f'(z_0)gh(z_0)$
$= f(z_0)[g(z_0)h'(z_0) + g'(z_0)h(z_0)] + f'(z_0)g(z_0)h(z_0)$
$= f'(z_0)g(z_0)h(z_0) + f(z_0)g'(z_0)h(z_0) + f(z_0)g(z_0)h'(z_0)$

## EXERCISES 2.4: The Cauchy-Riemann Equations

1. a. $\dfrac{\partial u}{\partial x} = 1 \neq \dfrac{\partial v}{\partial y} = -1$

   b. $\dfrac{\partial u}{\partial x} = 1 \neq \dfrac{\partial v}{\partial y} = 0$

   c. $\dfrac{\partial u}{\partial y} = 2 \neq -\dfrac{\partial v}{\partial x} = 1$

2. $\dfrac{\partial u}{\partial x} = 3x^2 + 3y^2 - 3 = \dfrac{\partial v}{\partial y}$, but $\dfrac{\partial u}{\partial y} = 6xy = \dfrac{\partial v}{\partial x}$. Therefore $\dfrac{\partial u}{\partial y} = -\dfrac{\partial v}{\partial x}$ only when $x = 0$ or $y = 0$. This means $h$ is differentiable on the axes but $h$ is nowhere analytic since lines are not open sets in the complex plane.

3. $\dfrac{\partial u}{\partial x} = 6x + 2 = \dfrac{\partial v}{\partial y}$, $\dfrac{\partial u}{\partial y} = -6y = -\dfrac{\partial v}{\partial x}$. Since these partial derivatives exist and are continuous for all $x$ and $y$, $g$ is analytic. $g$ can be written as $g(z) = 3z^2 + 2z - 1$.



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

4. $\dfrac{\partial u}{\partial x} = \lim\limits_{\Delta x \to 0} \dfrac{u(\Delta x, 0) - u(0,0)}{\Delta x} = \lim\limits_{\Delta x \to 0} \dfrac{0}{\Delta x} = 0$
$\dfrac{\partial u}{\partial y} = \lim\limits_{\Delta y \to 0} \dfrac{u(0, \Delta y) - u(0,0)}{\Delta y} = \lim\limits_{\Delta y \to 0} \dfrac{0}{\Delta y} = 0.$
Similarly $\dfrac{\partial v}{\partial x} = 0$ and $\dfrac{\partial v}{\partial y} = 0.$
However, when $\Delta z \to 0$ through real values ($\Delta z = \Delta x$)

$$\lim_{\Delta x \to 0} \frac{f(0 + \Delta x) - f(0)}{\Delta x} = 0,$$

while along the real line $y = x$ ($\Delta z = \Delta x + i\Delta x$)

$$\begin{aligned} \lim_{\Delta z \to 0} \frac{f(0 + \Delta z) - f(0)}{\Delta z} &= \lim_{\Delta x \to 0} \frac{\frac{(\Delta x)^{4/3}(\Delta x)^{5/3} + i(\Delta x)^{5/3}(\Delta x)^{4/3}}{2(\Delta x)^2}}{\Delta x(1+i)} \\ &= \frac{1}{2}. \end{aligned}$$

Therefore $f$ is not differentiable at $z = 0$.

5. $\dfrac{\partial u}{\partial x} = 2e^{x^2 - y^2}[x\cos(2xy) - y\sin(2xy)] = \dfrac{\partial v}{\partial y}$
$\dfrac{\partial u}{\partial y} = -2e^{x^2 - y^2}[y\cos(2xy) + x\sin(2xy)] = -\dfrac{\partial v}{\partial x}$
$f$ is entire because these first partials exist and are continuous for all $x$ and $y$.

$$\begin{aligned} f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} &= 2e^{x^2 - y^2}(x + iy)[\cos(2xy) + i\sin(2xy)] \\ &= 2e^{(x^2 - y^2)}e^{i2xy}(x + iy) \\ &= 2ze^{z^2} \end{aligned}$$

(This derivative could have been obtained directly, since $f(z) = e^{z^2}$.)

6. $z = re^{i\theta} \Longrightarrow x = r\cos\theta$ and $y = r\sin\theta$ and

$$f(z) = u(x(r,\theta), y(r,\theta)) + iv(x(r,\theta), y(r,\theta))$$

$$\frac{\partial u}{\partial r} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial r} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial r} = \frac{\partial u}{\partial x}\cos\theta + \frac{\partial u}{\partial y}\sin\theta$$


Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

Similar applications of the chain rule yield

$$\frac{\partial u}{\partial \theta} = \frac{\partial u}{\partial x}(-r\sin\theta) + \frac{\partial u}{\partial y} r\cos\theta$$
$$\frac{\partial v}{\partial r} = \frac{\partial v}{\partial x}\cos\theta + \frac{\partial v}{\partial y}\sin\theta$$
$$\frac{\partial v}{\partial \theta} = \frac{\partial v}{\partial x}(-r\sin\theta) + \frac{\partial v}{\partial y} r\cos\theta$$

Replace the partial derivatives on the right sides of the equations for $\frac{\partial u}{\partial r}$ and $\frac{\partial v}{\partial r}$ by their Cauchy-Riemann counterparts to obtain:

$$\frac{\partial u}{\partial r} = \frac{\partial v}{\partial y}\cos\theta - \frac{\partial v}{\partial x}\sin\theta = \frac{1}{r}\frac{\partial v}{\partial \theta}$$
$$\frac{\partial v}{\partial r} = -\frac{\partial u}{\partial y}\cos\theta + \frac{\partial u}{\partial x}\sin\theta = -\frac{1}{r}\frac{\partial u}{\partial \theta}$$

7. Let $h(z) = f(z) - g(z)$. Then $h$ is analytic in $D$ and $h'(z) = 0$ so $h$ is a constant function.

$$h(z) = c = f(z) - g(z) \Longrightarrow f(z) = g(z) + c$$

8. $u(x,y) = c$ in $D \Longrightarrow \frac{\partial u}{\partial x} = 0$ and $\frac{\partial u}{\partial y} = 0 = -\frac{\partial v}{\partial x}$. Hence $f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} = 0$ so $f$ is constant in $D$.

9. By contradiction. If $f$ is analytic in a domain D then $v(x,y) = 0$ (a constant) $\Rightarrow f$ is constant (by condition 8) $\Rightarrow u$ is constant. (However, there is no open set in which $u(x,y) = |z^2 - z|$ is constant).

10. $\operatorname{Im} f(z) = 0$ in $D \Longrightarrow \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} = 0 \Longrightarrow \frac{\partial u}{\partial x} = 0$
$\Longrightarrow f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} = 0 \Longrightarrow f$ is constant in D.

11. $\operatorname{Re} f(z) = \frac{1}{2}[f(z) + \overline{f(z)}]$ is real valued and analytic if both $f$ and $\overline{f}$ are analytic. Hence $\operatorname{Re} f(z)$ is constant by Exercise 10. It follows that $f(z)$ is constant by Exercise 8.



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

12. $|f(z)|$ constant in $D \Longrightarrow |f(z)|^2 = u^2 + v^2$ is constant in $D$. If $u = 0$ or $v = 0$ in $D$, then $f$ is constant by Exercises 8 and 10. Otherwise,

$$\begin{aligned}
\frac{\partial |f|^2}{\partial x} &= 2u\frac{\partial u}{\partial x} + 2v\frac{\partial v}{\partial x} = 0 \\
\frac{\partial |f|^2}{\partial y} &= 2u\frac{\partial u}{\partial y} + 2v\frac{\partial v}{\partial y} = -2u\frac{\partial v}{\partial x} + 2v\frac{\partial u}{\partial x} = 0
\end{aligned}$$

$$\Longrightarrow \frac{1}{2}v\frac{\partial |f|^2}{\partial x} - \frac{1}{2}u\frac{\partial |f|^2}{\partial y} = 0 = (u^2 + v^2)\frac{\partial v}{\partial x}$$

$$\Longrightarrow \frac{\partial v}{\partial x} = 0 \Longrightarrow \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} = \frac{\partial u}{\partial y} = 0$$

$$\Longrightarrow f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} = 0$$

$\Longrightarrow f$ is constant in $D$.

13. $|f(z)|$ is analytic and real-valued, so the result follows from Exercises 10 and 12.

14. If the line is vertical then $\operatorname{Re} f(z)$ is constant and this reduces to Problem 8. If the line is not vertical, then $v(x, y) = mu(x, y) + b$, and

$$\begin{aligned}
\frac{\partial v}{\partial x} &= m\frac{\partial u}{\partial x} = m\frac{\partial v}{\partial y}, \\
\frac{\partial v}{\partial y} &= m\frac{\partial u}{\partial y} = -m\frac{\partial v}{\partial x} = -m^2\frac{\partial v}{\partial y}.
\end{aligned}$$

It follows that

$$\frac{\partial v}{\partial y} = 0 = \frac{\partial u}{\partial x} = \frac{\partial v}{\partial x} = \frac{\partial u}{\partial y} \text{ and } f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} = 0.$$

Hence $f(z)$ is constant.

15. $$\begin{aligned}
J(x_0, y_0) &= \left.\frac{\partial u}{\partial x}\frac{\partial v}{\partial y} - \frac{\partial u}{\partial y}\frac{\partial v}{\partial x}\right|_{(x_0, y_0)} \\
&= \left[\frac{\partial u}{\partial x}(x_0, y_0)\right]^2 + \left[\frac{\partial v}{\partial x}(x_0, y_0)\right]^2 \\
&= |f'(z_0)|^2 \quad \text{(using Equation (1))}
\end{aligned}$$

Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

16. a. $$\frac{\partial \tilde{f}}{\partial \xi} = \frac{\partial f}{\partial x}\frac{\partial x}{\partial \xi} + \frac{\partial f}{\partial y}\frac{\partial y}{\partial \xi}$$
$$= \left(\frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x}\right)\frac{1}{2} + \left(\frac{\partial u}{\partial y} + i\frac{\partial v}{\partial y}\right)\frac{1}{2i}$$
$$= \frac{1}{2}\left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y}\right) + \frac{i}{2}\left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y}\right)$$
$$\frac{\partial \tilde{f}}{\partial \eta} = \frac{\partial f}{\partial x}\frac{\partial x}{\partial \eta} + \frac{\partial f}{\partial y}\frac{\partial y}{\partial \eta}$$
$$= \left(\frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x}\right)\frac{1}{2} + \left(\frac{\partial u}{\partial y} + i\frac{\partial v}{\partial y}\right)\left(\frac{-1}{2i}\right)$$
$$= \frac{1}{2}\left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y}\right) + \frac{i}{2}\left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x}\right)$$

b. $$\frac{\partial \tilde{f}}{\partial \eta} = 0 \Leftrightarrow 0 = \frac{1}{2}\left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y}\right) \text{ and } 0 = \frac{1}{2}\left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x}\right)$$
$$\Leftrightarrow \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

## EXERCISES 2.5: Harmonic Functions

1. a. $u(x,y) = x^2 - y^2 + 2x + 1$, $\dfrac{\partial^2 u}{\partial x^2} = 2 = -\dfrac{\partial^2 u}{\partial y^2} \Longrightarrow \Delta u = 0$
$v(x,y) = 2xy + 2y$, $\dfrac{\partial^2 v}{\partial x^2} = 0 = -\dfrac{\partial^2 v}{\partial y^2} \Longrightarrow \Delta v = 0$

b. $u(x,y) = \dfrac{x}{x^2+y^2}$, $\dfrac{\partial^2 u}{\partial x^2} = \dfrac{2x(x^2-3y^2)}{(x^2+y^2)^3} = -\dfrac{\partial^2 u}{\partial y^2} \Longrightarrow \Delta u = 0$
$v(x,y) = -\dfrac{y}{x^2+y^2}$, $\dfrac{\partial^2 v}{\partial x^2} = \dfrac{-2y(3x^2-y^2)}{(x^2+y^2)^3} = -\dfrac{\partial^2 v}{\partial y^2} \Longrightarrow \Delta v = 0$

c. $u(x,y) = e^x \cos y$, $\dfrac{\partial^2 u}{\partial x^2} = e^x \cos y = -\dfrac{\partial^2 u}{\partial y^2} \Longrightarrow \Delta u = 0$
$v(x,y) = e^x \sin y$, $\dfrac{\partial^2 v}{\partial x^2} = e^x \sin y = -\dfrac{\partial^2 v}{\partial y^2} \Longrightarrow \Delta v = 0$

2. $h(x,y) = ax^2 + bxy - ay^2$



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

3. a. $u = \text{Re}(-iz)$, $v = -x + a$, where $a$ is a constant

b. $u = \text{Re}(-ie^z)$, $v = -e^x \cos y + a$

c. $u = \text{Re}\left(\frac{-i}{2}z^2 - iz - z\right)$, $v = -\frac{1}{2}(x^2 - y^2) - (x + y) + a$

d. It is straightforward to verify that $\Delta u = 0$.
$\frac{\partial u}{\partial x} = \cos x \cosh y = \frac{\partial v}{\partial y}$
$\Rightarrow v(x, y) = \int \cos x \cosh y dy = \cos x \sinh y + \psi(x)$
$\frac{\partial u}{\partial y} = \sin x \sinh y = -\frac{\partial v}{\partial x} = \sin x \sinh y + \psi'(x) \Rightarrow \psi(x) = a$
Thus, $v(x, y) = \cos x \sinh y + a$.

e. It is straightforward to verify that $\Delta u = 0$.
$\frac{\partial u}{\partial x} = \frac{x}{x^2 + y^2} = \frac{\partial v}{\partial y} \Rightarrow$
$v(x, y) = \int \frac{x}{x^2 + y^2} dy = \tan^{-1}\left(\frac{y}{x}\right) + \psi(x)$
$\frac{\partial u}{\partial y} = \frac{y}{x^2 + y^2} = -\frac{\partial v}{\partial x} = \frac{y}{x^2 + y^2} - \psi'(x) \Rightarrow \psi(x) = a$
Thus, $v(x, y) = \tan^{-1}\left(\frac{y}{x}\right) + a$.

f. $u = \text{Re}\left(-ie^{z^2}\right), v = -e^{x^2 - y^2} \cos(2xy) + a$.

4. Suppose $v$ and $w$ are both harmonic conjugates of $u$, and consider $\phi(x, y) = w(x, y) - v(x, y)$. Then (using the Cauchy-Riemann equations for $v$ and $w$),

$$\frac{\partial \phi}{\partial x} = \frac{\partial w}{\partial x} - \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} - \left(-\frac{\partial u}{\partial y}\right) = 0$$

and similarly $\frac{\partial \phi}{\partial y} = 0$. Hence $\phi(x, y) = a$, from which it follows that

$$w(x, y) = v(x, y) + a.$$

5. If $f(z) = u(x, y) + iv(x, y)$ is analytic then $-if(z) = v(x, y) - iu(x, y)$ is analytic. Thus $-u$ is a harmonic conjugate of $v$.


Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

6. Since $f(z) = u + iv$ is analytic, $\frac{1}{2}[f(z)]^2 = \frac{1}{2}(u^2 - v^2) + iuv$ is analytic. Thus $uv = \text{Im}\frac{1}{2}[f(z)]^2$ is harmonic.

7. $\phi(x, y) = x + 1$

8. a. Yes, because $\Delta(u + v) = \Delta u + \Delta v = 0$.

   b. No. Take $u = x, v = x^2 - y^2$ as an example.

   c. Yes, because $\Delta(u_x) = u_{xxx} + u_{xyy} = u_{xxx} + u_{yyx}$

$$= \frac{\partial}{\partial x}(\Delta u) = \frac{\partial}{\partial x}(0) = 0.$$

9. $\phi(x, y) = xy - 1$ $\left(\text{this is Im}\left(\frac{1}{2}z^2 - i\right)\right)$

10. Let $x = r\cos\theta$ and $y = r\sin\theta$.

$$\begin{aligned}
\frac{\partial\phi}{\partial r} &= \frac{\partial\phi}{\partial x}\frac{\partial x}{\partial r} + \frac{\partial\phi}{\partial y}\frac{\partial y}{\partial r} = \frac{\partial\phi}{\partial x}\cos\theta + \frac{\partial\phi}{\partial y}\sin\theta \\
\frac{\partial^2\phi}{\partial r^2} &= \frac{\partial^2\phi}{\partial x^2}\frac{\partial x}{\partial r}\cos\theta + \frac{\partial^2\phi}{\partial y\partial x}\frac{\partial y}{\partial r}\cos\theta \\
&\quad + \frac{\partial^2\phi}{\partial x\partial y}\frac{\partial x}{\partial r}\sin\theta + \frac{\partial^2\phi}{\partial y^2}\frac{\partial y}{\partial r}\sin\theta \\
&= \frac{\partial^2\phi}{\partial x^2}\cos^2\theta + \frac{\partial^2\phi}{\partial y\partial x}2\sin\theta\cos\theta + \frac{\partial^2\phi}{\partial y^2}\sin^2\theta \\
\frac{\partial\phi}{\partial\theta} &= \frac{\partial\phi}{\partial x}\frac{\partial x}{\partial\theta} + \frac{\partial\phi}{\partial y}\frac{\partial y}{\partial\theta} = \frac{\partial\phi}{\partial x}(-r\sin\theta) + \frac{\partial\phi}{\partial y}r\cos\theta \\
\frac{\partial^2\phi}{\partial\theta^2} &= \frac{\partial^2\phi}{\partial x^2}\frac{\partial x}{\partial\theta}(-r\sin\theta) + \frac{\partial^2\phi}{\partial y\partial x}\frac{\partial y}{\partial\theta}(-r\sin\theta) + \frac{\partial\phi}{\partial x}(-r\cos\theta) \\
&\quad + \frac{\partial^2\phi}{\partial x\partial y}\frac{\partial x}{\partial\theta}(r\cos\theta) + \frac{\partial^2\phi}{\partial y^2}\frac{\partial y}{\partial\theta}(r\cos\theta) + \frac{\partial\phi}{\partial y}(-r\sin\theta) \\
&= \frac{\partial^2\phi}{\partial x^2}r^2\sin^2\theta + \frac{\partial^2\phi}{\partial y\partial x}(-2r^2\sin\theta\cos\theta) + \frac{\partial^2\phi}{\partial y^2}r^2\cos^2\theta \\
&\quad + \frac{\partial\phi}{\partial x}(-r\cos\theta) + \frac{\partial\phi}{\partial y}(-r\sin\theta).
\end{aligned}$$



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

Combining these partial derivatives, one gets

$$\frac{\partial^2\phi}{\partial r^2}+\frac{1}{r}\frac{\partial\phi}{\partial r}+\frac{1}{r^2}\frac{\partial^2\phi}{\partial\theta^2}=\frac{\partial^2\phi}{\partial x^2}+\frac{\partial^2\phi}{\partial y^2}=0.$$

11. $\operatorname{Im} f(z)=y-\dfrac{y}{x^2+y^2}=0\Longrightarrow yx^2+y^3-y=y(x^2+y^2-1)=0.$
The points satisfying $x^2+y^2-1=0$ lie on the circle $|z|=1$. The points (other than $z=0$) satisfying $y=0$ lie on the real axis.

12. $f(z)=z^n=r^n(\cos\theta+i\sin\theta)^n=r^n(\cos n\theta+i\sin n\theta)\Longrightarrow$
$\operatorname{Re} f(z)=r^n\cos n\theta$ and $\operatorname{Im} f(z)=r^n\sin n\theta$ are harmonic since $f$ is analytic.

13. $\phi(x,y)=\operatorname{Im} z^4=r^4\sin 4\theta=-4xy^3+4x^3y$

14. Let $\phi(x,y)=\ln|f(z)|=\dfrac{1}{2}\ln(u^2+v^2)$

$$\frac{\partial\phi}{\partial x}=\frac{\partial\phi}{\partial u}\frac{\partial u}{\partial x}+\frac{\partial\phi}{\partial v}\frac{\partial v}{\partial x}=\frac{u\dfrac{\partial u}{\partial x}+v\dfrac{\partial v}{\partial x}}{u^2+v^2}$$

$$\frac{\partial^2\phi}{\partial x^2}=\frac{(v^2-u^2)\left[\left(\dfrac{\partial u}{\partial x}\right)^2-\left(\dfrac{\partial v}{\partial x}\right)^2\right]-4uv\dfrac{\partial u}{\partial x}\dfrac{\partial v}{\partial x}}{(u^2+v^2)^2}+\frac{u\dfrac{\partial^2 u}{\partial x^2}+v\dfrac{\partial^2 v}{\partial x^2}}{u^2+v^2}$$

A similar calculation yields $\dfrac{\partial^2\phi}{\partial y^2}$. By applying Laplace's equation and the Cauchy-Riemann equations of $u$ and $v$ to $\dfrac{\partial^2\phi}{\partial x^2}+\dfrac{\partial^2\phi}{\partial y^2}$, the sum simplifies to reveal that $\Delta\phi=0$.

15. Consider $\varphi(z)=\operatorname{Re}(Az^n+Bz^{-n})+C$ which is harmonic for $1\le|z|\le2$.
Consider the polar form for z. $z=re^{i\theta}$ and select n=3 to agree with the cosine argument. $\varphi(re^{i\theta})=Ar^3\operatorname{Re}(e^{i3\theta})+Br^{-3}\operatorname{Re}(e^{-i3\theta})+C$.
$\varphi(re^{i\theta})=Ar^3\cos3\theta+Br^{-3}\cos3\theta+C=(Ar^3+Br^{-3})\cos3\theta+C$.

$r=1\Rightarrow(A+B)\cos3\theta+C=0\Rightarrow A+B=0,\ C=0.$
$r=2\Rightarrow(A*8+B/8)\cos3\theta=5\cos3\theta.\ A=40/63,\ B=-40/63$
$\varphi(re^{i\theta})=(40/63)(r^3-r^{-3})\cos3\theta=(40/63)\operatorname{Re}(z^3-z^{-3})$.

16. $\phi(x,y)=\dfrac{1}{\ln 3}\ln|z|-1$ or $\phi(x,y)=\ln\left|\dfrac{z}{3}\right|$ are two possibilities.


Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

17. a. $\phi(x,y) = \text{Re}(z^2+5z+1) = x^2 - y^2 + 5x + 1$

b. $\phi(x,y) = 2\text{Re}\left(\dfrac{z^2}{z+2i}\right) = \dfrac{2x(x^2+4y+y^2)}{x^2+y^2+4y+4}$

18. Let $u = \phi_x$, $v = -\phi_y$. Then

$$\frac{\partial u}{\partial x} = \phi_{xx} = -\phi_{yy} = \frac{\partial v}{\partial y}$$
$$\frac{\partial u}{\partial y} = \phi_{xy} = -\frac{\partial v}{\partial x}$$

19. $\cos^2\theta = (\frac{1}{2})\cos 2\theta + \frac{1}{2} = \varphi(z) = A\text{Re}(r^{-2}e^{-i2\theta})n + B = Ar^{-2}\cos 2\theta + B$. In the limit as $r\to\infty$ $\varphi(z) = \frac{1}{2} \Rightarrow B = \frac{1}{2}$. On the circle $|z|=1$, $r = 1$. $\Rightarrow A = \frac{1}{2}$. $\varphi(z) = (\frac{1}{2})r^{-2}\cos 2\theta + \frac{1}{2} = \text{Re}\,[1/(2z^2)] + \frac{1}{2}$.

20. In order that $\dfrac{\partial v}{\partial y} = \dfrac{\partial u}{\partial x}$, let $v(x,y) = \displaystyle\int_0^y \frac{\partial u}{\partial x}(x,\eta)d\eta + \psi(x)$. Then

$$\begin{aligned}\frac{\partial v}{\partial x} &= \int_0^y \frac{\partial^2 u}{\partial x^2}(x,\eta)d\eta + \psi'(x)\\ &= -\int_0^y \frac{\partial^2 u}{\partial y^2}(x,\eta)d\eta + \psi'(x) \quad \text{(because } u \text{ is harmonic)}\\ &= -\frac{\partial u}{\partial y}(x,y) + \frac{\partial u}{\partial y}(x,0) + \psi'(x).\end{aligned}$$

In order that $\dfrac{\partial v}{\partial x} = -\dfrac{\partial u}{\partial y}$, it must be true that $\psi'(x) = -\dfrac{\partial u}{\partial y}(x,0)$. Thus,

$$\psi(x) = -\int_0^x \frac{\partial u}{\partial y}(\zeta,0)d\zeta + a$$

and

$$v(x,y) = \int_0^y \frac{\partial u}{\partial x}(x,\eta)d\eta - \int_0^x \frac{\partial u}{\partial y}(\zeta,0)d\zeta + a.$$

21. It is easily verified that $u = \ln|z|$ satisfies Laplace's equation on $\mathbf{C}\setminus\{0\}$ and that $u+iv = \ln|z| + i\text{Arg}(z)$ satisfies the Cauchy-Riemann equations on the domain $D = \mathbf{C}\setminus\{\text{nonpositive real axis}\}$, so that

**Arg $(z)$ is a harmonic conjugate of $u$ on $D$. By Problem 4, any harmonic conjugate of $u$ has to be of the form Arg$(z)+a$ in $D$. It is impossible to have a harmonic conjugate of this form that is continuous on $\mathbf{C}\setminus\{0\}$.**


22. $$\begin{aligned}\frac{\partial u}{\partial x} &= \phi_{xx}\phi_y + \phi_x\phi_{yx} + \psi_{xx}\psi_y + \psi_x\psi_{yx}\\ &= -\phi_{yy}\phi_y + \phi_x\phi_{yx} - \psi_{yy}\psi_y + \psi_x\psi_{yx} = \frac{\partial v}{\partial y}\end{aligned}$$




Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

**EXERCISES 2.6: Steady-State Temperature as a Harmonic Function.**


1. a.





b.




c.





d.







Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

e.





2. This does not violate the maximum principle.






Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

3. This does not violate the maximum principle.






Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual

Exercises 2.7

1. $f(z) = z^2 + c$ where c is a real constant.
$\zeta_1 = (1 + \sqrt{(1-4c)})/2$, $\zeta_2 = (1 - \sqrt{(1-4c)})/2$
Only $\zeta_2$ is an attractor for $-3/4 < c < ¼$.

2. $f(\varsigma) = \varsigma$ and $f'(\varsigma) > 1$ Therefore we can pick a real number $\rho$ between 1 and $|f'(\varsigma)|$ such that $|f(z) - \varsigma| = \rho|z-\varsigma|$ for all z in a sufficiently small disk around $\varsigma$. If any point $z_0$ in this disk is the seed for an orbit $z_1 = f(z_0)$, $z_2 = f(z_1)$, … $z_n = f(z_{n-1})$, then we have $|z_n-\varsigma| \geq \rho|z_{n-1}-\varsigma| \geq \ldots \geq \rho^n|z_{n-1}-\varsigma|$. Because $\rho>1$, the point $z_n$ moves away from $\varsigma$ until the magnitude of the derivative becomes 1 or less. The orbit is out of the disk.

3. (a) Fixed points are $\zeta_1 = i$, $\zeta_2 = -i$. Both are repellors.
(b) Fixed points are $\zeta_1 = 1/2$, $\zeta_2 = -1/2$ , $\zeta_3 = -1$. Fixed points $\zeta_1$ and $\zeta_3$ are repellors, but fixed point $\zeta_2$ is an attractor.

4. $z_0 = e^{i2\pi\alpha}$ with $\alpha$ an irrational real number. $z_n = e^{i2\pi\alpha 2^{\wedge}n}$. Because $|z_n| = 1$, the trajectory will follow the unit circle. If iterations p and q coincide, $2\pi\alpha 2^p - 2\pi\alpha 2^q = 2\pi\alpha(2^p - 2^q) = 2\pi k$ for some integer k. But because $(2^p - 2^q)$ is an integer that can be represented by m, the equation $2\pi\alpha m = 2\pi k$ is satisfied only if $k=\alpha m$ or $\alpha = k/m$. Because $\alpha$ is irrational it cannot be represented by a rational number and no iterations repeat.

5. Fixed points are $\zeta_1 = -1/2 + i\sqrt{5}/2$ (an attractor) and $\zeta_2 = -1/2 - i\sqrt{5}/2$ (a repellor).

6. $f(z) = z^2$. The seed is $z_0$. $z_1 = z_0^2$ , $z_2 = z_0^4$, … $z_n = z_0^{\wedge}(2^n)$. To have an n cycle $z_n = z_0 = z_0^{\wedge}(2^n)$. Or $z_n/z_0 = z_0^{\wedge}(2^n-1) = 1 = e^{i2\pi}$. Solving gives $z_0 = e^{\wedge}(i2\pi/(2^n-1))$.

7. The cycle is 4. $2^4(2\pi/p) = 2\pi \bmod p \Rightarrow 2^4 = 1 \bmod p$. p=3,5,15. 3 will give repeated cycles of length 2. 5 and 15 will give the desired cycles of length 4.

8. Student Matlab: n=100;c=.253; zo=0;y(1)=zo;
for k=1:n-1,y(k+1)=y(k)^2+c;end
plot(y)

9. If $|\alpha| \leq 1$ the whole complex plane is the filled Julia set. If $|\alpha| \geq 1$ the origin is the filled Julia set.

10. $f(z) = z - F(z)/F'(z)$. $f(\zeta) = \zeta - F(\zeta)/F'(\zeta) = \zeta \Rightarrow F(\zeta)/F'(\zeta) = 0 \Rightarrow F(\zeta)=0$ with the possible exception of the points where $F'(\zeta) = 0$.
$f'(z) = 1 - F'(z)/F'(z) + F(z)F''(z)/(F'(z))^2 = F(z)F''(z)/(F'(z))^2$
$f'(\zeta) = F(\zeta)F''(\zeta)/(F'(\zeta))^2 = 0$ where $F'(\zeta) \neq 0$ and every zero of F(z) is an attractor as long as $F'(\zeta) \neq 0$.



Fundamentals of Complex Analysis with Applications to Engineering and Science 3rd edition Saff Solutions Manual