Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

SKU: 9780321977069-SOLUTIONS

CHAPTER 2: First Order Differential Equations

EXERCISES 2.2: Separable Equations

2. This equation is separable because we can separate variables by multiplying both sides by dx and dividing by $4y^2 - 3y + 1$.

4. This equation is separable because

$$\frac{dy}{dx} = \frac{ye^{x+y}}{x^2+2} = \left(\frac{e^x}{x^2+2}\right) ye^y = g(x)p(y).$$

6. Writing the equation in the form

$$\frac{ds}{dt} = \frac{s+1}{st} - s^2,$$

we see that the right-hand side cannot be represented in the form $g(t)p(s)$. Therefore, the equation is not separable.

8. To separate variables, we divide the equation by x and multiply by dt. Integrating yields

$$\frac{dx}{x} = 3t^2 dt \qquad \Rightarrow \qquad \ln|x| = t^3 + C_1 \qquad \Rightarrow \qquad |x| = e^{t^3+C_1} = e^{C_1}e^{t^3}$$
$$\Rightarrow \qquad |x| = C_2 e^{t^3} \qquad \Rightarrow \qquad x = \pm C_2 e^{t^3} = Ce^{t^3},$$

where C_1 is an arbitrary constant and, therefore, $C_2 := e^{C_1}$ is an arbitrary *positive* constant, $C = \pm C_2$ is any *nonzero* constant. Separating variables, we lost a solution $x \equiv 0$, which can be included in the above formula by taking $C = 0$. Thus, $x = Ce^{t^3}$, C – arbitrary constant, is a general solution.

10. Separating variables yields

$$y^2\,dy = \frac{x\,dx}{\sqrt{1+x}} \qquad \Rightarrow \qquad \int y^2\,dy = \int \frac{x\,dx}{\sqrt{1+x}}\,.$$

The second integral can be evaluated using a substitution $u = 1 + x$.

$$\int \frac{x\,dx}{\sqrt{1+x}} = \int \frac{(u-1)du}{\sqrt{u}} = \int \left(u^{1/2} - u^{-1/2}\right) du = \frac{2}{3}\,u^{3/2} - 2u^{1/2} + C.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

Thus,

$$\frac{y^3}{3} = \frac{2}{3}(1+x)^{3/2} - 2(1+x)^{1/2} + C \qquad \Rightarrow \qquad y = \sqrt[3]{2(1+x)^{3/2} - 6(1+x)^{1/2} + C}\,.$$

12. Separating variables, we obtain

$$\frac{dy}{\sec^2 y} = \frac{dx}{1+x^2}\,.$$

Using the trigonometric identities $\sec y = 1/\cos y$ and $\cos^2 y = (1+\cos 2y)/2$ and integrating, we get

$$\begin{aligned}
&\frac{dy}{\sec^2 y} = \frac{dx}{1+x^2} \qquad \Rightarrow \qquad \frac{(1+\cos 2y)dy}{2} = \frac{dx}{1+x^2} \\
&\Rightarrow \qquad \int \frac{(1+\cos 2y)dy}{2} = \int \frac{dx}{1+x^2} \\
&\Rightarrow \qquad \frac{1}{2}\left(y + \frac{1}{2}\sin 2y\right) = \arctan x + C_1 \\
&\Rightarrow \qquad 2y + \sin 2y = 4\arctan x + 4C_1 \qquad \Rightarrow \qquad 2y + \sin 2y = 4\arctan x + C.
\end{aligned}$$

The last equation defines implicit solutions to the given differential equation.

14. Writing the given equation in the form $dx/dt = x^3 + x$, we separate the variables to get

$$\frac{dx}{x^3+x} = \frac{dx}{x\,(x^2+1)} = \int \left(\frac{1}{x^2} - \frac{1}{x^2+1}\right) x\,dx = dt\,.$$

Thus,

$$\frac{1}{2}\left[\ln\left(x^2\right) - \ln\left(x^2+1\right)\right] = t + C_1 \qquad \Rightarrow \qquad \frac{x^2}{x^2+1} = e^{2C_1}e^{2t} = Ce^{2t}\,,$$

where $C > 0$ is an arbitrary constant. (Note that, separating variables, we lost a solution $x \equiv 0$, which can be included in the above formula by letting $C = 0$.) Solving for x, we get

$$x = \pm\sqrt{\frac{Ce^{2t}}{1 - Ce^{2t}}}\,, \quad C \geq 0\,.$$

16. To separate variables, we move the term, containing dx, to the right-hand side of the equation and divide both sides by y. This yields

$$y^{-1}dy = -ye^{\cos x}\sin x\,dx \qquad \Rightarrow \qquad y^{-2}dy = -e^{\cos x}\sin x\,dx.$$

Integrating the last equation, we obtain

$$\int y^{-2}dy = \int (-e^{\cos x}\sin x)\,dx \qquad \Rightarrow \qquad -y^{-1} + C = \int e^u du \quad (u = \cos x)$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

$$\Rightarrow \qquad -\frac{1}{y} + C = e^u = e^{\cos x} \qquad \Rightarrow \qquad y = \frac{1}{C - e^{\cos x}},$$

where C is an arbitrary constant.

18. First we find a general solution to the equation. Separating variables and integrating, we get

$$\frac{dy}{dx} = x^3(1-y) \qquad \Rightarrow \qquad \frac{dy}{1-y} = x^3 dx$$

$$\Rightarrow \qquad \int \frac{dy}{1-y} = \int x^3 dx \qquad \Rightarrow \qquad -\ln|1-y| + C_1 = \frac{x^4}{4}$$

$$\Rightarrow \qquad |1-y| = \exp\left(C_1 - \frac{x^4}{4}\right) = Ce^{-x^4/4}.$$

We use the initial condition, $y(0) = 3$, to find C. Thus, substitution $y = 3$ and $x = 0$ into the last equation yields

$$|1-3| = Ce^{-0^4/4} \qquad \Rightarrow \qquad 2 = C.$$

Therefore, $|1-y| = 2e^{-x^4/4}$. Finally, since $1-y(0) = 1-3 < 0$, on an interval containing $x = 0$ one has $1 - y(x) < 0$ so that $|1 - y(x)| = y(x) - 1$. Thus, the answer

$$y - 1 = 2e^{-x^4/4} \qquad \text{or} \qquad y = 2e^{-x^4/4} + 1.$$

20. Separating variables yields

$$(y+1)dy = \frac{4x^2 - x - 2}{x^2(x+1)}\,dx\,. \tag{2.1}$$

For integrating the right-hand side, we use partial fractions.

$$\frac{4x^2 - x - 2}{x^2(x+1)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x+1}$$

Solving for A, B, and C, we get $A = 1$, $B = -2$, and $C = 3$. Thus, integrating (2.1), we obtain

$$\frac{1}{2}(y+1)^2 = \ln x + \frac{2}{x} + 3\ln(x+1) + C.$$

With $y(1) = 1$, $C = -3\ln 2$ so that the answer is

$$\frac{1}{2}(y+1)^2 = \ln x + \frac{2}{x} + 3\ln(x+1) - 3\ln 2$$

or, solving for y,

$$y = -1 + \sqrt{\ln(x^2) + \frac{4}{x} + \ln\left[(x+1)^6\right] - 6\ln 2}\,.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

22. Writing $2ydy = -x^2dx$ and integrating, we find

$$y^2 = -\frac{x^3}{3} + C.$$

With $y(0) = 2$,

$$(2)^2 = -\frac{(0)^3}{3} + C \qquad \Rightarrow \qquad C = 4,$$

and so

$$y^2 = -\frac{x^3}{3} + 4 \qquad \Rightarrow \qquad y = \sqrt{-\frac{x^3}{3} + 4}.$$

We note that, taking the square root, we chose the positive sign because $y(0) > 0$.

24. For a general solution, we separate variables and integrate.

$$\int e^{2y}dy = \int 8x^3dx \qquad \Rightarrow \qquad \frac{e^{2y}}{2} = 2x^4 + C_1 \qquad \Rightarrow \qquad e^{2y} = 4x^4 + C.$$

We substitute now the initial condition, $y(1) = 0$, and obtain

$$1 = 4 + C \qquad \Rightarrow \qquad C = -3.$$

Hence, the answer is given by

$$e^{2y} = 4x^4 - 3 \qquad \Rightarrow \qquad y = \frac{1}{2}\ln\left(4x^4 - 3\right).$$

26. We separate variables and obtain

$$\int \frac{dy}{\sqrt{y}} = -\int \frac{dx}{1+x} \qquad \Rightarrow \qquad 2\sqrt{y} = -\ln|1+x| + C = -\ln(1+x) + C,$$

because at initial point, $x = 0$, $1 + x > 0$. Using the fact that $y(0) = 1$, we find C.

$$2 = 0 + C \qquad \Rightarrow \qquad C = 2,$$

and so $y = [2 - \ln(1+x)]^2/4$ is the answer.

28. We have

$$\frac{dy}{dt} = 2y(1-t) \qquad \Rightarrow \qquad \frac{dy}{y} = 2(1-t)dt \qquad \Rightarrow \qquad \ln|y| = -(t-1)^2 + C$$
$$\Rightarrow \qquad y = \pm e^C e^{-(t-1)^2} = C_1 e^{-(t-1)^2},$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

where $C_1 \neq 0$ is any constant. Separating variables, we lost the solution $y \equiv 0$, which can be included into the above formula by letting $C_1 = 0$. So, a general solution to the given equation is

$$y = C_1 e^{-(t-1)^2}, \quad C_1 \text{ is an arbitrary constant.}$$

Substituting $t = 0$ and $y = 3$, we find

$$3 = C_1 e^{-1} \qquad \Rightarrow \qquad C_1 = 3e \qquad \Rightarrow \qquad y = 3e^{1-(t-1)^2} = 3e^{2t-t^2}.$$

The graph of this function is given in Fig. **2**–**A** at the end of the chapter.

Since $y(t) > 0$ for any t, from the given equation we have $y'(t) > 0$ for $t < 1$ and $y'(t) < 0$ for $t > 1$. Thus $t = 1$ is the point of absolute maximum with $y_{\max} = y(1) = 3e$.

30. **(a)** Dividing by $(y+1)^{2/3}$, multiplying by dx, and integrating, we obtain

$$\int \frac{dy}{(y+1)^{2/3}} = \int (x-3)dx \qquad \Rightarrow \qquad 3(y+1)^{1/3} = \frac{x^2}{2} - 3x + C$$
$$\Rightarrow \qquad y = -1 + \left(\frac{x^2}{6} - x + C_1\right)^3.$$

(b) Substituting $y \equiv -1$ into the original equation yields

$$\frac{d(-1)}{dx} = (x-3)(-1+1)^{2/3} = 0,$$

and so the equation is satisfied.

(c) For $y \equiv -1$ for the solution in part (a), we must have

$$\left(\frac{x^2}{6} - x + C_1\right)^3 \equiv 0 \quad \Leftrightarrow \quad \frac{x^2}{6} - x + C_1 \equiv 0,$$

which is impossible since a quadratic polynomial has at most two zeros.

32. **(a)** The direction field of the given differential equation is shown in Fig. **2**–**B** at the end of the chapter. Using this picture we predict that $\lim_{x\to\infty} \phi(x) = 1$.

(b) In notation of Section 1.4, we have $x_0 = 0$, $y_0 = 1.5$, $f(x, y) = y^2 - 3y + 2$, and $h = 0.1$. With this step size, we need $(1-0)/0.1 = 10$ steps to approximate $\phi(1)$. The results of computation are given in Table **2**–**A** at the end of the chapter. From this table we conclude that $\phi(1) \approx 1.26660$.

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

(c) Separating variables and integrating, we obtain

$$\frac{dy}{y^2-3y+2}=dx \quad \Rightarrow \quad \int \frac{dy}{y^2-3y+2}=\int dx \quad \Rightarrow \quad \ln\left|\frac{y-2}{y-1}\right|=x+C\,,$$

where we have used a partial fractions decomposition

$$\frac{1}{y^2-3y+2}=\frac{1}{y-2}-\frac{1}{y-1}$$

to evaluate the integral. The initial condition, $y(0)=1.5$, implies that $C=0$, and so

$$\ln\left|\frac{y-2}{y-1}\right|=x \qquad \Rightarrow \qquad \left|\frac{y-2}{y-1}\right|=e^x \qquad \Rightarrow \qquad \frac{y-2}{y-1}=-e^x\,.$$

(We have chosen the negative sign because of the initial condition.) Solving for y yields

$$y=\phi(x)=\frac{e^x+2}{e^x+1}$$

The graph of this solution is shown in Fig. **2**–**B** at the end of the chapter.

(d) We find

$$\phi(1)=\frac{e+2}{e+1}\approx 1.26894\,.$$

Thus, the approximate value $\phi(1)\approx 1.26660$ found in part (b) differs from the actual value by less than 0.003.

(e) We find the limit of $\phi(x)$ at infinity writing

$$\lim_{x\to\infty}\frac{e^x+2}{e^x+1}=\lim_{x\to\infty}\left(1+\frac{1}{e^x+1}\right)=1\,,$$

which confirms our guess in part (a).

34. (a) Separating variables and integrating, we get

$$\frac{dT}{T-M}=-k dt \quad \Rightarrow \quad \int\frac{dT}{T-M}=-\int k dt \quad \Rightarrow \quad \ln|T-M|=-kt+C_1$$
$$\Rightarrow \qquad |T-M|=e^{C_1}e^{-kt} \qquad \Rightarrow \qquad T-M=\pm e^{C_1}e^{-kt}=Ce^{-kt},$$

where C is any nonzero constant. We can include the lost solution $T\equiv M$ into this formula by letting $C=0$. Thus, a general solution to the equation is

$$T=M+Ce^{-kt}.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

(b) Given that $M = 70°$, $T(0) = 100°$, $T(6) = 80°$, we form a system to determine C and k.

$$\begin{cases} 100 &= 70 + C \\ 80 &= 70 + Ce^{-6k} \end{cases} \quad \Rightarrow \quad \begin{cases} C &= 30 \\ k &= -(1/6)\ln[(80-70)/30] = (1/6)\ln 3. \end{cases}$$

Therefore,

$$T = 70 + 30e^{-(t\ln 3)/6} = 70 + (30)3^{-t/6},$$

and after 20 min the reading is

$$T(20) = 70 + (30)3^{-20/6} \approx 70.77°.$$

36. A general solution to the cooling equation found in Problem 34, that is, $T = M + Ce^{-kt}$. Since $T(0) = 100°$, $T(5) = 80°$, and $T(10) = 65°$, we determine M, C, and k from the system

$$\begin{cases} M + C &= 100 \\ M + Ce^{-5k} &= 80 \\ M + Ce^{-10k} &= 65 \end{cases} \quad \Rightarrow \quad \begin{cases} C(1 - e^{-5k}) &= 20 \\ Ce^{-5k}(1 - e^{-5k}) &= 15 \end{cases} \quad \Rightarrow \quad e^{-5k} = 3/4.$$

To find M, we can now use the first two equations in the above system.

$$\begin{cases} M + C &= 100 \\ M + (3/4)C &= 80 \end{cases} \quad \Rightarrow \quad M = 20.$$

38. With $m = 100$, $g = 9.8$, and $k = 5$, the equation becomes

$$100\frac{dv}{dt} = 100(9.8) - 5v \quad \Rightarrow \quad 20\frac{dv}{dt} = 196 - v.$$

Separating variables and integrating yields

$$\int \frac{dv}{v - 196} = -\frac{1}{20}\int dt \quad \Rightarrow \quad \ln|v - 196| = -\frac{t}{20} + C_1 \quad \Rightarrow \quad v = 196 + Ce^{-t/20},$$

where C is an arbitrary nonzero constant. With $C = 0$, this formula also gives the (lost) constant solution $v(t) \equiv 196$. From the initial condition, $v(0) = 10$, we find C.

$$196 + C = 10 \quad \Rightarrow \quad C = -186 \quad \Rightarrow \quad v(t) = 196 - 186e^{-t/20}.$$

The terminal velocity of the object can be found by letting $t \to \infty$.

$$v_\infty = \lim_{t\to\infty}\left(196 - 186e^{-t/20}\right) = 196\ (\text{m/sec}).$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

40. **(a)** Substituting $\rho = Mp/RT$ into $dp/dz = -\rho g$ yields

$$\frac{dp}{dz} = -\left(\frac{Mp}{RT}\right) g = -\frac{Mg}{RT}\, p.$$

Separating variables and integrating, we find

$$\frac{dp}{p} = -\frac{Mg}{RT}\, dz \tag{2.2}$$

$$\Rightarrow \qquad \ln p = -\frac{Mg}{RT}\, z + C_1 \qquad \Rightarrow \qquad p(z) = Ce^{-(Mg/RT)z}. \tag{2.3}$$

For $z = z_0$,

$$p\,(z_0) = Ce^{-(Mg/RT)z_0} \qquad \Rightarrow \qquad C = p\,(z_0)\, e^{(Mg/RT)z_0}\,.$$

Thus,

$$p(z) = p\,(z_0)\, e^{(Mg/RT)(z_0 - z)} = p\,(z_0)\, e^{-Mg(z-z_0)/(RT)}\,.$$

(b) If $T = T(z)$ varies, then integrating (2.2) from z_0 to z we obtain

$$\ln \frac{p(z)}{p\,(z_0)} = -\frac{Mg}{R} \int_{z_0}^{z} \frac{d\zeta}{T(\zeta)}$$

$$\Rightarrow \qquad p(z) = p\,(z_0) \exp\left[-\frac{Mg}{R} \int_{z_0}^{z} \frac{d\zeta}{T(\zeta)}\right].$$

(c) With the given formula for $T(z)$, the integral in part (b) gives

$$\int_{z_0}^{z} \frac{d\zeta}{288 - 0.0065\,(\zeta - z_0)} = -\frac{1}{0.0065} \ln \frac{288 - 0.0065\,(z - z_0)}{288}\,.$$

Therefore,

$$p(z) = p\,(z_0) \left[\frac{288 - 0.0065\,(z - z_0)}{288}\right]^{Mg/(0.0065R)}\,.$$

Substituting the given data into this equation and computing, we get the height

$$z - z_0 \approx 168 \text{ (m)}\,.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

EXERCISES 2.3: Linear Equations

2. Neither.

4. Linear.

6. Linear.

8. Writing the equation in standard form,

$$\frac{dy}{dx} - \frac{y}{x} = 2x + 1,$$

we see that

$$P(x) = -\frac{1}{x} \qquad \Rightarrow \qquad \mu(x) = \exp\left[\int \left(-\frac{1}{x}\right) dx\right] = \exp\left(-\ln x\right) = \frac{1}{x}\,.$$

Multiplying the given equation by $\mu(x)$, we get

$$\frac{d}{dx}\left(\frac{y}{x}\right) = 2 + \frac{1}{x} \qquad \Rightarrow \qquad y = x\int \left(2 + \frac{1}{x}\right) dx = x\left(2x + \ln|x| + C\right).$$

10. From the standard form of the given equation,

$$\frac{dy}{dx} + \frac{2}{x}\,y = x^{-4},$$

we find that

$$\mu(x) = \exp\left[\int (2/x)dx\right] = \exp\left(2\ln|x|\right) = x^2$$

$$\Rightarrow \quad \frac{d}{dx}\left(x^2 y\right) = x^{-2} \qquad \Rightarrow \qquad y = x^{-2}\int x^{-2}dx = x^{-2}\left(-x^{-1} + C\right) = \frac{Cx - 1}{x^3}\,.$$

12. Here, $P(x) = 4$, $Q(x) = x^2 e^{-4x}$. So, $\mu(x) = e^{4x}$ and

$$\frac{d}{dx}\left(e^{4x} y\right) = x^2 \qquad \Rightarrow \qquad y = e^{-4x}\int x^2 dx = e^{-4x}\left(\frac{x^3}{3} + C\right).$$

14. In standard form, we have

$$\frac{dy}{dx} + \frac{3}{x}\,y = \frac{\sin x}{x^2} - 3x\,.$$

Therefore, $P(x) = 3/x$,

$$\mu(x) = \exp\left(\int \frac{3\,dx}{x}\right) = x^3$$

and

$$y = \frac{1}{x^3}\int \left(x\sin x - 3x^4\right) dx = \frac{1}{x^3}\left(\sin x - x\cos x - \frac{3}{5}\,x^5 + C\right).$$

(Integration by parts was used to integrate the first term.)

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

16. Here, $|x| < 1$ and

$$P(x) = -\frac{x^2}{1-x^2} = 1 - \frac{1}{1-x^2}, \quad Q(x) = \frac{1+x}{\sqrt{1-x^2}}.$$

Therefore,

$$\mu(x) = \exp\left[\int \left(1 - \frac{1}{1-x^2}\right) dx\right] = e^x \sqrt{\frac{1-x}{1+x}}$$

so that $\mu(x)Q(x) = e^x$ and

$$y = e^{-x}\sqrt{\frac{1+x}{1-x}}\,(e^x + C) = \sqrt{\frac{1+x}{1-x}}\,(1 + Ce^{-x}).$$

18. Since $\mu(x) = \exp\left(\int 4dx\right) = e^{4x}$, we have

$$\frac{d}{dx}\left(e^{4x}y\right) = e^{4x}e^{-x} = e^{3x}$$
$$\Rightarrow \qquad y = e^{-4x}\int e^{3x}dx = \frac{e^{-x}}{3} + Ce^{-4x}.$$

Substituting the initial condition, $y = 4/3$ at $x = 0$, yields

$$\frac{4}{3} = \frac{1}{3} + C \qquad \Rightarrow \qquad C = 1,$$

and so $y = e^{-x}/3 + e^{-4x}$ is the solution to the given initial value problem.

20. We have

$$\mu(x) = \exp\left(\int \frac{3\,dx}{x}\right) = \exp(3\ln x) = x^3$$
$$\Rightarrow \qquad x^3 y = \int x^3(3x-2)\,dx = \frac{3x^5}{5} - \frac{x^4}{2} + C$$
$$\Rightarrow \qquad y = \frac{3x^2}{5} - \frac{x}{2} + Cx^{-3}.$$

With $y(1) = 1$,

$$1 = y(1) = \frac{3}{5} - \frac{1}{2} + C \qquad \Rightarrow \qquad C = \frac{9}{10} \qquad \Rightarrow \qquad y = \frac{3x^2}{5} - \frac{x}{2} + \frac{9}{10x^3}.$$

22. From the standard form of this equation,

$$\frac{dy}{dx} + y\cot x = x,$$

we find

$$\mu(x) = \exp\left(\int \cot x\,dx\right) = \exp(\ln\sin x) = \sin x.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

(Alternatively, one can notice that the left-hand side of the original equation is the derivative of the product $y \sin x$.) So, using integration by parts, we obtain

$$y \sin x = \int x \sin x \, dx = -x \cos x + \sin x + C$$
$$\Rightarrow \qquad y = -x \cot x + 1 + C \csc x.$$

We find C using the initial condition $y(\pi/2) = 2$:

$$2 = -\frac{\pi}{2} \cot \frac{\pi}{2} + 1 + C \csc \left(\frac{\pi}{2}\right) = 1 + C \qquad \Rightarrow \qquad C = 1,$$

and the solution is given by

$$y = -x \cot x + 1 + \csc x.$$

24. **(a)** The equation (12) of the text becomes

$$\frac{dy}{dt} + 20y = 50e^{-10t} \qquad \Rightarrow \qquad \mu(t) = e^{20t}$$
$$\Rightarrow \qquad y = e^{-20t} \int e^{20t} \left(50e^{-10t}\right) dt = e^{-20t} \int 50e^{10t} dt = 5e^{-10t} + Ce^{-20t}.$$

Since $y(0) = 40$, we have

$$40 = 5 + C \qquad \Rightarrow \qquad C = 35 \qquad \Rightarrow \qquad y = 5e^{-10t} + 35e^{-20t}.$$

The term $5e^{-10t}$ will eventually dominate.

(b) This time, the equation (12) has the form

$$\frac{dy}{dt} + 10y = 50e^{-10t} \qquad \Rightarrow \qquad \mu(t) = e^{10t}$$
$$\Rightarrow \qquad y = e^{-10t} \int e^{10t} \left(50e^{-10t}\right) dt = e^{-10t}(50t + C).$$

Substituting the initial condition yields

$$40 = y(0) = C \qquad \Rightarrow \qquad y = e^{-10t}(50t + 40).$$

26. Here

$$P(x) = \frac{\sin x \cos x}{1 + \sin^2 x}$$
$$\Rightarrow \quad \mu(x) = \exp\left(\int \frac{\sin x \cos x \, dx}{1 + \sin^2 x}\right) = \exp\left[\frac{1}{2} \ln\left(1 + \sin^2 x\right)\right] = \sqrt{1 + \sin^2 x}.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

Thus,

$$y\sqrt{1+\sin^2 x} = \int_0^x \sqrt{1+\sin^2 t}\,dt \quad \Rightarrow \quad y = (1+\sin^2 x)^{-1/2}\int_0^x (1+\sin^2 t)^{1/2}dt$$

and

$$y(1) = (1+\sin^2 1)^{-1/2}\int_0^1 (1+\sin^2 t)^{1/2}dt.$$

We now use the Simpson's Rule to find that $y(1) \approx 0.860$.

28. **(a)** Substituting $y = e^{-x}$ into the equation (16) yields

$$\frac{d(e^{-x})}{dx} + e^{-x} = -e^{-x} + e^{-x} = 0.$$

So, $y = e^{-x}$ is a solution to (16).

The function $y = x^{-1}$ is a solution to (17) because

$$\frac{d(x^{-1})}{dx} + (x^{-1})^2 = (-1)x^{-2} + x^{-2} = 0.$$

(b) For any constant C,

$$\frac{d(Ce^{-x})}{dx} + Ce^{-x} = -Ce^{-x} + Ce^{-x} = 0.$$

Thus $y = Ce^{-x}$ is a solution to (16).

Substituting $y = Cx^{-1}$ into (17), we obtain

$$\frac{d(Cx^{-1})}{dx} + (Cx^{-1})^2 = (-C)x^{-2} + C^2x^{-2} = C(C-1)x^{-2},$$

and so we must have $C(C-1) = 0$ in order that $y = Cx^{-1}$ is a solution to (17). Thus, either $C = 0$ or $C = 1$.

(c) For the function $y = C\hat{y}$, one has

$$\frac{d(C\hat{y})}{dx} + P(x)\,(C\hat{y}) = C\frac{d\hat{y}}{dx} + C\,(P(x)\hat{y}) = C\left(\frac{d\hat{y}}{dx} + P(x)\hat{y}\right) = 0$$

if $\hat{y}$ is a solution to $y' + P(x)y = 0$.

30. **(a)** Multiplying both sides of (18) by y^2, we get

$$y^2\frac{dy}{dx} + 2y^3 = x.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

If $v = y^3$, then $v' = 3y^2y'$. Thus, $y^2y' = v'/3$, and we have

$$\frac{1}{3}\frac{dv}{dx} + 2v = x,$$

which is equivalent to (19).

(b) The equation (19) is linear with $P(x) = 6$ and $Q(x) = 3x$. So,

$$\mu(x) = \exp\left(\int 6dx\right) = e^{6x}$$

$$\Rightarrow \qquad v(x) = e^{-6x}\int (3xe^{6x})dx = \frac{e^{-6x}}{2}\left(xe^{6x} - \int e^{6x}dx\right)$$

$$= \frac{e^{-6x}}{2}\left(xe^{6x} - \frac{e^{6x}}{6} + C_1\right) = \frac{x}{2} - \frac{1}{12} + Ce^{-6x},$$

where $C = C_1/2$ is an arbitrary constant. The back substitution yields

$$y = \sqrt[3]{\frac{x}{2} - \frac{1}{12} + Ce^{-6x}}.$$

32. In the given equation, $P(x) = 2$, which implies that $\mu(x) = e^{2x}$. Following guidelines, first we solve the equation on $[0, 3]$. On this interval, $Q(x) \equiv 2$. Therefore,

$$y_1(x) = e^{-2x}\int (2)e^{2x}dx = 1 + C_1e^{-2x}.$$

Since $y_1(0) = 0$, we get

$$1 + C_1e^0 = 0 \qquad \Rightarrow \qquad C_1 = -1 \qquad \Rightarrow \qquad y_1(x) = 1 - e^{-2x}.$$

For $x > 3$, $Q(x) = -2$ and so

$$y_2(x) = e^{-2x}\int (-2)e^{2x}dx = -1 + C_2e^{-2x}.$$

We now choose C_2 so that

$$y_2(3) = y_1(3) = 1 - e^{-6} \qquad \Rightarrow \qquad -1 + C_2e^{-6} = 1 - e^{-6} \qquad \Rightarrow \qquad C_2 = 2e^6 - 1.$$

Therefore, $y_2(x) = -1 + (2e^6 - 1)e^{-2x}$, and the continuous solution to the given initial value problem on $[0, \infty)$ is

$$y(x) = \begin{cases} 1 - e^{-2x}, & 0 \le x \le 3, \\ -1 + (2e^6 - 1)e^{-2x}, & x > 3. \end{cases}$$

The graph of this function is shown in Fig. **2–D** at the end of the chapter.

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

34. **(a)** Since $P(x)$ is continuous on (a,b), its antiderivatives given by $\int P(x)dx$ are continuously differentiable, and therefore continuous, functions on (a,b). Since the function e^x is continuous on $(-\infty,\infty)$, composite functions $\mu(x) = e^{\int P(x)dx}$ are continuous on (a,b). The range of the exponential function is $(0,\infty)$. This implies that $\mu(x)$ is positive with any choice of the integration constant. Using the chain rule, we conclude that

$$\frac{d\mu(x)}{dx} = e^{\int P(x)dx}\frac{d}{dx}\left(\int P(x)dx\right) = \mu(x)P(x)$$

for any x on (a,b).

(b) Differentiating (8), we apply the product rule and obtain

$$\frac{dy}{dx} = -\mu^{-2}\mu'\left(\int \mu Q\,dx + C\right) + \mu^{-1}\mu Q = -\mu^{-1}P\left(\int \mu Q\,dx + C\right) + Q\,,$$

and so

$$\frac{dy}{dx} + Py = \left[-\mu^{-1}P\left(\int \mu Q\,dx + C\right) + Q\right] + P\left[\mu^{-1}\left(\int \mu Q\,dx + C\right)\right] = Q\,.$$

(c) Suggested choice of the antiderivative and the constant C yields

$$y(x_0) = \mu(x)^{-1}\left(\int_{x_0}^{x} \mu Q dx + y_0\mu(x_0)\right)\Bigg|_{x=x_0} = \mu(x_0)^{-1}y_0\mu(x_0) = y_0\,.$$

(d) We assume that $y(x)$ is a solution to the initial value problem (15). Since $\mu(x)$ is a continuous positive function on (a,b), the equation (5) is equivalent to (4). Since, from the part (a), the left-hand side of (5) is the derivative of the product $\mu(x)y(x)$, this function must be an antiderivative of the right-hand side, which is $\mu(x)Q(x)$. Thus, we come up with (8), where the integral means one of the antiderivatives, for example, the one suggested in the part (c) (which has zero value at x_0). Substituting $x = x_0$ into (8), we conclude that

$$y_0 = y(x_0) = \mu(x_0)^{-1}\left(\int \mu Q dx + C\right)\Big|_{x=x_0} = C\mu(x_0)^{-1},$$

and so $C = y_0\mu(x_0)$ is uniquely defined.

36. **(a)** If $\mu(x) = \exp\left(\int P dx\right)$ and $y_h(x) = \mu(x)^{-1}$, then

$$\frac{dy_h}{dx} = (-1)\mu(x)^{-2}\frac{d\mu(x)}{dx} = -\mu(x)^{-2}\mu(x)P(x) = -\mu(x)^{-1}P(x)$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

and so

$$\frac{dy_h}{dx} + P(x)y_h = -\mu(x)^{-1}P(x) + P(x)\mu(x)^{-1} = 0,$$

i.e., y_h is a solution to the equation $y' + Py = 0$. Now, the formula (8) yields

$$y = \mu(x)^{-1}\left(\int \mu(x)Q(x)dx + C\right) = y_h(x)v(x) + Cy_h(x) = y_p(x) + Cy_h(x),$$

where $v(x) = \int \mu(x)Q(x)dx$.

(b) Separating variables in (22) and integrating, we obtain

$$\frac{dy}{y} = -\frac{3\,dx}{x} \qquad \Rightarrow \qquad \int \frac{dy}{y} = -\int \frac{3\,dx}{x} \qquad \Rightarrow \qquad \ln|y| = -3\ln x + C.$$

Since we need just one solution y_h, we take $C = 0$

$$\ln|y| = -3\ln x \qquad \Rightarrow \qquad y = \pm x^{-3},$$

and we choose, say, $y_h = x^{-3}$.

(c) Substituting $y_p = v(x)y_h(x) = v(x)x^{-3}$ into (21), we get

$$\frac{dv}{dx}\,y_h + v\frac{dy_h}{dx} + \frac{3}{x}\,vy_h = \frac{dv}{dx}\,y_h + v\left(\frac{dy_h}{dx} + \frac{3}{x}\,y_h\right) = \frac{dv}{dx}\,y_h = x^2.$$

Therefore, $dv/dx = x^2/y_h = x^5$.

(d) Integrating yields

$$v(x) = \int x^5 dx = \frac{x^6}{6}.$$

(We have chosen zero integration constant.)

(e) The function

$$y = Cy_h + vy_h = Cx^{-3} + \frac{x^3}{6}$$

is a general solution to (21) because

$$\begin{aligned}\frac{dy}{dx} + \frac{3}{x}\,y &= \frac{d}{dx}\left(Cx^{-3} + \frac{x^3}{6}\right) + \frac{3}{x}\left(Cx^{-3} + \frac{x^3}{6}\right)\\ &= \left(-3Cx^{-4} + \frac{x^2}{2}\right) + \left(3Cx^{-4} + \frac{x^2}{2}\right) = x^2.\end{aligned}$$

38. Dividing both sides of (6) by μ and multiplying by dx yields

$$\frac{d\mu}{\mu} = P\,dx \qquad \Rightarrow \qquad \int \frac{d\mu}{\mu} = \int P\,dx$$

$$\Rightarrow \qquad \ln|\mu| = \int P\,dx \qquad \Rightarrow \qquad \mu = \pm\exp\left(\int P\,dx\right).$$

Choosing the positive sign, we obtain (7).

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

40. **(a)** Figure 2-E at the end of this chapter depicts the direction field for $\alpha = \frac{1}{4}, \beta = \frac{1}{2}$. If $u = 1, u' = -\beta < 0; if\ u = 0, u' = \alpha > 0$. Thus $u(t)$ is confined to [0,1] if it ever enters this interval.

(b) The general solution $u(t) = \frac{\alpha}{\alpha+\beta} + Ce^{-(\alpha+\beta)t}$ approaches $\frac{\alpha}{\alpha+\beta}$ as $t \to \infty$.

EXERCISES 2.4: Exact Equations

2. This equation is not separable because the coefficient $x^{10/3} - 2y$ cannot be written as a product $p(x)q(y)$. Writing the equation in the form

$$\frac{dy}{dx} - 2x^{-1}y = -x^{7/3},$$

we see that the equation is linear. Since $M(x, y) = x^{10/3} - 2y$, $N(x, y) = x$,

$$\frac{\partial M}{\partial y} = -2 \neq \frac{\partial N}{\partial x} = 1,$$

and so the equation is not exact.

4. Here $M(x, y) = ye^{xy} + 2x$, $N(x, y) = xe^{xy} - 2y$. Thus

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y}\left(ye^{xy} + 2x\right) = e^{xy} + y\frac{\partial}{\partial y}\left(e^{xy}\right) = e^{xy} + ye^{xy}x = e^{xy}(1 + yx),$$
$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x}\left(xe^{xy} - 2y\right) = e^{xy} + x\frac{\partial}{\partial x}\left(e^{xy}\right) = e^{xy} + xe^{xy}y = e^{xy}(1 + xy),$$

Since $\partial M/\partial y = \partial N/\partial x$, the equation is exact.

Writing the equation in the form

$$\frac{dy}{dx} = -\frac{ye^{xy} + 2x}{xe^{xy} - 2y},$$

we conclude that it is not separable, because the right-hand side cannot be represented as a product of two functions of single variables x and y. Also, the right-hand side is not linear with respect to y, which implies that the equation is not linear with y as the dependent variable. Similarly, choosing x as the dependent variable (taking the reciprocals of both sides), we conclude that the equation is not linear in x either.

6. The differential equation is not separable because $(2xy + \cos y)$ cannot be factored. This equation can be put in standard form by defining x as the dependent variable and y as the independent variable. This gives

$$\frac{dx}{dy} + \frac{2}{y}x = \frac{-\cos y}{y^2},$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

and so we see that it is linear.

If we set $M(x, y) = y^2$ and $N(x, y) = 2xy + \cos y$, we see that the differential equation is also exact, because $M_y(x, y) = 2y = N_x(x, y)$.

8. In this problem, the variables are r and θ, $M(r, \theta) = \theta$, and $N(r, \theta) = 3r - \theta - 1$. Since

$$\frac{\partial M}{\partial \theta} = 1 \neq 3 = \frac{\partial N}{\partial r},$$

the equation is not exact. With r as the dependent variable, the equation takes the form

$$\frac{dr}{d\theta} = -\frac{3r - \theta - 1}{\theta} = -\frac{3}{\theta} r + \frac{\theta + 1}{\theta},$$

and so it is linear. Since the right-hand side in the above equation cannot be factored as $p(\theta)q(r)$, the equation is not separable.

10. In this problem, $M(x, y) = 2x + y$, $N(x, y) = x - 2y$. Thus, $M_y = N_x = 1$, and the equation is exact. We find

$$\begin{aligned}
&F(x, y) = \int (2x + y)dx = x^2 + xy + g(y), \\
&\frac{\partial F}{\partial y} = x + g'(y) = N(x, y) = x - 2y \\
&\Rightarrow \qquad g'(y) = -2y \qquad \Rightarrow \qquad g(y) = \int (-2y)dy = -y^2 \\
&\Rightarrow \qquad F(x, y) = x^2 + xy - y^2,
\end{aligned}$$

and so $x^2 + xy - y^2 = C$ is a general solution.

12. Taking partial derivatives of $M(x, y) = \cos x \cos y + 2x$ and $N(x, y) = -(\sin x \sin y + 2y)$, we obtain

$$\begin{aligned}
\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (\cos x \cos y + 2x) = -\cos x \sin y, \\
\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} [-(\sin x \sin y + 2y)] = -\cos x \sin y,
\end{aligned}$$

and so the equation is exact.

Integrating $M(x, y)$ with respect to x yields

$$F(x, y) = \int M(x, y)dx = \int (\cos x \cos y + 2x)\, dx$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

$$= \cos y \int \cos x\, dx + \int 2x\, dx = \sin x \cos y + x^2 + g(y).$$

To find $g(y)$, we compute the partial derivative of $F(x, y)$ with respect to y and compare the result with $N(x, y)$.

$$\frac{\partial F}{\partial y} = \frac{\partial}{\partial y}\left[\sin x \cos y + x^2 + g(y)\right] = -\sin x \sin y + g'(y) = -(\sin x \sin y + 2y)$$
$$\Rightarrow \qquad g'(y) = -2y \qquad \Rightarrow \qquad g(y) = \int (-2y)dy = -y^2.$$

(We have taken zero integration constant.) Therefore,

$$F(x, y) = \sin x \cos y + x^2 - y^2 = C$$

is a general solution to the given equation.

14. In this equation, the variables are y and t, $M(y, t) = t/y$, $N(y, t) = 1 + \ln y$. Since

$$\frac{\partial M}{\partial t} = \frac{\partial}{\partial t}\left(\frac{t}{y}\right) = \frac{1}{y} \qquad \text{and} \qquad \frac{\partial N}{\partial y} = \frac{\partial}{\partial y}(1 + \ln y) = \frac{1}{y},$$

the equation is exact.

Integrating $M(y, t)$ with respect to y, we get

$$F(y, t) = \int \frac{t}{y}\, dy = t \ln |y| + g(t) = t \ln y + g(t).$$

(From $N(y, t) = 1 + \ln y$ we conclude that $y > 0$.) Therefore,

$$\frac{\partial F}{\partial t} = \frac{\partial}{\partial t}\left[t \ln y + g(t)\right] = \ln y + g'(t) = 1 + \ln y$$
$$\Rightarrow \qquad g'(t) = 1 \qquad \Rightarrow \qquad g(t) = t$$
$$\Rightarrow \qquad F(y, t) = t \ln y + t,$$

and a general solution is given by $t \ln y + t = C$ (or, explicitly, $t = C/(\ln y + 1)$).

16. Computing

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y}\left(ye^{xy} - y^{-1}\right) = e^{xy} + xye^{xy} + y^{-2},$$
$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x}\left(xe^{xy} + xy^{-2}\right) = e^{xy} + xye^{xy} + y^{-2},$$

we see that the equation is exact. Therefore,

$$F(x, y) = \int \left(ye^{xy} - y^{-1}\right) dx = e^{xy} - xy^{-1} + g(y).$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

So,

$$\frac{\partial F}{\partial y} = xe^{xy} + xy^{-2} + g'(y) = N(x,y) \qquad \Rightarrow \qquad g'(y) = 0.$$

Thus, $g(y) = 0$, and the answer is $e^{xy} - xy^{-1} = C$.

18. Since

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} = 2y^2 + \sin(x+y),$$

the equation is exact. We find

$$F(x,y) = \int \left[2x + y^2 - \cos(x+y)\right] dx = x^2 + xy^2 - \sin(x+y) + g(y),$$

$$\frac{\partial F}{\partial y} = 2xy - \cos(x+y) + g'(y) = 2xy - \cos(x+y) - e^y$$

$$\Rightarrow \qquad g'(y) = -e^y \qquad \Rightarrow \qquad g(y) = -e^y.$$

Therefore,

$$F(x,y) = x^2 + xy^2 - \sin(x+y) - e^y = C$$

gives a general solution.

20. We find

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y}\left[y\cos(xy)\right] = \cos(xy) - xy\sin(xy),$$

$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x}\left[x\cos(xy)\right] = \cos(xy) - xy\sin(xy).$$

Therefore, the equation is exact and

$$F(x,y) = \int \left(x\cos(xy) - y^{-1/3}\right) dy = \sin(xy) - \frac{3}{2}\,y^{2/3} + h(x)$$

$$\frac{\partial F}{\partial x} = y\cos(xy) + h'(x) = \frac{2}{\sqrt{1-x^2}} + y\cos(xy)$$

$$\Rightarrow \qquad h'(x) = \frac{2}{\sqrt{1-x^2}} \qquad \Rightarrow \qquad h(x) = 2\arcsin x,$$

and a general solution is given by

$$\sin(xy) - \frac{3}{2}\,y^{2/3} + 2\arcsin x = C.$$

22. In Problem 16, we found that a general solution to this equation is

$$e^{xy} - xy^{-1} = C.$$

Substituting the initial condition, $y(1) = 1$, yields $e - 1 = C$. So, the answer is

$$e^{xy} - xy^{-1} = e - 1.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

24. First, we check the given equation for exactness.

$$\frac{dM}{dx} = e^t = \frac{\partial N}{\partial t}.$$

So, it is exact. We find

$$\begin{aligned} F(t,x) &= \int \left(e^t - 1\right) dx = x\left(e^t - 1\right) + g(t), \\ \frac{\partial F}{\partial t} &= xe^t + g'(t) = xe^t + 1 \qquad \Rightarrow \qquad g(t) = \int dt = t \\ \Rightarrow &\qquad x\left(e^t - 1\right) + t = C \end{aligned}$$

is a general solution. With $x(1) = 1$, we get

$$(1)\,(e-1) + 1 = C \qquad \Rightarrow \qquad C = e,$$

and the solution is given by

$$x = \frac{e - t}{e^t - 1}.$$

26. Taking partial derivatives M_y and N_x, we find that the equation is exact. So,

$$\begin{aligned} F(x,y) &= \int (\tan y - 2)\, dx = x(\tan y - 2) + g(y), \\ \frac{\partial F}{\partial y} &= x\sec^2 y + g'(y) = x\sec^2 y + y^{-1} \\ \Rightarrow &\qquad g'(y) = y^{-1} \qquad \Rightarrow \qquad g(y) = \ln|y|, \end{aligned}$$

and

$$x(\tan y - 2) + \ln|y| = C$$

is a general solution. Substituting $y(0) = 1$ yields $C = 0$. Therefore, the answer is

$$x(\tan y - 2) + \ln y = 0.$$

(We removed the absolute value sign in the logarithmic function because $y(0) > 0$.)

28. **(a)** Computing

$$\frac{\partial M}{\partial y} = \cos(xy) - xy\sin(xy),$$

which must be equal to $\partial N/\partial x$, we find that

$$\begin{aligned} N(x,y) &= \int \left[\cos(xy) - xy\sin(xy)\right] dx \\ &= \int \left[x\cos(xy)\right]'_x\, dx = x\cos(xy) + g(y). \end{aligned}$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

(b) Since

$$\frac{\partial M}{\partial y} = (1+xy)e^{xy} - 4x^3 = \frac{\partial N}{\partial x}\,,$$

we conclude that

$$N(x,y) = \int \left[(1+xy)e^{xy} - 4x^3\right] dx = xe^{xy} - x^4 + g(y).$$

30. (a) Differentiating, we find that

$$\frac{\partial M}{\partial y} = 5x^2 + 12x^3y + 8xy\,,$$
$$\frac{\partial N}{\partial x} = 6x^2 + 12x^3y + 6xy\,.$$

Since $M_y \neq N_x$, the equation is not exact.

(b) Multiplying given equation by $x^n y^m$ and taking partial derivatives of new coefficients yields

$$\begin{aligned}
&\frac{d}{dy}\left(5x^{n+2}y^{m+1} + 6x^{n+3}y^{m+2} + 4x^{n+1}y^{m+2}\right)\\
&\qquad = 5(m+1)x^{n+2}y^m + 6(m+2)x^{n+3}y^{m+1} + 4(m+2)x^{n+1}y^{m+1}\\
&\frac{d}{dx}\left(2x^{n+3}y^m + 3x^{n+4}y^{m+1} + 3x^{n+2}y^{m+1}\right)\\
&\qquad = 2(n+3)x^{n+2}y^m + 3(n+4)x^{n+3}y^{m+1} + 3(n+2)x^{n+1}y^{m+1}.
\end{aligned}$$

In order that these polynomials are equal, we must have equal coefficients at similar monomials. Thus, n and m must satisfy the system

$$\begin{cases} 5(m+1) = 2(n+3) \\ 6(m+2) = 3(n+4) \\ 4(m+2) = 3(n+2). \end{cases}$$

Solving, we obtain $n = 2$ and $m = 1$. Therefore, multiplying the given equation by x^2y yields an exact equation.

(c) We find

$$\begin{aligned}
F(x,y) &= \int \left(5x^4y^2 + 6x^5y^3 + 4x^3y^3\right) dx\\
&= x^5y^2 + x^6y^3 + x^4y^3 + g(y).
\end{aligned}$$

Therefore,

$$\frac{\partial F}{\partial y} = 2x^5y + 3x^6y^2 + 3x^4y^2 + g'(y)$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

$$= \quad 2x^5y + 3x^6y^2 + 3x^4y^2 \qquad \Rightarrow \qquad g(y) = 0,$$

and a general solution to the given equation is

$$x^5y^2 + x^6y^3 + x^4y^3 = C.$$

32. **(a)** The slope of the orthogonal curves, say $m_\perp$, must be $-1/m$, where m is the slope of the original curves. Therefore, we have

$$m_\perp = \frac{F_y(x,y)}{F_x(x,y)} \quad \Rightarrow \quad \frac{dy}{dx} = \frac{F_y(x,y)}{F_x(x,y)} \quad \Rightarrow \quad F_y(x,y)\,dx - F_x(x,y)\,dy = 0.$$

(b) Let $F(x,y) = x^2 + y^2$. Then we have $F_x(x,y) = 2x$ and $F_y(x,y) = 2y$. Substituting these expressions into the result of part (a) gives us

$$2y\,dx - 2x\,dy = 0 \qquad \Rightarrow \qquad y\,dx - x\,dy = 0.$$

To find the orthogonal trajectories, we must solve this differential equation. To this end, note that this equation is separable and thus

$$\int \frac{1}{x}\,dx = \int \frac{1}{y}\,dy \qquad \Rightarrow \qquad \ln|x| = \ln|y| + C$$
$$\Rightarrow \qquad e^{\ln|x| - C} = e^{\ln|y|} \qquad \Rightarrow \qquad y = kx, \quad \text{where } k = \pm e^{-C}.$$

Therefore, the orthogonal trajectories are lines through the origin.

(c) Let $F(x,y) = xy$. Then we have $F_x(x,y) = y$ and $F_y(x,y) = x$. Substituting these expressions into the result of part (a) yields

$$x\,dx - y\,dy = 0.$$

To find the orthogonal trajectories, we must solve this differential equation. To this end, note that this equation is separable and thus

$$\int x\,dx = \int y\,dy \qquad \Rightarrow \qquad \frac{x^2}{2} = \frac{y^2}{2} + C \qquad \Rightarrow \qquad x^2 - y^2 = k\,,$$

where $k := 2C$. Therefore, the orthogonal trajectories are hyperbolas.

34. To use the method described in Problem 32, we rewrite the equation $x^2 + y^2 = kx$ in the form $x + x^{-1}y^2 = k$. Thus, $F(x,y) = x + x^{-1}y^2$,

$$\frac{\partial F}{\partial x} = 1 - x^{-2}y^2, \qquad \frac{\partial F}{\partial y} = 2x^{-1}y.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

Substituting these derivatives in the equation given in Problem 32(b), we get the required. Multiplying the equation by $x^n y^m$, we obtain

$$2x^{n-1}y^{m+1}dx + \left(x^{n-2}y^{m+2} - x^n y^m\right) dy = 0.$$

Therefore,

$$\frac{\partial M}{\partial y} = 2(m+1)x^{n-1}y^m\,,$$
$$\frac{\partial N}{\partial x} = (n-2)x^{n-3}y^{m+2} - nx^{n-1}y^m\,.$$

Thus, to have an exact equation, n and m must satisfy

$$\begin{cases} n-2=0 \\ 2(m+1) = -n\,. \end{cases}$$

Solving, we obtain $n = 2$, $m = -2$. With this choice, the equation becomes

$$2xy^{-1}dx + \left(1 - x^2y^{-2}\right) dy = 0,$$

and so

$$G(x,y) = \int M(x,y)dx = \int 2xy^{-1}dx = x^2y^{-1} + g(y),$$
$$\frac{\partial G}{\partial y} = -x^2y^{-2} + g'(y) = N(x,y) = 1 - x^2y^{-2}.$$

Therefore, $g(y) = y$, and the family of orthogonal trajectories is given by $x^2y^{-1} + y = C$. Writing this equation in the form $x^2 + y^2 - Cy = 0$, we see that, given C, the trajectory is the circle centered at $(0, C/2)$ and of radius $C/2$.

Several given curves and their orthogonal trajectories are shown in Fig. **2**–**E** at the end of the chapter.

36. The first equation in (4) follows from (9) and the Fundamental Theorem of Calculus.

$$\frac{\partial F}{\partial x} = \frac{\partial}{\partial x}\left[\int_{x_0}^{x} M(t,y)dt + g(y)\right] = M(t,y)|_{t=x} = M(x,y).$$

For the second equation in (4),

$$\frac{\partial F}{\partial y} = \frac{\partial}{\partial y}\left[\int_{x_0}^{x} M(t,y)dt + g(y)\right]$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

$$= \frac{\partial}{\partial y}\int_{x_0}^{x} M(t,y)dt + g'(y)$$

$$= \frac{\partial}{\partial y}\int_{x_0}^{x} M(t,y)dt + \left[N(x,y) - \frac{\partial}{\partial y}\int_{x_0}^{x} M(t,y)dt\right] = N(x,y).$$

EXERCISES 2.5: Special Integrating Factors

2. Here, $M(x,y) = 2y^3 + 2y^2$ and $N(x,y) = 3y^2x + 2xy$. Computing

$$\frac{\partial M}{\partial y} = 6y^2 + 4y \qquad \text{and} \qquad \frac{\partial N}{\partial x} = 3y^2 + 2y\,,$$

we conclude that this equation is not exact. Note that these derivatives, as well as M itself, depend on y only. Then, clearly, so does the expression $(\partial N/\partial x - \partial M/\partial y)/M$, and the given equation has an integrating factor depending on y alone. Also, since

$$\frac{\partial M/\partial y - \partial N/\partial x}{N} = \frac{(6y^2+4y)-(3y^2+2y)}{3y^2x+2xy} = \frac{3y^2+2y}{x(3y^2+2y)} = \frac{1}{x}\,,$$

the equation has an integrating factor depending on x only.

Writing the equation in the form

$$\frac{dx}{dy} = -\frac{3y^2x+2xy}{2y^3+y^2} = -\frac{xy(3y+2)}{2y^2(y+1)} = -\frac{y(3y+2)}{2y^2(y+1)}\,x\,,$$

we conclude that it is separable and linear with x as the dependent variable.

4. This equation is not separable, because of the factor (y^2+2xy). It is not linear in either variable because of the terms y^2 and x^2. To see if it is exact, we compute $M_y(x,y)$ and $N_x(x,y)$, and find that

$$M_y(x,y) = 2y + 2x \neq -2x = N_x(x,y).$$

Therefore, the equation is not exact. To see if we can find an integrating factor of the form $\mu(x)$, we compute

$$\frac{\partial M/\partial y - \partial N/\partial x}{N} = \frac{2y+4x}{-x^2}\,,$$

which is not a function of x alone. To see if we can find an integrating factor of the form $\mu(y)$, we compute

$$\frac{\partial N/\partial x - \partial M/\partial y}{M} = \frac{-4x-2y}{y^2+2xy} = \frac{-2(2x+y)}{y(y+2x)} = \frac{-2}{y}\,.$$

Thus the equation has an integrating factor that is a function of y alone.

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

6. In this problem, $M(x,y) = 2y^2x - y$ and $N(x,y) = x$. Therefore,

$$\frac{\partial M}{\partial y} = 4yx - 1 \qquad \text{and} \qquad \frac{\partial N}{\partial x} = 1 \qquad \Rightarrow \qquad \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} = 2 - 4yx\,.$$

The equation is not exact, because $\partial M/\partial y \neq \partial N/\partial x$, but it has an integrating factor, which depends on y because

$$\frac{\partial N/\partial x - \partial M/\partial y}{M} = \frac{2 - 4yx}{2y^2x - y} = \frac{-2(2yx-1)}{y(2yx-1)} = \frac{-2}{y}\,.$$

Isolating dy/dx, we obtain

$$\frac{dy}{dx} = \frac{y - 2y^2x}{x} = \frac{y}{x} - 2y^2\,.$$

The right-hand side cannot be written as $p(x)q(y)$, so the equation is not separable. Also, it is not linear with y as the dependent variable (because of $2y^2$ term). Taking the reciprocals, we conclude that it is not linear with the dependent variable x.

8. The equation $(3x^2 + y)\,dx + (x^2y - x)\,dy = 0$ is not separable or linear. To see if it is exact, we compute

$$\frac{\partial M}{\partial y} = 1 \neq 2xy - 1 = \frac{\partial N}{\partial x}\,.$$

Thus, the equation is not exact. To see if we can find an integrating factor, we compute

$$\frac{\partial M/\partial y - \partial N/\partial x}{N} = \frac{2 - 2xy}{x^2y - x} = \frac{-2(xy-1)}{x(xy-1)} = \frac{-2}{x}\,.$$

Thus, an integrating factor $\mu(x)$ is given by

$$\mu(x) = \exp\left(\int \frac{-2}{x}\,dx\right) = \exp\left(-2\ln|x|\right) = x^{-2}.$$

To solve the equation, we multiply it by the integrating factor x^{-2} and get

$$(3 + yx^{-2})\,dx + (y - x^{-1})\,dy = 0.$$

This new equation is now exact. Thus, we find $F(x,y)$ integrating $M(x,y) = 3 + yx^{-2}$ with respect to x.

$$\begin{aligned} F(x,y) &= \int \left(3 + yx^{-2}\right)dx = 3x - yx^{-1} + g(y) \\ \Rightarrow \qquad & F_y(x,y) = -x^{-1} + g'(y) = N(x,y) = y - x^{-1} \\ \Rightarrow \qquad & g'(y) = y \qquad \Rightarrow \qquad g(y) = \frac{y^2}{2}\,. \end{aligned}$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

Therefore,

$$F(x,y) = 3x - yx^{-1} + \frac{y^2}{2},$$

and an implicit solution to the given equation is

$$\frac{y^2}{2} - \frac{y}{x} + 3x = C.$$

Since $\mu(x) = x^{-2}$ we must check if the solution $x \equiv 0$ was lost. The function $x \equiv 0$ is a solution to the original equation, but is not given by the above implicit solution for any choice of C. Hence,

$$\frac{y^2}{2} - \frac{y}{x} + 3x = C \qquad \text{and} \qquad x \equiv 0$$

give a general solution.

10. We compute the partial derivatives of $M(x,y) = 2y^2 + 2y + 4x^2$ and $N(x,y) = 2xy + x$.

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y}\left(2y^2 + 2y + 4x^2\right) = 4y + 2, \qquad \frac{\partial N}{\partial x} = \frac{\partial}{\partial x}\left(2xy + x\right) = 2y + 1.$$

Although the equation is not exact ($\partial M/\partial y \neq \partial N/\partial x$), the quotient

$$\frac{\partial M/\partial y - \partial N/\partial x}{N} = \frac{(4y+2) - (2y+1)}{2xy + x} = \frac{2y+1}{x(2y+1)} = \frac{1}{x}$$

depends on x only, and so the equation has an integrating factor, which can be found by applying formula (8). Namely,

$$\mu(x) = \exp\left(\int \frac{1}{x}\,dx\right) = \exp\left(\ln|x|\right) = |x|.$$

Note that if μ is an integrating factor, then $-\mu$ is an integrating factor as well. This observation allows us to take $\mu(x) = x$. Multiplying the given differential equation by x yields an exact equation

$$\left(2y^2 + 2y + 4x^2\right) x\,dx + x^2\left(2y + 1\right)\,dy = 0.$$

Therefore,

$$\begin{aligned}
&F(x,y) = \int x^2(2y+1)\,dy = x^2\left(y^2 + y\right) + h(x)\\
\Rightarrow \qquad &\frac{\partial F}{\partial x} = 2x\left(y^2 + y\right) + h'(x) = \left(2y^2 + 2y + 4x^2\right)x\\
\Rightarrow \qquad &h'(x) = 4x^3 \qquad \Rightarrow \qquad h(x) = \int 4x^3 dx = x^4\\
\Rightarrow \qquad &F(x,y) = x^2\left(y^2 + y\right) + x^4 = x^2y^2 + x^2y + x^4,
\end{aligned}$$

and $x^2y^2 + x^2y + x^4 = C$ is a general solution.

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

12. Here, $M(x,y) = 2xy^3 + 1$, $N(x,y) = 3x^2y^2 - y^{-1}$. Since

$$\frac{\partial M}{\partial y} = 6xy^2 = \frac{\partial N}{\partial x},$$

the equation is exact. So, we find that

$$\begin{aligned} F(x,y) &= \int \left(2xy^3 + 1\right) dx = x^2y^3 + x + g(y), \\ \frac{\partial F}{\partial y} &= 3x^2y^2 + g'(y) = 3x^2y^2 - y^{-1} \\ \Rightarrow \qquad g'(y) &= -y^{-1} \qquad \Rightarrow \qquad g(y) = -\ln|y|, \end{aligned}$$

and the given equation has a general solution

$$x^2y^3 + x - \ln|y| = C.$$

14. Multiplying the given equation by $x^n y^m$ yields

$$\left(12x^n y^m + 5x^{n+1}y^{m+1}\right) dx + \left(6x^{n+1}y^{m-1} + 3x^{n+2}y^m\right) dy = 0.$$

Therefore,

$$\begin{aligned} \frac{\partial M}{\partial y} &= 12mx^n y^{m-1} + 5(m+1)x^{n+1}y^m, \\ \frac{\partial N}{\partial x} &= 6(n+1)x^n y^{m-1} + 3(n+2)x^{n+1}y^m. \end{aligned}$$

Matching the coefficients, we get a system

$$\begin{cases} 12m = 6(n+1) \\ 5(m+1) = 3(n+2) \end{cases}$$

to determine n and m. This system has the solution $n = 3$, $m = 2$. Thus, the given equation multiplied by x^3y^2, that is,

$$\left(12x^3y^2 + 5x^4y^3\right) dx + \left(6x^4y + 3x^5y^2\right) dy = 0,$$

is exact. We compute

$$\begin{aligned} F(x,y) &= \int \left(12x^3y^2 + 5x^4y^3\right) dx = 3x^4y^2 + x^5y^3 + g(y), \\ \frac{\partial F}{\partial y} &= 6x^4y + 3x^5y^2 + g'(y) = 6x^4y + 3x^5y^2 \\ \Rightarrow \qquad g'(y) &= 0 \qquad \Rightarrow \qquad g(y) = 0, \end{aligned}$$

and so $3x^4y^2 + x^5y^3 = C$ is a general solution to the given equation.

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

16. **(a)** An equation $Mdx + Ndy = 0$ has an integrating factor $\mu(x+y)$ if and only if the equation

$$\mu(x+y)M(x,y)dx + \mu(x+y)N(x,y)dy = 0$$

is exact. According to Theorem 2, Section 2.4, this means that

$$\frac{\partial}{\partial y}\left[\mu(x+y)M(x,y)\right] = \frac{\partial}{\partial x}\left[\mu(x+y)N(x,y)\right].$$

Applying the product and chain rules yields

$$\mu'(x+y)M(x,y) + \mu(x+y)\frac{\partial M(x,y)}{\partial y} = \mu'(x+y)N(x,y) + \mu(x+y)\frac{\partial N(x,y)}{\partial x}.$$

Collecting similar terms yields

$$\mu'(x+y)\left[M(x,y) - N(x,y)\right] = \mu(x+y)\left[\frac{\partial N(x,y)}{\partial x} - \frac{\partial M(x,y)}{\partial y}\right]$$

$$\Leftrightarrow \qquad \frac{\partial N/\partial x - \partial M/\partial y}{M-N} = \frac{\mu'(x+y)}{\mu(x+y)}. \tag{2.4}$$

The right-hand side of (2.4) depends on $x+y$ only so the left-hand side does.

To find an integrating factor, we let $s = x+y$ and denote

$$G(s) = \frac{\partial N/\partial x - \partial M/\partial y}{M-N}.$$

Then (2.4) implies that

$$\frac{\mu'(s)}{\mu(s)} = G(s) \qquad \Rightarrow \qquad \ln|\mu(s)| = \int G(s)\,ds$$

$$\Rightarrow \quad |\mu(s)| = \exp\left[\int G(s)\,ds\right] \quad \Rightarrow \quad \mu(s) = \pm\exp\left[\int G(s)\,ds\right]. \tag{2.5}$$

In this formula, we can choose either sign and any integration constant.

(b) We compute

$$\frac{\partial N/\partial x - \partial M/\partial y}{M-N} = \frac{(1+y)-(1+x)}{(3+y+xy)-(3+x+xy)} = 1.$$

Applying formula (2.5), we obtain

$$\mu(s) = \exp\left[\int (1)ds\right] = e^s \qquad \Rightarrow \qquad \mu(x+y) = e^{x+y},$$

Multiplying the given equation by $\mu(x+y)$, we get an exact equation

$$e^{x+y}(3+y+xy)dx + e^{x+y}(3+x+xy)dy = 0$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

and follow the procedure of solving exact equations, Section 2.4.

$$\begin{aligned} F(x,y) &= \int e^{x+y}(3+y+xy)\,dx = e^y\left[(3+y)\int e^x dx + y\int xe^x dx\right] \\ &= e^y\left[(3+y)e^x + y(x-1)e^x\right] + h(y) = e^{x+y}(3+xy) + h(y)\,. \end{aligned}$$

Taking the partial derivative of F with respect to y, we find $h(y)$.

$$\begin{aligned} &\frac{\partial F}{\partial y} = e^{x+y}(3+xy+x) + h'(y) = e^{x+y}N(x,y) = e^{x+y}(3+x+xy) \\ &\Rightarrow \qquad h'(y) = 0 \qquad \Rightarrow \qquad h(y) = 0\,. \end{aligned}$$

Thus, a general solution is

$$e^{x+y}(3+xy) = C\,.$$

18. The given condition, $xM(x,y) + yN(x,y) \equiv 0$, is equivalent to $yN(x,y) \equiv -xM(x,y)$. In particular, substituting $x = 0$, we obtain

$$yN(0,y) \equiv -(0)M(0,y) \equiv 0.$$

This implies that $x \equiv 0$ is a solution to the given equation.

To obtain other solutions, we multiply the equation by $x^{-1}y$. This gives

$$\begin{aligned} x^{-1}yM(x,y)dx + x^{-1}yN(x,y)dy &= x^{-1}yM(x,y)dx - x^{-1}xM(x,y)dy \\ &= xM(x,y)\left(x^{-2}ydx - x^{-1}dy\right) = -xM(x,y)d\left(x^{-1}y\right) = 0\,. \end{aligned}$$

Therefore, $x^{-1}y = C$ or $y = Cx$.

Thus, a general solution is

$$y = Cx \quad \text{and} \quad x \equiv 0.$$

20. For the equation

$$e^{\int P(x)dx}\left[P(x)y - Q(x)\right]dx + e^{\int P(x)dx}dy = 0,$$

we compute

$$\begin{aligned} &\frac{\partial M}{\partial y} = \frac{\partial}{\partial y}\left(e^{\int P(x)dx}\left[P(x)y - Q(x)\right]\right) = e^{\int P(x)dx}P(x), \\ &\frac{\partial N}{\partial x} = \frac{\partial}{\partial x}\left(e^{\int P(x)dx}\right) = e^{\int P(x)dx}\frac{d}{dx}\left(\int P(x)dx\right) = e^{\int P(x)dx}P(x). \end{aligned}$$

Therefore, $\partial M/\partial y = \partial N/\partial x$, and the equation is exact.

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

EXERCISES 2.6: Substitutions and Transformations

2. We can write the equation in the form

$$\frac{dy}{dx} = (y - 4x - 1)^2 = [(y - 4x) - 1]^2 = G(y - 4x),$$

where $G(t) = (t - 1)^2$. Thus, it is of the form $dy/dx = G(ax + by)$.

4. In this equation, the variables are x and t. Its coefficients, $t + x + 2$ and $3t - x - 6$, are linear functions of x and t. Therefore, the given equation is an equation with linear coefficients.

6. The given differential equation is not homogeneous due to the e^{-2x} terms. Since it can be written in the form $dy/dx + P(x)y = Q(x)y^n$, namely,

$$\frac{dy}{dx} - y = e^{2x}y^3.$$

it is a Bernoulli equation. The differential equation does not have linear coefficients, and it is not of the form $y' = G(ax + by)$ either.

8. Here, the variables are y and θ. Writing

$$\frac{dy}{d\theta} = -\frac{y^3 - \theta y^2}{2\theta^2 y} = -\frac{(y/\theta)^3 - (y/\theta)^2}{2(y/\theta)},$$

we see that the right-hand side is a function of y/θ alone. Hence, the equation is homogeneous.

10. First, we write the equation in the form

$$\frac{dy}{dx} = \frac{-3x^2 + y^2}{xy - x^3y^{-1}} = \frac{y^3 - 3x^2y}{xy^2 - x^3} = \frac{(y/x)^3 - 3(y/x)}{(y/x)^2 - 1}.$$

Therefore, it is homogeneous, and we we make a substitution $y/x = u$ or $y = xu$. Then $y' = u + xu'$, and the equation becomes

$$u + x\frac{du}{dx} = \frac{u^3 - 3u}{u^2 - 1}.$$

Separating variables and integrating yields

$$x\frac{du}{dx} = \frac{u^3 - 3u}{u^2 - 1} - u = -\frac{2u}{u^2 - 1} \qquad \Rightarrow \qquad \frac{u^2 - 1}{u}\,du = -\frac{2}{x}\,dx$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

$$\Rightarrow \qquad \int \frac{u^2-1}{u}\,du = -\int \frac{2}{x}\,dx \qquad \Rightarrow \qquad \int \left(u - \frac{1}{u}\right) du = -2\int \frac{dx}{x}$$
$$\Rightarrow \qquad \frac{1}{2}\,u^2 - \ln|u| = -2\ln|x| + C_1 \qquad \Rightarrow \qquad u^2 - \ln\left(u^2\right) + \ln(x^4) = C.$$

Substituting back y/x for u and simplifying, we finally get

$$\left(\frac{y}{x}\right)^2 - \ln\left(\frac{y^2}{x^2}\right) + \ln(x^4) = C \qquad \Rightarrow \qquad \frac{y^2}{x^2} - \ln\left(\frac{y^2}{x^6}\right) = C.$$

12. From

$$\frac{dy}{dx} = -\frac{x^2+y^2}{2xy} = -\frac{1}{2}\left(\frac{x}{y} + \frac{y}{x}\right),$$

making the substitution $v = y/x$, we obtain

$$v + x\frac{dv}{dx} = -\frac{1}{2}\left(\frac{1}{v} + v\right) = -\frac{1+v^2}{2v} \qquad \Rightarrow \qquad x\frac{dv}{dx} = -\frac{1+v^2}{2v} - v = -\frac{1+3v^2}{2v}$$
$$\Rightarrow \qquad \frac{2v\,dv}{1+3v^2} = -\frac{dx}{x} \qquad \Rightarrow \qquad \int \frac{2v\,dv}{1+3v^2} = -\int \frac{dx}{x}$$
$$\Rightarrow \qquad \frac{1}{3}\ln\left(1+3v^2\right) = -\ln|x| + C_2 \qquad \Rightarrow \qquad 1+3v^2 = C_1|x|^{-3},$$

where $C_1 = e^{3C_2}$ is any positive constant. Making the back substitution, we finally get

$$1 + 3\left(\frac{y}{x}\right)^2 = \frac{C_1}{|x|^3} \qquad \Rightarrow \qquad 3\left(\frac{y}{x}\right)^2 = \frac{C_1}{|x|^3} - 1 = \frac{C_1 - |x|^3}{|x|^3}$$
$$\Rightarrow \quad 3|x|y^2 = C_1 - |x|^3 \quad \Rightarrow \quad 3|x|y^2 + |x|^3 = C_1 \quad \Rightarrow \quad 3xy^2 + x^3 = C\,,$$

where $C = \pm C_1$ is any nonzero constant.

14. Substituting $v = y/\theta$ yields

$$v + \theta\frac{dv}{d\theta} = \sec v + v \qquad \Rightarrow \qquad \theta\frac{dv}{d\theta} = \sec v$$
$$\Rightarrow \qquad \cos v\,dv = \frac{d\theta}{\theta} \qquad \Rightarrow \qquad \int \cos v\,dv = \int \frac{d\theta}{\theta}$$
$$\Rightarrow \qquad \sin v = \ln|\theta| + C \qquad \Rightarrow \qquad y = \theta\arcsin\left(\ln|\theta| + C\right).$$

16. We rewrite the equation in the form

$$\frac{dy}{dx} = \frac{y}{x}\left(\ln\frac{y}{x} + 1\right)$$

and substitute $v = y/x$ to get

$$v + x\frac{dv}{dx} = v\left(\ln v + 1\right) \qquad \Rightarrow \qquad x\frac{dv}{dx} = v\ln v \qquad \Rightarrow \qquad \int \frac{dv}{v\ln v} = \int \frac{dx}{x}$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

$$\Rightarrow \qquad \ln|\ln v| = \ln|x| + C_1 \qquad \Rightarrow \qquad \ln v = \pm e^{C_1} x = Cx \qquad \Rightarrow \qquad v = e^{Cx},$$

where $C \neq 0$ is any constant. Note that, separating variables, we lost a solution, $v \equiv 1$, which can be included in the above formula by letting $C = 0$. Thus we have $v = e^{Cx}$. where C is any constant. Substituting back $y = xv$ yields a general solution

$$y = xe^{Cx}$$

to the given equation.

18. With $z = x + y + 2$ and $z' = 1 + y'$, we have

$$\frac{dz}{dx} = z^2 + 1 \qquad \Rightarrow \qquad \frac{dz}{z^2+1} = dx \qquad \Rightarrow \qquad \int \frac{dz}{z^2+1} = \int dx$$
$$\Rightarrow \qquad \arctan z = x + C \qquad \Rightarrow \qquad x + y + 2 = z = \tan(x + C)$$
$$\Rightarrow \qquad y = \tan(x + C) - x - 2.$$

20. Substitution $z = x - y$ yields

$$1 - \frac{dz}{dx} = \sin z \qquad \Rightarrow \qquad \frac{dz}{dx} = 1 - \sin z \qquad \Rightarrow \qquad \frac{dz}{1-\sin z} = dx$$
$$\Rightarrow \qquad \int \frac{dz}{1-\sin z} = \int dx = x + C.$$

The left-hand side integral can be found as follows.

$$\begin{aligned} \int \frac{dz}{1-\sin z} &= \int \frac{(1+\sin z)dz}{1-\sin^2 z} = \int \frac{(1+\sin z)dz}{\cos^2 z} \\ &= \int \sec^2 z + \int \tan z \sec z \, dz = \tan z + \sec z. \end{aligned}$$

Thus, a general solution is given implicitly by

$$\tan(x - y) + \sec(x - y) = x + C.$$

22. Dividing the equation by y^3 yields

$$y^{-3}\frac{dy}{dx} - y^{-2} = e^{2x}.$$

We now make a substitution $v = y^{-2}$ so that $v' = -2y^{-3}y'$, and get

$$\frac{dv}{dx} + 2v = -2e^{2x}.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

This is a linear equation. So,

$$\mu(x) = \exp\left(\int 2dx\right) = e^{2x},$$
$$v(x) = e^{-2x}\int \left(-2e^{2x}\right)e^{2x}dx = -(1/2)e^{-2x}\left(e^{4x}+C\right) = -\frac{e^{2x}+Ce^{-2x}}{2}.$$

Therefore,

$$\frac{1}{y^2} = -\frac{e^{2x}+Ce^{-2x}}{2} \qquad \Rightarrow \qquad y = \pm\sqrt{-\frac{2}{e^{2x}+Ce^{-2x}}}.$$

Dividing the equation by y^3, we lost a constant solution $y \equiv 0$.

24. We divide this Bernoulli equation by $y^{1/2}$ and make a substitution $v = y^{1/2}$.

$$y^{-1/2}\frac{dy}{dx} + \frac{1}{x-2}\,y^{1/2} = 5(x-2)$$
$$\Rightarrow \qquad 2\frac{dv}{dx} + \frac{1}{x-2}\,v = 5(x-2) \qquad \Rightarrow \qquad \frac{dv}{dx} + \frac{1}{2(x-2)}\,v = \frac{5(x-2)}{2}.$$

An integrating factor for this linear equation is

$$\mu(x) = \exp\left[\int \frac{dx}{2(x-2)}\right] = \sqrt{|x-2|}.$$

Therefore,

$$\begin{aligned} v(x) &= \frac{1}{\sqrt{|x-2|}}\int \frac{5(x-2)\sqrt{|x-2|}}{2}\,dx \\ &= \frac{1}{\sqrt{|x-2|}}\left(|x-2|^{5/2}+C\right) = (x-2)^2 + C|x-2|^{-1/2}. \end{aligned}$$

Since $y = v^2$, we finally get

$$y = \left[(x-2)^2 + C|x-2|^{-1/2}\right]^2.$$

In addition, $y \equiv 0$ is a (lost) solution.

26. Multiplying the equation by y^2, we get

$$y^2\frac{dy}{dx} + y^3 = e^x.$$

With $v = y^3$, $v' = 3y^2y'$, the equation becomes

$$\frac{1}{3}\frac{dv}{dx} + v = e^x \qquad \Rightarrow \qquad \frac{dv}{dx} + 3v = 3e^x \qquad \Rightarrow \qquad \frac{d}{dx}\left(e^{3x}v\right) = 3e^{4x}$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

$$\Rightarrow \qquad v = e^{-3x}\int 3e^{4x}dx = \frac{3e^x}{4} + Ce^{-3x}.$$

Therefore,

$$y = \sqrt[3]{v} = \sqrt[3]{\frac{3e^x}{4} + Ce^{-3x}}\,.$$

28. First, we note that $y \equiv 0$ is a solution, which will be lost when we divide the equation by y^3 and make a substitution $v = y^{-2}$ to get a linear equation

$$y^{-3}\frac{dy}{dx} + y^{-2} + x = 0 \qquad \Rightarrow \qquad \frac{dv}{dx} - 2v = 2x.$$

An integrating factor for this equation is

$$\mu(x) = \exp\left[\int(-2)dx\right] = e^{-2x}.$$

Thus,

$$\begin{aligned} v &= e^{2x}\int 2xe^{-2x}dx = e^{2x}\left(-xe^{-2x} + \int e^{-2x}dx\right) \\ &= e^{2x}\left(-xe^{-2x} - \frac{e^{-2x}}{2} + C\right) = -x - \frac{1}{2} + Ce^{2x}. \end{aligned}$$

So,

$$y = \pm v^{-1/2} = \pm\frac{1}{\sqrt{-x - \frac{1}{2} + Ce^{2x}}}\,.$$

30. We make a substitution $x = u + h, \quad y = v + k$, where h and k satisfy system (14) in the text:

$$\begin{aligned} -4h - k - 1 &= 0 \\ h + k + 3 &= 0. \end{aligned}$$

Solving yields $h = \frac{2}{3}$, $k = -\frac{11}{3}$. Thus, $x = u + \frac{2}{3}$, $y = v - \frac{11}{3}$. Since $dx = du$, $dy = dv$, the substitution leads to

$$(-4u - v)du + (u + v)dv = 0 \qquad \Rightarrow \qquad \frac{dv}{du} = \frac{4u + v}{u + v}\,.$$

Therefore with $z = \frac{v}{u}$, $v' = z + uz'$ we have

$$\begin{aligned} z + uz' &= \frac{4 + z}{1 + z} \text{ or } u\frac{dz}{du} = \frac{4 + z}{1 + z} - z = \frac{4 - z^2}{1 + z} \\ \Rightarrow \qquad & \frac{1 + z}{4 - z^2}dz = \frac{du}{u} \end{aligned}$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

$$\Rightarrow \qquad -\frac{3}{4}\log|2-z| - \frac{1}{4}\log|z+2| = \log|u| + C_1$$
$$\Rightarrow \qquad |u||2-z|^{3/4}|z+2|^{1/4} = e^{-C_1} = C_2$$
$$\Rightarrow \qquad \left|x-\frac{2}{3}\right|\left|2-\frac{y+\frac{11}{3}}{x-\frac{2}{3}}\right|^{2/3}\left|\frac{y+\frac{11}{3}}{x-\frac{2}{3}}+2\right|^{1/4} = C_2$$

where C_1 and C_2 are constants. In dividing by $4-z^2$, we lost the constant solutions $z \equiv 2$ and $z \equiv -2$, but these are included in our answer if we let $C_2 = 0$.

32. We make a substitution $x = u + h, \quad y = v + k$, where h and k satisfy system (14) in the text:

$$2h - k + 4 = 0$$
$$h - 2k - 2 = 0.$$

Solving yields $h = -\frac{10}{3}$, $k = -\frac{8}{3}$. Thus, $x = u - \frac{10}{3}$, $y = v - \frac{8}{3}$. Since $dx = du$, $dy = dv$, the substitution leads to

$$(2u - v)du + (u - 2v)dv = 0 \qquad \Rightarrow \qquad \frac{dv}{du} = \frac{-v + 2u}{2v - u}\,.$$

Therefore with $z = \frac{v}{u}$, $v' = z + uz'$ we have

$$z + uz' = \frac{-z+2}{2z-1} \text{ or } u\frac{dz}{du} = \frac{-z+2}{2z-1} - z = \frac{-2z^2+2}{2z-1}$$
$$\Rightarrow \qquad \frac{2z-1}{z^2-1}dz = -2\frac{du}{u}$$
$$\Rightarrow \qquad \frac{1}{2}\log|1-z| + \frac{3}{2}\log|z+1| = -2\log|u| + C_1$$
$$\Rightarrow \qquad |1-z|^{\frac{1}{2}}|z+1|^{\frac{3}{2}}u^2 = \left|1-\frac{y+\frac{8}{3}}{x+\frac{10}{3}}\right|^{\frac{1}{2}}\left|1+\frac{y+\frac{8}{3}}{x+\frac{10}{3}}\right|^{\frac{3}{2}}\left(x+\frac{10}{3}\right)^2 = e^{c_1} = C_2,$$

where C_1 and C_2 are constants. In dividing by $z^2 - 1$, we lost the constant solutions $z \equiv 1$ and $z \equiv -1$, but these are included in our answer if we let $C_2 = 0$.

34. In Problem 2, we found that the given equation is of the form $dy/dx = G(y - 4x)$ with $G(u) = (u-1)^2$. Thus we make a substitution $u = y - 4x$ to get

$$\frac{dy}{dx} = (y - 4x - 1)^2 \qquad \Rightarrow \qquad 4 + \frac{du}{dx} = (u-1)^2$$
$$\Rightarrow \qquad \frac{du}{dx} = (u-1)^2 - 4 = (u-3)(u+1) \qquad \Rightarrow \qquad \int \frac{du}{(u-3)(u+1)} = \int dx\,.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

To integrate the left-hand side, we use partial fractions decomposition,

$$\frac{1}{(u-3)(u+1)} = \frac{1}{4}\left(\frac{1}{u-3} - \frac{1}{u+1}\right).$$

Thus,

$$\begin{aligned}
&\frac{1}{4}\left(\ln|u-3| - \ln|u+1|\right) = x + C_1 \qquad \Rightarrow \qquad \ln\left|\frac{u-3}{u+1}\right| = 4x + C_2 \\
&\Rightarrow \qquad \frac{u-3}{u+1} = Ce^{4x} \qquad \Rightarrow \qquad u = \frac{Ce^{4x}+3}{1-Ce^{4x}} \\
&\Rightarrow \qquad y = 4x + \frac{Ce^{4x}+3}{1-Ce^{4x}}, \qquad (2.6)
\end{aligned}$$

where $C \neq 0$ is an arbitrary constant. Separating variables, we lost the constant solutions $u \equiv 3$ and $u \equiv -1$, that is, $y = 4x+3$ and $y = 4x-1$. While $y = 4x+3$ can be obtained from (2.6) by setting $C = 0$, the solution $y = 4x - 1$ is not included in (2.6). Therefore, a general solution to the given equation is

$$y = 4x + \frac{Ce^{4x}+3}{1-Ce^{4x}} \qquad \text{and} \qquad y = 4x - 1.$$

36. This equation has linear coefficients. Thus we make a substitution $t = u+h$ and $x = v+k$ with h and k satisfying

$$\begin{aligned} h + k + 2 &= 0, \\ 3h - k - 6 &= 0 \end{aligned} \qquad \Rightarrow \qquad \begin{aligned} h &= 1, \\ k &= -3. \end{aligned}$$

As $dt = du$ and $dx = dv$, the substitution yields

$$(u+v)dv + (3u-v)du = 0 \qquad \Rightarrow \qquad \frac{du}{dv} = -\frac{u+v}{3u-v} = -\frac{(u/v)+1}{3(u/v)-1}.$$

With $z = u/v$, we have $u = vz$, $u' = z + vz'$, and the equation becomes

$$\begin{aligned}
&z + v\frac{dz}{dv} = -\frac{z+1}{3z-1} \qquad \Rightarrow \qquad v\frac{dz}{dv} = -\frac{3z^2+1}{3z-1} \\
&\Rightarrow \qquad \frac{3z-1}{3z^2+1}\,dz = -\frac{1}{v}\,dv \qquad \Rightarrow \qquad \int \frac{3z-1}{3z^2+1}\,dz = -\int \frac{1}{v}\,dv \\
&\Rightarrow \qquad \int \frac{3z\,dz}{3z^2+1} - \int \frac{dz}{3z^2+1} = -\ln|v| + C_1 \\
&\Rightarrow \qquad \frac{1}{2}\ln\left(3z^2+1\right) - \frac{1}{\sqrt{3}}\arctan\left(z\sqrt{3}\right) = -\ln|v| + C_1 \\
&\Rightarrow \qquad \ln\left[(3z^2+1)v^2\right] - \frac{2}{\sqrt{3}}\arctan\left(z\sqrt{3}\right) = C_2.
\end{aligned}$$

Making back substitution, after some algebra we get

$$\ln\left[(x+3)^2 + 3(t-1)^2\right] + \frac{2}{\sqrt{3}}\arctan\left[\frac{x+3}{\sqrt{3}(t-1)}\right] = C.$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

38. In Problem 6, we have written the equation in the form

$$\frac{dy}{dx} - y = e^{2x}y^3 \qquad \Rightarrow \qquad y^{-3}\frac{dy}{dx} - y^{-2} = e^{2x}.$$

Making a substitution $u = y^{-2}$ (and so $u' = -2y^{-3}y'$) in this Bernoulli equation, we get

$$\frac{du}{dx} + 2u = -2e^{2x} \qquad \Rightarrow \qquad \mu(x) = \exp\left(\int 2dx\right) = e^{2x}$$

$$\Rightarrow \quad \frac{d\left(e^{2x}u\right)}{dx} = -2e^{2x}e^{2x} = -2e^{4x} \qquad \Rightarrow \qquad e^{2x}u = \int \left(-2e^{4x}\right) dx = -\frac{1}{2}\,e^{4x} + C$$

$$\Rightarrow \quad u = -\frac{1}{2}\,e^{2x} + Ce^{-2x} \qquad \Rightarrow \qquad y^{-2} = -\frac{1}{2}\,e^{2x} + Ce^{-2x}\,.$$

Also, we lost the constant solution $y \equiv 0$ when divided the equation by y^3.

40. Since the equation is homogeneous, we make a substitution $u = y/\theta$. Thus, we get

$$\frac{dy}{d\theta} = -\frac{(y/\theta)^3 - (y/\theta)^2}{2(y/\theta)} \qquad \Rightarrow \qquad u + \theta\,\frac{du}{d\theta} = -\frac{u^3 - u^2}{2u} = -\frac{u^2 - u}{2}$$

$$\Rightarrow \qquad \theta\,\frac{du}{d\theta} = -\frac{u^2 + u}{2} \qquad \Rightarrow \qquad \frac{2du}{u(u+1)} = -\frac{d\theta}{\theta}$$

$$\Rightarrow \qquad \int \frac{2du}{u(u+1)} = -\int \frac{d\theta}{\theta} \qquad \Rightarrow \qquad \ln\frac{u^2}{(u+1)^2} = -\ln|\theta| + C_1\,,$$

which gives

$$\frac{u^2}{(u+1)^2} = \frac{C}{\theta}, \qquad C \neq 0\,.$$

Back substitution $u = y/\theta$ yields

$$\frac{y^2}{(\theta+y)^2} = \frac{C}{\theta} \qquad \Rightarrow \qquad \theta y^2 = C(\theta+y)^2\,, \qquad C \neq 0.$$

When $C = 0$, the above formula gives $\theta \equiv 0$ or $y \equiv 0$, which were lost in separating variables. Also, we lost another solution, $u + 1 \equiv 0$ or $y = -\theta$. Thus, the answer is

$$\theta y^2 = C(\theta+y)^2 \quad \text{and} \quad y = -\theta,$$

where C is an arbitrary constant.

42. Suggested substitution, $y = vx^2$ (so that $y' = 2xv + x^2v'$), yields

$$2xv + x^2\frac{dv}{dx} = 2vx + \cos v \qquad \Rightarrow \qquad x^2\frac{dv}{dx} = \cos v.$$

Solving this separable equation, we obtain

$$\frac{dv}{\cos v} = \frac{dx}{x^2} \qquad \Rightarrow \qquad \ln|\sec v + \tan v| = -x^{-1} + C_1$$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

$$\Rightarrow \qquad \sec v + \tan v = \pm e^{C_1} e^{-1/x} = Ce^{-1/x}$$
$$\Rightarrow \qquad \sec\left(\frac{y}{x^2}\right) + \tan\left(\frac{y}{x^2}\right) = Ce^{-1/x},$$

where $C = \pm e^{C_1}$ is an arbitrary nonzero constant. With $C = 0$, this formula also includes lost solutions

$$y = \left[\frac{\pi}{2} + (2k+1)\pi\right] x^2, \quad k = 0, \pm 1, \pm 2, \ldots .$$

So, together with the other set of lost solutions,

$$y = \left(\frac{\pi}{2} + 2k\pi\right) x^2, \quad k = 0, \pm 1, \pm 2, \ldots ,$$

we get a general solution to the given equation.

44. From

$$\frac{dy}{dx} = -\frac{a_1 x + b_1 y + c_1}{a_2 x + b_2 y + c_2},$$

using that $a_2 = ka_1$ and $b_2 = kb_1$, we obtain

$$\frac{dy}{dx} = -\frac{a_1 x + b_1 y + c_1}{ka_1 x + kb_1 y + c_2} = -\frac{a_1 x + b_1 y + c_1}{k(a_1 x + b_1 y) + c_2} = G\left(a_1 x + b_1 y\right),$$

where

$$G(t) = -\frac{t + c_1}{kt + c_2}.$$

48. Apply formula (8), page 50 of Section 2.3, to the transformed Bernoulli equation below equation (10) of Section 2.6, page 74. (If $n = 1$, the Bernoulli equation is linear and formula (8) can be applied directly.)

REVIEW PROBLEMS

2. $y = -8x^2 - 4x - 1 + Ce^{4x}$

4. $\dfrac{x^3}{6} - \dfrac{4x^2}{5} + \dfrac{3x}{4} - Cx^{-3}$

6. $y^{-2} = 2\ln|1 - x^2| + C$ and $y \equiv 0$

8. $y = (Cx^2 - 2x^3)^{-1}$ and $y \equiv 0$

10. $x + y + 2y^{1/2} + \arctan(x + y) = C$

12. $2ye^{2x} + y^3 e^x = C$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

14. $x = \dfrac{t^2(t-1)}{2} + t(t-1) + 3(t-1)\ln|t-1| + C(t-1)$

16. $y = \cos x \ln|\cos x| + C\cos x$

18. $y = 1 - 2x + \sqrt{2}\tan\left(\sqrt{2}\,x + C\right)$

20. $y = \left(C\theta^{-3} - \dfrac{12\theta^2}{5}\right)^{1/3}$

22. $(3y - 2x + 9)(y + x - 2)^4 = C$

24. $2\sqrt{xy} + \sin x - \cos y = C$

26. $y = Ce^{-x^2/2}$

28. $(y+3)^2 + 2(y+3)(x+2) - (x+2)^2 = C$

30. $y = Ce^{4x} - x - \dfrac{1}{4}$

32. $y^2 = x^2\ln(x^2) + 16x^2$

34. $y = x^2\sin x + \dfrac{2x^2}{\pi^2}$

36. $\sin(2x+y) - \dfrac{x^3}{3} + e^y = \sin 2 + \dfrac{2}{3}$

38. $y = \left[2 - \left(\dfrac{1}{4}\right)\arctan\left(\dfrac{x}{2}\right)\right]^2$

40. $y = \dfrac{8}{1 - 3e^{-4x} - 4x}$

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual

TABLES

n	x_n	y_n	n	x_n	y_n
1	0.1	1.475	6	0.6	1.353368921
2	0.2	1.4500625	7	0.7	1.330518988
3	0.3	1.425311875	8	0.8	1.308391369
4	0.4	1.400869707	9	0.9	1.287062756
5	0.5	1.376852388	10	1.0	1.266596983

Table 2–A: Euler's approximations to $y' = x - y$, $y(0) = 0$, on $[0, 1]$ with $h = 0.1$.

FIGURES




Figure 2–A: The graph of the solution in Problem 28.




Figure 2–B: The direction field and solution curve in Problem 32.

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual




Figure 2–C: The graph of the solution in Problem 32.




Figure 2–D: The direction field for Problem 40 ($\alpha = \frac{1}{4}, \beta = \frac{1}{2}$).




Figure 2–E: Curves and their orthogonal trajectories in Problem 34.

Copyright © 2018 Pearson Education, Inc.

Fundamentals of Differential Equations 9th edition Nagle Solutions Manual