# *Clinical Calculations 10th edition Kee Test Bank*

SKU: 9780323809832-TEST-BANK

**OTHER**

**Body Surface Area by Square Root**

1. Order: cyclophosphamide 500 mg/$m^2$ in 500 mL of normal saline solution (NSS) over 90 minutes
   Patient height and weight: 5 ft 10 in, 142 lb
   Drug available: cyclophosphamide 100 mg dilute with 5 mL of sterile water; yields 20 mg/mL
   a. What is the patient's body surface area (BSA) ($m^2$)?
   b. What is the total dose?
   c. How many milliliters should you prepare?

   ANS:
   a. $\sqrt{\frac{70 \times 142}{3131}} = 1.78\ m^2$
   b. 500 mg/$m^2$ × 1.78 $m^2$ = 890 mg
   c. FE: 890 mg/100 mg × 5 mL = 44.5 mL

   **OR**

   BF: $\frac{D}{H} \times V = \frac{890\ mg}{100\ mg} \times 5\ mL = 44.5\ mL$

2. Order: cisplatin 50 mg/$m^2$ in 500 mL of NSS intravenously over 90 minutes
   Patient height and weight: 5 ft 6 in, 160 lb
   Drug available: cisplatin 100 mg/100 mL
   a. What is the patient's BSA ($m^2$)?
   b. What is the total dose?
   c. How many milliliters should you prepare?


   ANS:
   a. $\sqrt{\frac{66 \times 160}{3131}} = 1.84\ m^2$
   b. 50 mg × 1.84 $m^2$ = 92 mg
   c. FE: 92 mg/100 mg × 100 mL = 92 mL

   **OR**

   BF: $\frac{D}{H} \times V = \frac{92\ mg}{100\ mg} \times 100\ mL = 92\ mL$

3. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give dacarbazine 250 mg/m$^2$/day × 5 days.
Patient height: 5 ft 10 in
Patient weight: 173 lb
What is the daily dose with
a. square root method?
b. nomogram?


ANS:

a. $\sqrt{\frac{70 \times 173}{3131}} = \sqrt{3.8677} = 1.97\,\text{m}^2$

250 mg/m$^2$/day × 1.97 m$^2$ = 493 mg/day
b. Height 70 in, weight 173 lb, intersects 2.02 m$^2$
250 mg/m$^2$/day × 2.02 m$^2$ = 505 ≈ 500 mg/day

4. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give 5-fluorouracil 450 mg/m$^2$/wk.
Patient height: 5 ft 6 in
Patient weight: 210 lb
What is the weekly dose with
a. square root method?
b. nomogram?


ANS:

a. $\sqrt{\frac{66 \times 210}{3131}} = \sqrt{4.43} = 2.10\ \text{m}^2$

450 mg/m$^2$/wk × 2.10 m$^2$ = 945 mg/wk
b. Height 66 in, weight 210 lb, intersects 2.04 m$^2$
450 mg/m$^2$/wk × 2.04 m$^2$ = 918 ≈ 920 mg/wk

5. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give leucovorin 200 mg/m$^2$/wk.
Patient height: 5 ft 6 in
Patient weight: 210 lb
What is the weekly dose with
a. square root method?
b. nomogram?


ANS:

a. $\sqrt{\frac{66 \times 210}{3131}} = \sqrt{4.43} = 2.10\ \text{m}^2$

200 mg/m$^2$/wk × 2.10 m$^2$ = 420 mg/wk
b. Height 66 in, weight 210 lb, intersects 2.04 m$^2$
200 mg/m$^2$/wk × 2.04 m$^2$ = 408.00 ≈ 400 mg/wk

6. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give cisplatin 30 mg/m$^2$/day × 3 days.
Patient height: 70 in
Patient weight: 80 kg
What is the daily dose with
a. square root method?
b. nomogram?

ANS:
a. 80 kg × 2.2 = 176 lb


$$\sqrt{\frac{70 \times 176}{3131}} = \sqrt{3.93} = 1.98\ \text{m}^2$$

30 mg/m$^2$/day × 1.98 m$^2$ = 59.4 ≈ 59 mg/day

b. Height 70 in, weight 80 kg, intersects 2.08 m$^2$
30 mg/m$^2$/day × 2.08 m$^2$ = 62.4 ≈ 62 mg/day

7. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give cisplatinum 80 mg/m$^2$/wk.
Patient height: 6 ft 2 in
Patient weight: 186 lb
What is the weekly dose with
a. square root method?
b. nomogram?


ANS:

a. $\sqrt{\frac{74 \times 186}{3131}} = \sqrt{4.39} = 2.09\ \text{m}^2$

80 mg/m$^2$/wk × 2.09 m$^2$ = 167.2 mg/wk
b. Height 74 in, weight 186 lb, intersects 2.10 m$^2$
80 mg/m$^2$/wk × 2.10 m$^2$ = 168 ≈ 170 mg/wk

8. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give etoposide 120 mg/m$^2$/wk.
Patient height: 74 in
Patient weight: 70 kg
What is the weekly dose with
a. square root method?
b. nomogram?


ANS:
a. 70 kg × 2.2 = 154 lb

$$\sqrt{\frac{74 \times 154}{3131}} = \sqrt{3.639} = 1.91 \text{ m}^2$$

120 mg/m$^2$/wk × 1.91 m$^2$ = 229.2 mg/wk
b. Height 74 in, weight 70 kg, intersects 2.06 m$^2$
120 mg/m$^2$/wk × 2.06 m$^2$ = 247.2 ≈ 250 mg/wk

9. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give Cytoxan 600 mg/m$^2$/wk.
Patient height: 70 in
Patient weight: 85 kg
What is the weekly dose with
a. square root method?
b. nomogram?

ANS:


a. 85 kg × 2.2 = 187 lb

$$\sqrt{\frac{70 \times 187}{3131}} = \sqrt{4.18} = 2.04 \text{ m}^2$$

600 mg/m$^2$/wk × 2.04 m$^2$ = 1224 or 1225 mg/wk
b. Height 70 in, weight 85 kg, intersects 2.08 m$^2$
600 mg/m$^2$/wk × 2.08 m$^2$ = 1248 ≈ 1250 mg/wk

10. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give Adriamycin 60 mg/m$^2$/wk.
Patient height: 70 in
Patient weight: 80 kg
What is the weekly dose with
a. square root method?
b. nomogram?


ANS:
a. 80 kg × 2.2 = 176 lb

$$\sqrt{\frac{70 \times 176}{3131}} = \sqrt{3.93} = 1.98\ \text{m}^2$$

60 mg/m$^2$/wk × 1.98 m$^2$ = 118.8 mg/wk = 119 mg/wk
b. Height 70 in, weight 80 kg, intersects 2.04 m$^2$
60 mg/m$^2$/wk × 2.04 m$^2$ = 122.4 mg/wk

11. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give vincristine 2 mg/m$^2$/week.
Patient height: 62 in
Patient weight: 75 kg
What is the weekly dose with
a. square root method?
b. nomogram?


ANS:
a. 75 kg × 2.2 = 165 lb

$$\sqrt{\frac{62 \times 165}{3131}} = \sqrt{3.27} = 1.81 \text{ m}^2$$

2 mg/m$^2$/wk × 1.81 m$^2$ = 3.62 = 3.6 mg/wk
b. Height 62 in, weight 75 kg, intersects 1.78 m$^2$
2 mg/m$^2$/wk × 1.78 m$^2$ = 3.56 ≈ 3.6 mg/wk

12. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give mitomycin 15 mg/m$^2$/week.
Patient height: 65 in
Patient weight: 64 kg
What is the weekly dose with
a. square root method?
b. nomogram?


ANS:
a. 64 kg × 2.2 = 140.8 or 141 lb

$$\sqrt{\frac{65 \times 141}{3131}} = \sqrt{2.93} = 1.71 \text{ m}^2$$

15 mg/m$^2$/wk × 1.71 m$^2$ = 25.6 or 26 mg/wk
b. Height 65 in, weight 64 kg, intersects 1.75 m$^2$
15 mg/m$^2$/wk × 1.75 m$^2$ = 26.25 ≈ 26 mg/wk

13. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give mitoxantrone 12 mg/m$^2$/day × 3 days.
Patient height: 5 ft 8 in
Patient weight: 150 lb
What is the daily dose with
a. square root method?
b. nomogram?


ANS:

a. $\sqrt{\frac{68 \times 150}{3131}} = \sqrt{3.25} = 1.8\ \text{m}^2$

12 mg/m$^2$/day × 1.80 m$^2$ = 21.6 or 22 mg/day
b. Height 68 in, weight 150 lb, intersects 1.85 m$^2$
12 mg/m$^2$/day × 1.85 m$^2$ = 22.2 ≈ 22 mg/day

14. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give cytosine arabinoside 100 mg/m$^2$/day × 7 days.
Patient height: 5 ft 2 in
Patient weight: 130 lb
What is the weekly dose with
a. square root method?
b. nomogram?


ANS:

a. $\sqrt{\frac{62 \times 130}{3131}} = \sqrt{2.57} = 1.6\ \text{m}^2$

100 mg/m$^2$/day × 1.6 m$^2$ = 160 mg/day
b. Height 62 in, weight 130 lb, intersects 1.65 m$^2$
100 mg/m$^2$/day × 1.65 m$^2$ = 165 mg/day

15. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give methotrexate 3.3 mg/m$^2$/day × 7 days.
Patient height: 72 in
Patient weight: 82 kg
What is the daily dose with
a. square root method?
b. nomogram?


ANS:
a. 82 kg × 2.2 = 180.4 lb

$$\sqrt{\frac{72 \times 180.4}{3131}} = \sqrt{4.15} = 2.04 \text{ m}^2$$

3.3 mg/m$^2$/day × 2.04 m$^2$ = 6.7 mg/day
b. Height 72 in, weight 82 kg, intersects 2.10 m$^2$
3.3 mg/m$^2$/day × 2.10 m$^2$ = 6.93 ≈ 6.9 mg/day

16. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give prednisone 60 mg/m$^2$/day × 7 days.
Patient height: 72 in
Patient weight: 84 kg
What is the daily dose with
a. square root method?
b. nomogram?


ANS:
a. 84 kg × 2.2 = 184.8 or 185 lb

$$\sqrt{\frac{72 \times 185}{3131}} = \sqrt{4.25} = 2.06\ \text{m}^2$$

60 mg/m$^2$/day × 2.06 m$^2$ = 123.6 or 124 mg/day
b. Height 72 in, weight 84 kg, intersects 2.10 m$^2$
60 mg/m$^2$/day × 2.10 m$^2$ = 126 mg/day

17. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give idarubicin hydrochloride 12 mg/m$^2$/day × 3 days.
Patient height: 60 in
Patient weight: 60 kg
What is the daily dose with
a. square root method?
b. nomogram?


ANS:
a. 60 kg × 2.2 = 132 lb

$$\sqrt{\frac{60 \times 132}{3131}} = \sqrt{2.53} = 1.59 \text{ m}^2$$

12 mg/m$^2$/day × 1.59 m$^2$ = 19.1 or 19 mg/day
b. Height 60 in, weight 60 kg, intersects 1.60 m$^2$
12 mg/m$^2$/day × 1.60 m$^2$ = 19.2 ≈ 19 mg/day

18. For the following questions, use the square root method and/or nomogram. Note discrepancies between methods.

Clinical Calculations 10th edition Kee Test Bank




Give cytarabine 100 mg/m$^2$/day × 7 days.
Patient height: 64 in
Patient weight: 65 kg
What is the daily dose with
a. square root method?
b. nomogram?

ANS:
a. 65 kg × 2.2 = 143 lb

$$\sqrt{\frac{64 \times 143}{3131}} = \sqrt{2.92} = 1.71 \text{ m}^2$$

100 mg/m$^2$/day × 1.71 m$^2$ = 171 mg/day
b. Height 64 in, weight 60 kg, intersects 1.69 m$^2$
100 mg/m$^2$/day × 1.69 m$^2$ = 169 ≈ 170 mg/day

19. Order: streptozocin 1000 mg/m$^2$ in 100 mL $D_5W$ over 2 hours
Patient's height and weight: 5 ft 2 in and 210 lb

Clinical Calculations 10th edition Kee Test Bank

Drug available: streptozocin 1 g powdered vial, reconstitute with 9.5 mL NS; yields 100 mg/mL
a. What is the patient's BSA ($m^2$)?
b. What is the total dose?
c. How many milliliters should you prepare?

ANS:
a. $\sqrt{\frac{62 \times 210}{3131}} = \sqrt{4.16} = 2.04\ m^2$
b. 2.04 $m^2$ × 1000 mg/$m^2$ = 2040 mg
c. BF: $\frac{D}{H} \times V = \frac{2040\ mg}{100\ mg \times 1\ mL} = 20.4\ mL$

20. Order: methotrexate 3 mg/$m^2$ PO × 2 weekly
Patient's height and weight: 5 ft 2 in and 130 lb
Drug available: methotrexate tablets 2.5, 5, 7.5 mg
a. What is the patient's BSA ($m^2$)?
b. What is the total dose?

ANS:
a. $\sqrt{\frac{62 \times 130}{3131}} = \sqrt{\frac{8060}{3131}} = \sqrt{2.57} = 1.60\ m^2$
b. 1.6 $m^2$ × 3 mg/$m^2$ = 4.8 mg or 5 mg tablet twice a week

21. Order: sargramostim 250 mcg/$m^2$/day, dilute in 50 mL and infuse over 2 hours
Patient's height and weight: 5 ft 10 in and 285 lb
Drug available: sargramostim 500 mcg/mL
a. What is the patient's BSA ($m^2$)?
b. What is the total dose?
c. How many milliliters should you prepare?

ANS:
a. $\sqrt{\frac{70 \times 285}{3131}} = \sqrt{\frac{19950}{3131}} = \sqrt{6.37} = 2.52\ m^2$
b. 2.52 $m^2$ × 250 mcg/$m^2$ = 630 mcg
c. $\frac{630\ mcg}{500\ mcg} \times 1\ mL = 1.26\ mL$

22. Order: clofarabine 52 mg/$m^2$ daily and infuse in 250 mL $D_5W$ over 2 hours
Patient's height and weight: 5 ft 4 in and 115 lb
Drug available: 20 mg/20 mL
a. What is the patient's BSA ($m^2$)?
b. What is the total dose?
c. How many milliliters should you prepare?

ANS:
a. $\sqrt{\frac{64 \times 115}{3131}} = \sqrt{\frac{7360}{3131}} = \sqrt{2.35} = 1.53\ m^2$

Clinical Calculations 10th edition Kee Test Bank

b. $1.53\ m^2 \times 52\ mg/m^2 = 79.56$ or 80 mg

c. $\frac{80\ mcg}{20\ mcg} \times 20\ mL = 80\ mL$