Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

SKU: 9781119169789-SOLUTIONS

Chapter Review Sheets for
Elementary Differential Equations and Boundary Value Problems, 11e

Chapter 2: First Order Differential Equations

Definitions:

- First-order linear differential equation
- Integrating factor, integral curves
- Variation of parameters
- Separable, differential form
- Homogeneous differential equations
- Implicit solutions
- Bernoulli equations
- Logistic equations, intrinsic growth rate
- Existence and uniqueness of solutions, general solutions
- Autonomous, logistic equation, rate of growth or decline
- Intrinsic growth rate, equilibrium solutions, critical points
- Threshold level, phase line, saturation level or carrying capacity
- Stable solutions, asymptotically stable solutions, unstable equilibrium solution
- Integrating factors, equations
- Tangent line method (Euler's method)
- First order difference equation
- Method of successive approximations

Theorems:

- Theorem 2.4.1: Existence and uniqueness of solutions to linear first order ODE's.
- Theorem 2.4.2: Existence and uniqueness of solutions to first order IVP's.
- Theorem 2.6.1: Existence and uniqueness of solutions to exact first order ODE's.
- Theorem 2.8.1: Restatement and elaboration of Theorem 2.4.2.

Important Skills:

- Be able to determine if a first order differential equation is linear or nonlinear. (Sec. 2.1)
- If the differential equation is linear, compute the integrating factor, and then the general solution. (Sec. 2.1, Ex. 4)
- Be able to graph integral curves for an ODE. (Sec. 2.1, Ex. 4)
- If it's nonlinear, determine if it is separable. (Sec. 2.1, Ex. 1)
- It is crucial to know integration of basic functions and integral methods from calculus. Various substitutions, integration by parts, and partial fractions will all be utilized. (Techniques are used in most examples.)
- Understand the three steps in the process of mathematical modeling: construction of the model, analysis of the model, and comparison with experiment or observation. (Section 2.3, Ex. 1, 2, 3, 4)
- If it isn't separable or exact, check for substitutions that would convert it into a linear equation, nonlinear equation that is then separable. (Sec. 2.4)
- Determine the existence and uniqueness of solutions to differential equations. (Sec. 2.4, Ex. 1, 2)
- These important qualitative classifications are at the heart of dynamical systems. Important with this is the concept of a threshold value. (Sec. 2.5)

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

- Be able to determine what happens to solutions as time tends to infinity. Understand stability, asymptotic stability and instability. (Sec. 2.5)
- Know how to recognize autonomous equations, and utilize the direction field to represent solution to them. Be able to determine asymptotically stable, semi-stable, and unstable equilibrium solutions. (Sec. 2.5, Ex. 1)
- If the differential equation is not separable, determine if it is exact. (Sec. 2.6, Ex. 2)
- Recognize that Bernoulli equations can be transformed into linear equations.
- Know how to obtain approximate solutions using Euler's method if an analytical solution cannot be found. (Sec. 2.7, Ex. 2)

Relevant Applications:

- Mixing problems, compound interest, motion in a gravitational field, radioactive carbon dating, population dynamics

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual


CHAPTER

2

First-Order Differential Equations

2.1

5.(a)




(b) If $y(0) > -3$, solutions eventually have positive slopes, and hence increase without bound. If $y(0) \le -3$, solutions have negative slopes and decrease without bound.

(c) The integrating factor is $\mu(t) = e^{-\int 2dt} = e^{-2t}$. The differential equation can be written as $e^{-2t}y' - 2e^{-2t}y = 3e^{-t}$, that is, $(e^{-2t}y)' = 3e^{-t}$. Integration of both sides of the equation results in the general solution $y(t) = -3e^{t} + c\,e^{2t}$. It follows that all solutions will increase exponentially if $c > 0$ and will decrease exponentially

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

if $c \le 0$. Letting $c = 0$ and then $t = 0$, we see that the boundary of these behaviors is at $y(0) = -3$.


6.(a)



(b) For $y > 0$, the slopes are all positive, and hence the corresponding solutions increase without bound. For $y < 0$, almost all solutions have negative slopes, and hence solutions tend to decrease without bound.

(c) First divide both sides of the equation by t ($t > 0$). From the resulting standard form, the integrating factor is $\mu(t) = e^{-\int (1/t)\,dt} = 1/t$. The differential equation can be written as $y'/t - y/t^2 = t\,e^{-t}$, that is, $(y/t)' = t\,e^{-t}$. Integration leads to the general solution $y(t) = -te^{-t} + c\,t$. For $c \neq 0$, solutions diverge, as implied by the direction field. For the case $c = 0$, the specific solution is $y(t) = -te^{-t}$, which evidently approaches zero as $t \to \infty$.

8.(a)



(b) All solutions eventually have positive slopes, and hence increase without bound.


(c) The integrating factor is $\mu(t) = e^{t/2}$. The differential equation can be written as $e^{t/2}y' + e^{t/2}y/2 = 3t^2/2$, that is, $(e^{t/2}\,y/2)' = 3t^2/2$. Integration of both sides of the equation results in the general solution $y(t) = 3t^2 - 12t + 24 + c\,e^{-t/2}$. It follows that all solutions converge to the specific solution $3t^2 - 12t + 24$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

10. The integrating factor is $\mu(t) = e^{2t}$. After multiplying both sides by $\mu(t)$, the equation can be written as $(e^{2t}\, y)' = t$. Integrating both sides of the equation results in the general solution $y(t) = t^2 e^{-2t}/2 + c\, e^{-2t}$. Invoking the specified condition, we require that $e^{-2}/2 + c\, e^{-2} = 0$. Hence $c = -1/2$, and the solution to the initial value problem is $y(t) = (t^2 - 1)e^{-2t}/2$.

11. The integrating factor is $\mu(t) = e^{\int (2/t)\, dt} = t^2$. Multiplying both sides by $\mu(t)$, the equation can be written as $(t^2\, y)' = \cos t$. Integrating both sides of the equation results in the general solution $y(t) = \sin t/t^2 + c\, t^{-2}$. Substituting $t = \pi$ and setting the value equal to zero gives $c = 0$. Hence the specific solution is $y(t) = \sin t/t^2$.

14.(a)




Solutions appear to grow infinitely large in absolute value, with signs depending on the initial value $y(0) = a_0$. The direction field appears horizontal for $a_0 \approx -1/8$.

(b) Dividing both sides of the given equation by 3, the integrating factor is $\mu(t) = e^{-2t/3}$. Multiplying both sides of the original differential equation by $\mu(t)$ and integrating results in $y(t) = (2\, e^{2t/3} - 2\, e^{-\pi t/2} + a(4 + 3\pi)\, e^{2t/3})/(4 + 3\pi)$. The qualitative behavior of the solution is determined by the terms containing $e^{2t/3}$: $2\, e^{2t/3} + a(4 + 3\pi)\, e^{2t/3}$. The nature of the solutions will change when $2 + a(4 + 3\pi) = 0$. Thus the critical initial value is $a_0 = -2/(4 + 3\pi)$.

(c) In addition to the behavior described in part (a), when $y(0) = -2/(4 + 3\pi)$, the solution is $y(t) = (-2\, e^{-\pi t/2})/(4 + 3\pi)$, and that specific solution will converge to $y = 0$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

15.(a)



As $t \to 0$, solutions increase without bound if $y(1) = a > 0.4$, and solutions decrease without bound if $y(1) = a < 0.4$.

(b) The integrating factor is $\mu(t) = e^{\int (t+1)/t\,dt} = t\,e^t$. The general solution of the differential equation is $y(t) = t\,e^{-t} + c\,e^{-t}/t$. Since $y(1) = a$, we have that $1 + c = ae$. That is, $c = ae - 1$. Hence the solution can also be expressed as $y(t) = t\,e^{-t} + (ae - 1)\,e^{-t}/t$. For small values of t, the second term is dominant. Setting $ae - 1 = 0$, the critical value of the parameter is $a_0 = 1/e$.

(c) When $a = 1/e$, the solution is $y(t) = t\,e^{-t}$, which approaches 0 as $t \to 0$.

17. The integrating factor is $\mu(t) = e^{\int (1/2)\,dt} = e^{t/2}$. Therefore the general solution is $y(t) = (4\cos t + 8\sin t)/5 + c\,e^{-t/2}$. Invoking the initial condition, the specific solution is $y(t) = (4\cos t + 8\sin t - 9\,e^{-t/2})/5$. Differentiating, it follows that $y'(t) = (-4\sin t + 8\cos t + 4.5\,e^{-t/2})/5$ and $y''(t) = (-4\cos t - 8\sin t - 2.25\,e^{-t/2})/5$. Setting $y'(t) = 0$, the first solution is $t_1 = 1.3643$, which gives the location of the first stationary point. Since $y''(t_1) < 0$, the first stationary point in a local maximum. The coordinates of the point are $(1.3643, 0.82008)$.

18. The integrating factor is $\mu(t) = e^{\int (2/3)\,dt} = e^{2t/3}$, and the differential equation can be written as $(e^{2t/3}\,y)' = e^{2t/3} - t\,e^{2t/3}/2$. The general solution is $y(t) = (21 - 6t)/8 + c\,e^{-2t/3}$. Imposing the initial condition, we have $y(t) = (21 - 6t)/8 + (y_0 - 21/8)e^{-2t/3}$. Since the solution is smooth, the desired intersection will be a point of tangency. Taking the derivative, $y'(t) = -3/4 - (2y_0 - 21/4)e^{-2t/3}/3$. Setting $y'(t) = 0$, the solution is $t_1 = (3/2)\ln[(21 - 8y_0)/9]$. Substituting into the solution, the respective value at the stationary point is $y(t_1) = 3/2 + (9/4)\ln 3 - (9/8)\ln(21 - 8y_0)$. Setting this result equal to zero, we obtain the required initial value $y_0 = (21 - 9\,e^{4/3})/8 \approx -1.643$.

19.(a) The integrating factor is $\mu(t) = e^{t/4}$, and the differential equation can be written as $(e^{t/4}\,y)' = 3\,e^{t/4} + 2\,e^{t/4}\cos 2t$. After integration, we get that the general solution is $y(t) = 12 + (8\cos 2t + 64\sin 2t)/65 + ce^{-t/4}$. Invoking the initial condition, $y(0) = 0$, the specific solution is $y(t) = 12 + (8\cos 2t + 64\sin 2t - 788\,e^{-t/4})/65$. As $t \to \infty$, the exponential term will decay, and the solution will oscillate about an average value of 12, with an amplitude of $8/\sqrt{65}$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

(b) Solving $y(t) = 12$, we obtain the desired value $t \approx 10.0658$.

21. The integrating factor is $\mu(t) = e^{-3t/2}$, and the differential equation can be written as $(e^{-3t/2}\ y)' = 3t\, e^{-3t/2} + 2\, e^{-t/2}$. The general solution is $y(t) = -2t - 4/3 - 4\, e^t + c\, e^{3t/2}$. Imposing the initial condition, $y(t) = -2t - 4/3 - 4\, e^t + (y_0 + 16/3)\, e^{3t/2}$. Now as $t \to \infty$, the term containing $e^{3t/2}$ will dominate the solution. Its sign will determine the divergence properties. Hence the critical value of the initial condition is $y_0 = -16/3$. The corresponding solution, $y(t) = -2t - 4/3 - 4\, e^t$, will also decrease without bound.

Note on Problems 24-27:

Let $g(t)$ be given, and consider the function $y(t) = y_1(t) + g(t)$, in which $y_1(t) \to 0$ as $t \to \infty$. Differentiating, $y'(t) = y_1'(t) + g'(t)$. Letting a be a constant, it follows that $y'(t) + ay(t) = y_1'(t) + ay_1(t) + g'(t) + ag(t)$. Note that the hypothesis on the function $y_1(t)$ will be satisfied, if $y_1'(t) + ay_1(t) = 0$. That is, $y_1(t) = c\, e^{-at}$. Hence $y(t) = c\, e^{-at} + g(t)$, which is a solution of the equation $y' + ay = g'(t) + ag(t)$. For convenience, choose $a = 1$.

24. Here $g(t) = 3$, and we consider the linear equation $y' + y = 3$. The integrating factor is $\mu(t) = e^t$, and the differential equation can be written as $(e^t\, y)' = 3e^t$. The general solution is $y(t) = 3 + c\, e^{-t}$.

26. Here $g(t) = 2t - 5$. Consider the linear equation $y' + y = 2 + 2t - 5$. The integrating factor is $\mu(t) = e^t$, and the differential equation can be written as $(e^t\, y)' = (2t - 3)e^t$. The general solution is $y(t) = 2t - 5 + c\, e^{-t}$.

27. $g(t) = 4 - t^2$. Consider the linear equation $y' + y = 4 - 2t - t^2$ .The integrating factor is $\mu(t) = e^t$, and the equation can be written as $(e^t\, y)' = (4 - 2t - t^2)e^t$. The general solution is $y(t) = 4 - t^2 + c\, e^{-t}$.

28.(a) Differentiating y and using the fundamental theorem of calculus we obtain that $y' = Ae^{-\int p(t)dt} \cdot (-p(t))$, and then $y' + p(t)y = 0$.

(b) Differentiating y we obtain that

$$y' = A'(t)e^{-\int p(t)dt} + A(t)e^{-\int p(t)dt} \cdot (-p(t)).$$

If this satisfies the differential equation then

$$y' + p(t)y = A'(t)e^{-\int p(t)dt} = g(t)$$

and the required condition follows.

(c) Let us denote $\mu(t) = e^{\int p(t)dt}$. Then clearly $A(t) = \int \mu(t)g(t)dt$, and after substitution $y = \int \mu(t)g(t)dt \cdot (1/\mu(t))$, which is just Eq. (33).

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

30. We assume a solution of the form $y = A(t)e^{-\int(1/t)\,dt} = A(t)e^{-\ln t} = A(t)t^{-1}$, where $A(t)$ satisfies $A'(t) = 3t\cos 2t$. This implies that

$$A(t) = \frac{3\cos 2t}{4} + \frac{3t\sin 2t}{2} + c$$

and the solution is

$$y = \frac{3\cos 2t}{4t} + \frac{3\sin 2t}{2} + \frac{c}{t}.$$

2.2

Problems 1 through 16 follow the pattern of the examples worked in this section. The first eight problems, however, do not have an initial condition, so the integration constant c cannot be found.

2. The differential equation may be written as $y^{-2}dy = -\sin x\,dx$. Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation $-y^{-1} = \cos x + c$. That is, $(c - \cos x)y = 1$, in which c is an arbitrary constant. Solving for the dependent variable, explicitly, $y(x) = 1/(c - \cos x)$.

3. Write the differential equation as $\cos^{-2} 2y\,dy = \cos^2 x\,dx$, which also can be written as $\sec^2 2y\,dy = \cos^2 x\,dx$. Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation $\tan 2y = \sin x\cos x + x + c$.

5. The differential equation may be written as $(y + e^y)dy = (x - e^{-x})dx$. Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation $y^2 + 2e^y = x^2 + 2e^{-x} + c$.

6. Write the differential equation as $(1 + y^2)dy = x^2\,dx$. Integrating both sides of the equation, we obtain the relation $y + y^3/3 = x^3/3 + c$.


7. Write the differential equation as $y^{-1}\,dy = x^{-1}\,dx$. Integrating both sides of the equation, we obtain the relation $\ln|y| = \ln|x| + c$. Solving for y explicitly gives $y(x) = kx$. Note that k may be positive or negative due to the absolute values in the integrated equation.

8. Write the differential equation as $y\,dy = -x\,dx$. Integrating both sides of the equation, we obtain the relation $(1/2)y^2 = -(1/2)x^2 + c$. The explicit form of the solution is $y(x) = \pm\sqrt{x^2 + c}$. The initial condition would then be used to determine whether the positive of negative solution is to be used for a specific initial value problem.

9.(a) The differential equation is separable, with $y^{-2}dy = (1 - 2x)dx$. Integration yields $-y^{-1} = x - x^2 + c$. Substituting $x = 0$ and $y = -1/6$, we find that $c = 6$. Hence the specific solution is $y = 1/(x^2 - x - 6)$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual


(b)



(c) Note that $x^2 - x - 6 = (x + 2)(x - 3)$. Hence the solution becomes singular at $x = -2$ and $x = 3$, so the interval of existence is $(-2, 3)$.

12.(a) Write the differential equation as $r^{-2}dr = \theta^{-1}\,d\theta$. Integrating both sides of the equation results in the relation $-r^{-1} = \ln\theta + c$. Imposing the condition $r(1) = 2$, we obtain $c = -1/2$. The explicit form of the solution is $r = 2/(1 - 2\ln\theta)$.

(b)



(c) Clearly, the solution makes sense only if $\theta > 0$. Furthermore, the solution becomes singular when $\ln\theta = 1/2$, that is, $\theta = \sqrt{e}$.

18. The differential equation can be written as $(3y^2 - 4)dy = 3x^2dx$. Integrating both sides, we obtain $y^3 - 4y = x^3 + c$. Imposing the initial condition, the specific solution is $y^3 - 4y = x^3 - 1$. Referring back to the differential equation, we find that $y' \to \infty$ as $y \to \pm 2/\sqrt{3}$. The respective values of the abscissas are $x \approx -1.276$, 1.598. Hence the solution is valid for $-1.276 < x < 1.598$.

22.(a) Write the differential equation as $y^{-1}(4 - y)^{-1}dy = t(1 + t)^{-1}dt$. Integrating both sides of the equation, we obtain $\ln|y| - \ln|y - 4| = 4t - 4\ln|1 + t| + c$. Taking the exponential of both sides $|y/(y - 4)| = c\,e^{4t}/(1 + t)^4$. It follows that as $t \to \infty$, $|y/(y - 4)| = |1 + 4/(y - 4)| \to \infty$. That is, $y(t) \to 4$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual


(b) Setting $y(0) = 2$, we obtain that $c = 1$. Based on the initial condition, the solution may be expressed as $y/(y-4) = -e^{4t}/(1+t)^4$. Note that $y/(y-4) < 0$, for all $t \geq 0$. Hence $y < 4$ for all $t \geq 0$. Referring back to the differential equation, it follows that y' is always positive. This means that the solution is monotone increasing. We find that the root of the equation $e^{4t}/(1+t)^4 = 399$ is near $t = 2.844$.

(c) Note the $y(t) = 4$ is an equilibrium solution. Examining the local direction field we see that if $y(0) > 0$, then the corresponding solutions converge to $y = 4$. Referring back to part (a), we have $y/(y-4) = [y_0/(y_0-4)]\, e^{4t}/(1+t)^4$, for $y_0 \neq 4$. Setting $t = 2$, we obtain $y_0/(y_0-4) = (3/e^2)^4 y(2)/(y(2)-4)$. Now since the function $f(y) = y/(y-4)$ is monotone for $y < 4$ and $y > 4$, we need only solve the equations $y_0/(y_0-4) = -399(3/e^2)^4$ and $y_0/(y_0-4) = 401(3/e^2)^4$. The respective solutions are $y_0 = 3.6622$ and $y_0 = 4.4042$.

29.(a) Observe that $-(4x+3y)/(2x+y) = -2 - (y/x)\,[2 + (y/x)]^{-1}$. Hence the differential equation is homogeneous.

(b) The substitution $y = x\,v$ results in $v + x\,v' = -2 - v/(2+v)$. The transformed equation is $v' = -(v^2+5v+4)/(2+v)x$. This equation is separable, with general solution $(v+4)^2\,|v+1| = c/x^3$. In terms of the original dependent variable, the solution is $(4x+y)^2\,|x+y| = c$.

(c) The integral curves are symmetric with respect to the origin.




30.(a) The differential equation can be expressed as $y' = (1/2)(y/x)^{-1} - (3/2)(y/x)$. Hence the equation is homogeneous. The substitution $y = xv$ results in $xv' = (1-5v^2)/2v$. Separating variables, we have $2v\,dv/(1-5v^2) = dx/x$.

(b) Integrating both sides of the transformed equation yields $-(\ln|1-5v^2|)/5 = \ln|x| + c$, that is, $1 - 5v^2 = c/|x|^5$. In terms of the original dependent variable, the general solution is $5y^2 = x^2 - c/|x|^3$.

(c) The integral curves are symmetric with respect to the origin.


Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual



31.(a) The differential equation can be expressed as $y' = (3/2)(y/x) - (1/2)(y/x)^{-1}$. Hence the equation is homogeneous. The substitution $y = x\,v$ results in $x\,v' = (v^2 - 1)/2v$, that is, $2vdv/(v^2 - 1) = dx/x$.

(b) Integrating both sides of the transformed equation yields $\ln\left|v^2 - 1\right| = \ln|x| + c$, that is, $v^2 - 1 = c\,|x|$. In terms of the original dependent variable, the general solution is $y^2 = c\,x^2\,|x| + x^2$.

(c) The integral curves are symmetric with respect to the origin.



2.3

1. Let $Q(t)$ be the amount of dye in the tank at time t. Clearly, $Q(0) = 200$ g. The differential equation governing the amount of dye is $Q'(t) = -2Q(t)/200$. The solution of this separable equation is $Q(t) = Q(0)e^{-t/100} = 200e^{-t/100}$. We need the time T such that $Q(T) = 2$ g. This means we have to solve $2 = 200e^{-T/100}$ and we obtain that $T = -100\ln(1/100) = 100\ln 100 \approx 460.5$ min.

3.(a) Let Q be the amount of salt in the tank. Salt enters the tank of water at a rate of $2\,(1/4)(1 + (1/2)\sin t) = 1/2 + (1/4)\sin t$ oz/min. It leaves the tank at a rate of $2\,Q/100$ oz/min. Hence the differential equation governing the amount of


Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

salt at any time is

$$\frac{dQ}{dt} = \frac{1}{2} + \frac{1}{4}\sin t - \frac{Q}{50}\,.$$

The initial amount of salt is $Q_0 = 50$ oz. The governing differential equation is linear, with integrating factor $\mu(t) = e^{t/50}$. Write the equation as $(e^{t/50}Q)' = e^{t/50}(1/2 + (1/4)\sin t)$. The specific solution is $Q(t) = 25 + (12.5\sin t - 625\cos t + 63150\,e^{-t/50})/2501$ oz.

(b)



(c) The amount of salt approaches a steady state, which is an oscillation of approximate amplitude 1/4 about a level of 25 oz.

4.(a) Using the Principle of Conservation of Energy, the speed v of a particle falling from a height h is given by

$$\frac{1}{2}mv^2 = mgh\,.$$

(b) The outflow rate is (outflow cross-section area)×(outflow velocity): $\alpha\, a\sqrt{2gh}$. At any instant, the volume of water in the tank is $V(h) = \int_0^h A(u)du$. The time rate of change of the volume is given by $dV/dt = (dV/dh)(dh/dt) = A(h)dh/dt$. Since the volume is decreasing, $dV/dt = -\alpha\, a\sqrt{2gh}$.

(c) With $A(h) = \pi$, $a = 0.01\,\pi$, $\alpha = 0.6$, the differential equation for the water level h is $\pi(dh/dt) = -0.006\,\pi\sqrt{2gh}$, with solution $h(t) = 0.000018gt^2 - 0.006\sqrt{2gh(0)}\,t + h(0)$. Setting $h(0) = 3$ and $g = 9.8$, $h(t) = 0.0001764\,t^2 - 0.046\,t + 3$, resulting in $h(t) = 0$ for $t \approx 130.4$ s.

5.(a) The equation governing the value of the investment is $dS/dt = r\,S$. The value of the investment, at any time, is given by $S(t) = S_0e^{rt}$. Setting $S(T) = 2S_0$, the required time is $T = \ln(2)/r$.

(b) For the case $r = .07$, $T \approx 9.9$ yr.

(c) Referring to part (a), $r = \ln(2)/T$. Setting $T = 8$, the required interest rate is to be approximately $r = 8.66\%$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

8.(a) Using Eq.(15) we have $dS/dt - 0.005S = -(800 + 10t)$, $S(0) = 150,000$. Using an integrating factor and integration by parts we obtain that $S(t) = 560,000 - 410,000e^{0.005t} + 2000t$. Setting $S(t) = 0$ and solving numerically for t yields $t = 146.54$ months.


(b) The solution we obtained in part (a) with a general initial condition $S(0) = S_0$ is $S(t) = 560,000 - 560,000e^{0.005t} + S_0e^{0.005t} + 2000t$. Solving the equation $S(240) = 0$ yields $S_0 = 246,758$.

9.(a) Let $Q' = -r\,Q$. The general solution is $Q(t) = Q_0e^{-rt}$. Based on the definition of half-life, consider the equation $Q_0/2 = Q_0e^{-5730\,r}$. It follows that $-5730\,r = \ln(1/2)$, that is, $r = 1.2097 \times 10^{-4}$ per year.

(b) The amount of carbon-14 is given by $Q(t) = Q_0\,e^{-1.2097\times10^{-4}t}$.

(c) Given that $Q(T) = Q_0/5$, we have the equation $1/5 = e^{-1.2097\times10^{-4}T}$. Solving for the decay time, the apparent age of the remains is approximately $T = 13,305$ years.

11.(a) The differential equation $dy/dt = r(t)\,y - k$ is linear, with integrating factor $\mu(t) = e^{-\int r(t)dt}$. Write the equation as $(\mu\,y)' = -k\,\mu(t)$. Integration of both sides yields the general solution $y = \left[-k\int \mu(\tau)d\tau + y_0\,\mu(0)\right]/\mu(t)$. In this problem, the integrating factor is $\mu(t) = e^{(\cos t - t)/5}$.



(b) The population becomes extinct, if $y(t^*) = 0$, for some $t = t^*$. Referring to part (a), we find that $y(t^*) = 0$ when

$$\int_0^{t^*} e^{(\cos\,\tau - \tau)/5}d\tau = 5\,e^{1/5}y_c.$$

It can be shown that the integral on the left hand side increases monotonically, from zero to a limiting value of approximately 5.0893. Hence extinction can happen only if $5\,e^{1/5}y_0 < 5.0893$. Solving $5e^{1/5}y_c = 5.0893$ yields $y_c = 0.8333$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

(c) Repeating the argument in part (b), it follows that $y(t^*) = 0$ when


$$\int_0^{t^*} e^{(\cos\,\tau - \tau)/5} d\tau = \frac{1}{k}\, e^{1/5} y_c.$$

Hence extinction can happen only if $e^{1/5} y_0/k < 5.0893$, so $y_c = 4.1667\,k$.

(d) Evidently, y_c is a linear function of the parameter k.

13.(a) The solution of the governing equation satisfies $u^3 = u_0^3/(\,3\,\alpha\,u_0^3\,t + 1\,)$. With the given data, it follows that $u(t) = 2000/\sqrt[3]{6\,t/125 + 1}$.

(b)




(c) Numerical evaluation results in $u(t) = 600$ for $t \approx 750.77$ s.

18.(a) The differential equation for the upward motion is $m dv/dt = -\mu v^2 - mg$, in which $\mu = 1/1325$. This equation is separable, with $m/(\mu\,v^2 + mg)\,dv = -dt$. Integrating both sides and invoking the initial condition, $v(t) = 44.133\,\tan(0.425 - 0.222\,t)$. Setting $v(t_1) = 0$, the ball reaches the maximum height at $t_1 = 1.916$ s. Integrating $v(t)$, the position is given by $x(t) = 198.75\,\ln\,[\cos(0.222\,t - 0.425)] + 48.57$. Therefore the maximum height is $x(t_1) = 48.56$ m.

(b) The differential equation for the downward motion is $m\,dv/dt = +\mu v^2 - mg$. This equation is also separable, with $m/(mg - \mu\,v^2)\,dv = -dt$. For convenience, set $t = 0$ at the top of the trajectory. The new initial condition becomes $v(0) = 0$. Integrating both sides and invoking the initial condition, we obtain $\ln((44.13 - v)/(44.13 + v)) = t/2.25$. Solving for the velocity, $v(t) = 44.13(1 - e^{t/2.25})/(1 + e^{t/2.25})$. Integrating $v(t)$, we obtain $x(t) = 99.29\,\ln(e^{t/2.25}/(1 + e^{t/2.25})^2) + 186.2$. To estimate the duration of the downward motion, set $x(t_2) = 0$, resulting in $t_2 = 3.276$ s. Hence the total time that the ball spends in the air is $t_1 + t_2 = 5.192$ s.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

(c)



19.(a) Measure the positive direction of motion upward. The equation of motion is given by $m dv/dt = -k\,v - mg$. The initial value problem is $dv/dt = -kv/m - g$, with $v(0) = v_0$. The solution is $v(t) = -mg/k + (v_0 + mg/k)e^{-kt/m}$. Setting $v(t_m) = 0$, the maximum height is reached at time $t_m = (m/k)\ln[(mg + k\,v_0)/mg]$. Integrating the velocity, the position of the body is

$$x(t) = -mg\,t/k + \left[(\frac{m}{k})^2 g + \frac{m\,v_0}{k}\right](1 - e^{-kt/m}).$$

Hence the maximum height reached is

$$x_m = x(t_m) = \frac{m\,v_0}{k} - g(\frac{m}{k})^2 \ln\left[\frac{mg + k\,v_0}{mg}\right].$$

(b) Recall that for $\delta \ll 1$, $\ln(1+\delta) = \delta - \delta^2/2 + \delta^3/3 - \delta^4/4 + \ldots$.

(c) The dimensions of the quantities involved are $[k] = MT^{-1}$, $[v_0] = LT^{-1}$, $[m] = M$ and $[g] = LT^{-2}$. This implies that kv_0/mg is dimensionless.


23.(a) Both equations are linear and separable. Initial conditions: $v(0) = u\cos A$ and $w(0) = u\sin A$. We obtain the solutions $v(t) = (u\cos A)e^{-rt}$ and $w(t) = -g/r + (u\sin A + g/r)e^{-rt}$.

(b) Integrating the solutions in part (a), and invoking the initial conditions, the coordinates are $x(t) = u\cos A(1 - e^{-rt})/r$ and

$$y(t) = -\frac{gt}{r} + \frac{g + ur\,\sin\,A + hr^2}{r^2} - (\frac{u}{r}\sin\,A + \frac{g}{r^2})e^{-rt}.$$

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

(c)



(d) Let T be the time that it takes the ball to go 350 ft horizontally. Then from above, $e^{-T/5} = (u\,\cos\,A - 70)/u\,\cos\,A$. At the same time, the height of the ball is given by

$$y(T) = -160T + 803 + 5u\sin A - \frac{(800 + 5u\sin A)(u\cos A - 70)}{u\cos A}.$$

Hence A and u must satisfy the equality

$$800\ln\left[\frac{u\,\cos\,A - 70}{u\,\cos\,A}\right] + 803 + 5u\sin\,A - \frac{(800 + 5u\sin A)(u\cos A - 70)}{u\cos A} = 10$$

for the ball to touch the top of the wall. To find the optimal values for u and A, consider u as a function of A and use implicit differentiation in the above equation to find that

$$\frac{du}{dA} = -\frac{u(u^2\cos A - 70u - 11200\sin A)}{11200\cos A}.$$

Solving this equation simultaneously with the above equation yields optimal values for u and A: $u \approx 145.3\,\text{ft/s}$, $A \approx 0.644\,\text{rad}$.

24.(a) Solving equation (i), $y'(x) = \left[(k^2 - y)/y\right]^{1/2}$. The positive answer is chosen, since y is an increasing function of x.

(b) Let $y = k^2\sin^2 t$. Then $dy = 2k^2\sin t\cos t dt$. Substituting into the equation in part (a), we find that

$$\frac{2k^2\sin t\cos t dt}{dx} = \frac{\cos t}{\sin t}.$$

Hence $2k^2\sin^2 t dt = dx$.

(c) Setting $\theta = 2t$, we further obtain $k^2\sin^2(\theta/2)\,d\theta = dx$. Integrating both sides of the equation and noting that $t = \theta = 0$ corresponds to the origin, we obtain the solutions $x(\theta) = k^2(\theta - \sin\theta)/2$ and (from part (b)) $y(\theta) = k^2(1 - \cos\theta)/2$.

(d) Note that $y/x = (1 - \cos\,\theta)/(\theta - \sin\,\theta)$. Setting $x = 1$, $y = 2$, the solution of the equation $(1 - \cos\,\theta)/(\theta - \sin\,\theta) = 2$ is $\theta \approx 1.401$. Substitution into either of the expressions yields $k \approx 2.193$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

2.4

2. The function $\tan t$ is discontinuous at odd multiples of $\pi/2$. Since $\pi/2 < \pi < 3\pi/2$, the initial value problem has a unique solution on the interval $(\pi/2, 3\pi/2)$.

4. The function $\ln t$ is defined and continuous on the interval $(0, \infty)$. At $t = 1$, $\ln t = 0$, so the normal form of the differential equation has a singularity there. Also, $\cot t$ is not defined at integer multiples of π, so the initial value problem will have a solution on the interval $(1, \pi)$.

6. The function $f(t, y)$ is discontinuous along the coordinate axes, and on the hyperbola $t^2 - y^2 = 1$. Furthermore,

$$\frac{\partial f}{\partial y} = \frac{\pm 1}{y(1 - t^2 + y^2)} - 2\frac{y \ln |ty|}{(1 - t^2 + y^2)^2}$$

has the same points of discontinuity.

7. $f(t, y)$ is continuous everywhere on the plane. The partial derivative $\partial f/\partial y$ is also continuous everywhere.


10. The equation is separable, with $dy/y^2 = 2t\,dt$. Integrating both sides, the solution is given by $y(t) = y_0/(1 - y_0 t^2)$. For $y_0 > 0$, solutions exist as long as $t^2 < 1/y_0$. For $y_0 \le 0$, solutions are defined for all t.

11. The equation is separable, with $dy/y^3 = -\,dt$. Integrating both sides and invoking the initial condition, $y(t) = y_0/\sqrt{2y_0^2 t + 1}$. Solutions exist as long as $2y_0^2 t + 1 > 0$, that is, $2y_0^2 t > -1$. If $y_0 \neq 0$, solutions exist for $t > -1/2y_0^2$. If $y_0 = 0$, then the solution $y(t) = 0$ exists for all t.

12. The function $f(t, y)$ is discontinuous along the straight lines $t = -1$ and $y = 0$. The partial derivative $\partial f/\partial y$ is discontinuous along the same lines. The equation is separable, with $y\,dy = t^2\,dt/(1 + t^3)$. Integrating and invoking the initial condition, the solution is $y(t) = \left[(2/3)\ln\left|1 + t^3\right| + y_0^2\right]^{1/2}$. Solutions exist as long as $(2/3)\ln\left|1 + t^3\right| + y_0^2 \ge 0$, that is, $y_0^2 \ge -(2/3)\ln\left|1 + t^3\right|$. For all y_0 (it can be verified that $y_0 = 0$ yields a valid solution, even though Theorem 2.4.2 does not guarantee one), solutions exist as long as $\left|1 + t^3\right| \ge e^{-3y_0^2/2}$. From above, we must have $t > -1$. Hence the inequality may be written as $t^3 \ge e^{-3y_0^2/2} - 1$. It follows that the solutions are valid for $(e^{-3y_0^2/2} - 1)^{1/3} < t < \infty$.


Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

14.




Based on the direction field, and the differential equation, for $y_0 < 0$, the slopes eventually become negative, and hence solutions tend to $-\infty$. For $y_0 > 0$, solutions increase without bound if $t_0 < 0$. Otherwise, the slopes eventually become negative, and solutions tend to zero. Furthermore, $y_0 = 0$ is an equilibrium solution. Note that slopes are zero along the curves $y = 0$ and $ty = 3$.

15.



For initial conditions (t_0, y_0) satisfying $ty < 3$, the respective solutions all tend to zero. For $y_0 \leq 9$, the solutions tend to 0; for $y_0 > 9$, the solutions tend to ∞. Also, $y_0 = 0$ is an equilibrium solution.

16.



Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

Solutions with $t_0 < 0$ all tend to $-\infty$. Solutions with initial conditions (t_0, y_0) to the right of the parabola $t = 1 + y^2$ asymptotically approach the parabola as $t \to \infty$. Integral curves with initial conditions above the parabola (and $y_0 > 0$) also approach the curve. The slopes for solutions with initial conditions below the parabola (and $y_0 < 0$) are all negative. These solutions tend to $-\infty$.

17.(a) No. There is no value of $t_0 \geq 0$ for which $(2/3)(t - t_0)^{2/3}$ satisfies the condition $y(1) = 1$.

(b) Yes. Let $t_0 = 1/2$ in Eq.(19).

(c) For $t_0 > 0$, $|y(2)| \leq (4/3)^{3/2} \approx 1.54$.

20. The assumption is $\phi'(t) + p(t)\phi(t) = 0$. But then $c\phi'(t) + p(t)c\phi(t) = 0$ as well.

22.(a) Recalling Eq.(33) in Section 2.1,

$$y = \frac{1}{\mu(t)} \int_{t_0}^{t} \mu(s)g(s)\,ds + \frac{c}{\mu(t)}.$$

It is evident that $y_1(t) = 1/\mu(t)$ and $y_2(t) = (1/\mu(t)) \int_{t_0}^{t} \mu(s)g(s)\,ds$.

(b) By definition, $1/\mu(t) = e^{-\int p(t)dt}$. Hence $y_1' = -p(t)/\mu(t) = -p(t)y_1$. That is, $y_1' + p(t)y_1 = 0$.

(c) $y_2' = (-p(t)/\mu(t)) \int_0^t \mu(s)g(s)\,ds + \mu(t)g(t)/\mu(t) = -p(t)y_2 + g(t)$. This implies that $y_2' + p(t)y_2 = g(t)$.

25. Since $n = 3$, set $v = y^{-2}$. It follows that $v' = -2y^{-3}y'$ and $y' = -(y^3/2)v'$. Substitution into the differential equation yields $-(y^3/2)v' - \varepsilon y = -\sigma y^3$, which further results in $v' + 2\varepsilon v = 2\sigma$. The latter differential equation is linear, and can be written as $(ve^{2\varepsilon t})' = 2\sigma e^{2\varepsilon t}$. The solution is given by $v(t) = \sigma/\varepsilon + ce^{-2\varepsilon t}$. Converting back to the original dependent variable, $y = \pm v^{-1/2} = \pm(\sigma/\varepsilon + ce^{-2\varepsilon t})^{-1/2}$.

27. The solution of the initial value problem $y_1' + 2y_1 = 0$, $y_1(0) = 1$ is $y_1(t) = e^{-2t}$. Therefore $y(1^-) = y_1(1) = e^{-2}$. On the interval $(1, \infty)$, the differential equation is $y_2' + y_2 = 0$, with $y_2(t) = ce^{-t}$. Therefore $y(1^+) = y_2(1) = ce^{-1}$. Equating the limits $y(1^-) = y(1^+)$, we require that $c = e^{-1}$. Hence the global solution of the initial value problem is

$$y(t) = \begin{cases} e^{-2t}, & 0 \leq t \leq 1 \\ e^{-1-t}, & t > 1 \end{cases}.$$


Note the discontinuity of the derivative


$$y'(t) = \begin{cases} -2e^{-2t}, & 0 < t < 1 \\ -e^{-1-t}, & t > 1 \end{cases}.$$

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

2.5


1.






The equilibrium points are $y^* = -a/b$ and $y^* = 0$, and $y' > 0$ when $y > 0$ or $y < -a/b$, and $y' < 0$ when $-a/b < y < 0$, therefore the equilibrium solution $y = -a/b$ is asymptotically stable and the equilibrium solution $y = 0$ is unstable.

3.





Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual



The only equilibrium point is $y^* = 0$, and $y' > 0$ when $y > 0$, $y' < 0$ when $y < 0$, hence the equilibrium solution $y = 0$ is unstable.

7.







The equilibrium points are $y^* = 0, \pm 1$, and $y' > 0$ for $y < -1$ or $0 < y < 1$ and $y' < 0$ for $-1 < y < 0$ or $y > 1$. The equilibrium solution $y = 0$ is unstable, and the remaining two are asymptotically stable.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual


8.






The equilibrium points are $y^* = 0, \pm 2$, and $y' < 0$ when $y < -2$ or $y > 2$, and $y' > 0$ for $-2 < y < 0$ or $0 < y < 2$. The equilibrium solutions $y = -2$ and $y = 2$ are unstable and asymptotically stable, respectively. The equilibrium solution $y = 0$ is semistable.

9.



Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual



The equilibrium points are $y^* = 0, 1$. $y' > 0$ for all y except $y = 0$ and $y = 1$. Both equilibrium solutions are semistable.

10. Eq.(10) is $y/(1-(y/K)) = Ce^{rt}$. Clearing the denominator gives $y = Ce^{rt}(1-(y/K)) = Ce^{rt} - Ce^{rt}(y/K)$. Thus $y + Ce^{rt}(y/K) = Ce^{rt}$, or $(1+(C/K)e^{rt})y = Ce^{rt}$. This last equation may easily be solved for y to give $y = Ce^{rt}/(1+(C/K)e^{rt})$. Applying the initial condition $y(0) = y_0$ gives $y_0 = C/(1+(C/K))$, which may be solved for C to give $C = y_0/(1-(y_0/K)) = Ky_0/(K-y_0)$. Using this last value of C in the solution for y gives $y(x) = ((Ky_0e^{rt}/(K-y_0))/(1+(y_0e^{rt}/(K-y_0))$, which may be simplified to yield Eq.(11).

11. To solve Eq.(12) for t, multiply each side of the equation by $(y_0/K)+[1-(y_0/K)]e^{-rt}$ to obtain $y((y_0/K)+[1-(y_0/K)]e^{-rt}) = y_0$, or $y_0/y = (y_0/K)+[1-(y_0/K)]e^{-rt}$. Multiplying each side of this equation by K gives $(y_0K)/y = y_0 + (K-y_0)e^{-rt}$, which may be solved for e^{-rt} to find that

$$e^{-rt} = \frac{(y_0K)/y - y_0}{K-y_0} = \frac{(y_0/y)-(y_0/K)}{1-(y_0/K)} = \frac{(y_0/K)[1-(y/K)]}{(y/K)[1-(y_0/K)]}$$

as in the text. Taking logarithms and dividing by $-r$ gives

$$t = -\frac{1}{r}\ln\frac{(y_0/K)[1-(y/K)]}{(y/K)[1-(y_0/K)]}$$

as given in Eq.(13).

12. To locate the time at which the solution given in Eq.(15) reaches its vertical asymptote, determine when the denominator of the solution is zero. Solving $y_0 + (T-y_0)e^{rt} = 0$ for e^{rt} gives $e^{rt} = y_0/(y_0-T)$, where the fact that $y_0 > T$ ensures that $y_0/(y_0-T) > 0$. Thus the equation $e^{rt} = y_0/(y_0-T)$ has a solution, which is $t = (1/r)\ln(y_0/(y_0-T))$ as given in Eq.(16).

13. To find the inflection points of a solution y to the differential equation $y' = f(y)$, first compute $y'' = f'(y)(dy/dt) = f'(y)f(y)$. Thus possible inflection points occur at solutions to $f'(y) = 0$ or $f(y) = 0$. The function $f(y) = -r(1-(y/T))(1-(y/K))y = 0$ at $y = 0$, $y = T$, and $y = K$, but Figure 2.5.7 shows that y'' does not change sign at any of these points, so they are not inflection points. To consider the points at which $f'(y) = 0$, note that $f(y) = (-r/(KT))(y-T)(y-K)y$, so $f'(y) = (-r/(KT))((y-K)y + (y-T)y + (y-T)(y-K)) = (-r/KT)(3y^2 - 2(K+T) +$

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

KT). Setting $f'(y) = 0$ and using the quadratic formula gives that $f'(y) = 0$ when $y = (K + T \pm \sqrt{K^2 - KT + T^2})/3$ as given in Eq.(18).

21.(a) The equilibrium points are at $y^* = 0$ and $y^* = 1$. Since $f'(y) = \alpha - 2\alpha y$, the equilibrium solution $y = 0$ is unstable and the equilibrium solution $y = 1$ is asymptotically stable.

(b) The differential equation is separable, with $[y(1-y)]^{-1}\, dy = \alpha\, dt$. Integrating both sides and invoking the initial condition, the solution is

$$y(t) = \frac{y_0\, e^{\alpha t}}{1 - y_0 + y_0\, e^{\alpha t}} = \frac{y_0}{y_0 + (1 - y_0)e^{-\alpha t}}.$$

It is evident that (independent of y_0) $\lim_{t \to -\infty} y(t) = 0$ and $\lim_{t \to \infty} y(t) = 1$.


22.(a) $y(t) = y_0\, e^{-\beta t}$.

(b) From part (a), $dx/dt = -\alpha x y_0 e^{-\beta t}$. Separating variables, $dx/x = -\alpha y_0 e^{-\beta t} dt$. Integrating both sides, the solution is $x(t) = x_0\, e^{-\alpha\, y_0 (1 - e^{-\beta t})/\beta}$.


(c) As $t \to \infty$, $y(t) \to 0$ and $x(t) \to x_0\, e^{-\alpha\, y_0/\beta}$. Over a long period of time, the proportion of carriers vanishes. Therefore the proportion of the population that escapes the epidemic is the proportion of susceptibles left at that time, $x_0\, e^{-\alpha\, y_0/\beta}$.


25.(a) For $a < 0$, the only critical point is at $y = 0$, which is asymptotically stable. For $a = 0$, the only critical point is at $y = 0$, which is asymptotically stable. For $a > 0$, the three critical points are at $y = 0, \pm\sqrt{a}$. The critical point at $y = 0$ is unstable, whereas the other two are asymptotically stable.

(b) Below, we graph solutions in the case $a = -1$, $a = 0$ and $a = 1$ respectively.




Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual






(c)




26.(a) $f(y) = y(a - y)$; $f'(y) = a - 2y$. For $a < 0$, the critical points are at $y = a$ and $y = 0$. Observe that $f'(a) > 0$ and $f'(0) < 0$. Hence $y = a$ is unstable and $y = 0$ asymptotically stable. For $a = 0$, the only critical point is at $y = 0$, which is semistable since $f(y) = -y^2$ is concave down. For $a > 0$, the critical points are at $y = 0$ and $y = a$. Observe that $f'(0) > 0$ and $f'(a) < 0$. Hence $y = 0$ is unstable and $y = a$ asymptotically stable.

(b) Below, we graph solutions in the case $a = -1$, $a = 0$ and $a = 1$ respectively.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual



(c)



Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

2.6

1. $M(x,y) = 2x + 3$ and $N(x,y) = 2y - 2$. Since $M_y = N_x = 0$, the equation is exact. Integrating M with respect to x, while holding y constant, yields $\psi(x,y) = x^2 + 3x + h(y)$. Now $\psi_y = h'(y)$, and equating with N results in the possible function $h(y) = y^2 - 2y$. Hence $\psi(x,y) = x^2 + 3x + y^2 - 2y$, and the solution is defined implicitly as $x^2 + 3x + y^2 - 2y = c$.

2. $M(x,y) = 2x + 4y$ and $N(x,y) = 2x - 2y$. Note that $M_y \neq N_x$, and hence the differential equation is not exact.

5. Write the equation as $(ax - by)dx + (bx - cy)dy = 0$. Now $M(x,y) = ax - by$ and $N(x,y) = bx - cy$. Since $M_y \neq N_x$, the differential equation is not exact.

7. $M(x,y) = y/x + 6x$ and $N(x,y) = \ln x - 2$. Since $M_y = N_x = 1/x$, the given equation is exact. Integrating N with respect to y, while holding x constant, results in $\psi(x,y) = y \ln x - 2y + h(x)$. Differentiating with respect to x, $\psi_x = y/x + h'(x)$. Setting $\psi_x = M$, we find that $h'(x) = 6x$, and hence $h(x) = 3x^2$. Therefore the solution is defined implicitly as $3x^2 + y \ln x - 2y = c$.

9. $M(x,y) = 2x - y$ and $N(x,y) = 2y - x$. Since $M_y = N_x = -1$, the equation is exact. Integrating M with respect to x, while holding y constant, yields $\psi(x,y) = x^2 - xy + h(y)$. Now $\psi_y = -x + h'(y)$. Equating ψ_y with N results in $h'(y) = 2y$, and hence $h(y) = y^2$. Thus $\psi(x,y) = x^2 - xy + y^2$, and the solution is given implicitly as $x^2 - xy + y^2 = c$. Invoking the initial condition $y(1) = 3$, the specific solution is $x^2 - xy + y^2 = 7$. The explicit form of the solution is $y(x) = (x + \sqrt{28 - 3x^2})/2$. Hence the solution is valid as long as $3x^2 \leq 28$.

12. $M(x,y) = y\,e^{2xy} + x$ and $N(x,y) = bx\,e^{2xy}$. Note that $M_y = e^{2xy} + 2xy\,e^{2xy}$, and $N_x = b\,e^{2xy} + 2bxy\,e^{2xy}$. The given equation is exact, as long as $b = 1$. Integrating N with respect to y, while holding x constant, results in $\psi(x,y) = e^{2xy}/2 + h(x)$. Now differentiating with respect to x, $\psi_x = y\,e^{2xy} + h'(x)$. Setting $\psi_x = M$, we find that $h'(x) = x$, and hence $h(x) = x^2/2$. We conclude that $\psi(x,y) = e^{2xy}/2 + x^2/2$. Hence the solution is given implicitly as $e^{2xy} + x^2 = c$.

13. Note that ψ is of the form $\psi(x\,,y) = f(x) + g(y)$, since each of the integrands is a function of a single variable. It follows that $\psi_x = f'(x)$ and $\psi_y = g'(y)$. That is, $\psi_x = M(x\,,y_0)$ and $\psi_y = N(x_0\,,y)$. Furthermore,

$$\frac{\partial^2 \psi}{\partial x \partial y}(x_0\,,y_0\,) = \frac{\partial M}{\partial y}(x_0\,,y_0\,) \text{ and } \frac{\partial^2 \psi}{\partial y \partial x}(x_0\,,y_0\,) = \frac{\partial N}{\partial x}(x_0\,,y_0\,),$$

based on the hypothesis and the fact that the point (x_0, y_0) is arbitrary, $\psi_{xy} = \psi_{yx}$ and $M_y(x,y) = N_x(x,y)$.

14. Observe that $(M(x))_y = (N(y))_x = 0$.

21. The equation is not exact, since $N_x - M_y = 2y - 1$. However, $(N_x - M_y)/M =$

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

$(2y-1)/y$ is a function of y alone. Hence there exists $\mu = \mu(y)$, which is a solution of the differential equation $\mu' = (2-1/y)\mu$. The latter equation is separable, with $d\mu/\mu = 2 - 1/y$. One solution is $\mu(y) = e^{2y-\ln y} = e^{2y}/y$. Now rewrite the given ODE as $e^{2y}dx + (2x\,e^{2y} - 1/y)dy = 0$. This equation is exact, and it is easy to see that $\psi(x,y) = x\,e^{2y} - \ln|y|$. Therefore the solution of the given equation is defined implicitly by $x\,e^{2y} - \ln|y| = c$.

22. Multiplying both sides of the ODE by $\mu = [xy(2x+y)]^{-1}$, the given equation is equivalent to $\left[(3x+y)/(2x^2+xy)\right]dx + \left[(x+y)/(2xy+y^2)\right]dy = 0$. Rewrite the differential equation as

$$\left[\frac{2}{x} + \frac{2}{2x+y}\right]dx + \left[\frac{1}{y} + \frac{1}{2x+y}\right]dy = 0\,.$$

It is easy to see that $M_y = N_x$. Integrating M with respect to x, while keeping y constant, results in $\psi(x,y) = 2\ln|x| + \ln|2x+y| + h(y)$. Now taking the partial derivative with respect to y, $\psi_y = (2x+y)^{-1} + h'(y)$. Setting $\psi_y = N$, we find that $h'(y) = 1/y$, and hence $h(y) = \ln|y|$. Therefore $\psi(x,y) = 2\ln|x| + \ln|2x+y| + \ln|y|$, and the solution of the given equation is defined implicitly by $2x^3y + x^2y^2 = c$.

2.7

2. The Euler formula is given by $y_{n+1} = y_n + h(2y_n - 1) = (1+2h)y_n - h$.


(a) 1.1, 1.22, 1.364, 1.5368

(b) 1.105, 1.23205, 1.38578, 1.57179

(c) 1.10775, 1.23873, 1.39793, 1.59144

(d) The differential equation is linear with solution $y(t) = (1+e^{2t})/2$. The values are 1.1107, 1.24591, 1.41106, 1.61277.


5.



All solutions seem to converge to $y = 25/9$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

7.



Solutions with initial conditions $|y(0)| > 2.5$ seem to diverge. On the other hand, solutions with initial conditions $|y(0)| < 2.5$ seem to converge to zero. Also, $y = 0$ is an equilibrium solution.

9. The Euler formula is $y_{n+1} = y_n - 3h\sqrt{y_n} + 5h$. The initial value is $y_0 = 2$.

(a) 2.30800, 2.49006, 2.60023, 2.66773, 2.70939, 2.73521

(b) 2.30167, 2.48263, 2.59352, 2.66227, 2.70519, 2.73209

(c) 2.29864, 2.47903, 2.59024, 2.65958, 2.70310, 2.73053

(d) 2.29686, 2.47691, 2.58830, 2.65798, 2.70185, 2.72959

10. The Euler formula is $y_{n+1} = (1 + 3h)y_n - ht_n y_n^2$. The initial value is $(t_0, y_0) = (0, 0.5)$.


(a) 1.70308, 3.06605, 2.44030, 1.77204, 1.37348, 1.11925

(b) 1.79548, 3.06051, 2.43292, 1.77807, 1.37795, 1.12191

(c) 1.84579, 3.05769, 2.42905, 1.78074, 1.38017, 1.12328

(d) 1.87734, 3.05607, 2.42672, 1.78224, 1.38150, 1.12411

14.(a)



Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

(b) The iteration formula is $y_{n+1} = y_n + h\,y_n^2 - h\,t_n^2$. The critical value α_0 appears to be between 0.67 and 0.68. For $y_0 > \alpha_0$, the iterations diverge.

15.(a) The ODE is linear, with general solution $y(t) = t + ce^t$. Invoking the specified initial condition, $y(t_0) = y_0$, we have $y_0 = t_0 + ce^{t_0}$. Hence $c = (y_0 - t_0)e^{-t_0}$. Thus the solution is given by $\phi(t) = (y_0 - t_0)e^{t-t_0} + t$.

(b) The Euler formula is $y_{n+1} = (1+h)y_n + h - h\,t_n$. Now set $k = n+1$.

(c) We have $y_1 = (1+h)y_0 + h - ht_0 = (1+h)y_0 + (t_1 - t_0) - ht_0$. Rearranging the terms, $y_1 = (1+h)(y_0 - t_0) + t_1$. Now suppose that $y_k = (1+h)^k(y_0 - t_0) + t_k$, for some $k \geq 1$. Then $y_{k+1} = (1+h)y_k + h - ht_k$. Substituting for y_k, we find that

$$y_{k+1} = (1+h)^{k+1}(y_0 - t_0) + (1+h)t_k + h - ht_k = (1+h)^{k+1}(y_0 - t_0) + t_k + h.$$

Noting that $t_{k+1} = t_k + h$, the result is verified.

(d) Substituting $h = (t - t_0)/n$, with $t_n = t$, $y_n = (1 + (t - t_0)/n)^n(y_0 - t_0) + t$. Taking the limit of both sides, and using the fact that $\lim_{n\to\infty}(1 + a/n)^n = e^a$, pointwise convergence is proved.

16. The exact solution is $y(t) = e^t$. The Euler formula is $y_{n+1} = (1+h)y_n$. It is easy to see that $y_n = (1+h)^n y_0 = (1+h)^n$. Given $t > 0$, set $h = t/n$. Taking the limit, we find that $\lim_{n\to\infty} y_n = \lim_{n\to\infty}(1 + t/n)^n = e^t$.

2.8

2. Let $z = y - 3$ and $\tau = t + 1$. It follows that $dz/d\tau = (dz/dt)(dt/d\tau) = dz/dt$. Furthermore, $dz/dt = dy/dt = 1 - y^3$. Hence $dz/d\tau = 1 - (z+3)^3$. The new initial condition is $z(0) = 0$.

3.(a) The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t 2\,[\phi_n(s) + 1]\,ds\,.$$


Setting $\phi_0(t) = 0$, $\phi_1(t) = 2t$. Continuing, $\phi_2(t) = 2t^2 + 2t$, $\phi_3(t) = 4t^3/3 + 2t^2 + 2t$, $\phi_4(t) = 2t^4/3 + 4t^3/3 + 2t^2 + 2t$, ... Based upon these we conjecture that $\phi_n(t) = \sum_{k=1}^n 2^k t^k/k!$ and use mathematical induction to verify this form for $\phi_n(t)$. First, let $n = 1$, then $\phi_n(t) = 2t$, so it is certainly true for $n = 1$. Then, using Eq.(7) again we have

$$\phi_{n+1}(t) = \int_0^t 2\,[\phi_n(s) + 1]\,ds = \int_0^t 2\left[\sum_{k=1}^n \frac{2^k}{k!}s^k + 1\right] ds = \sum_{k=1}^{n+1} \frac{2^k}{k!}t^k,$$

and we have verified our conjecture.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual


(b)



(c) Recall from calculus that $e^{at} = 1 + \sum_{k=1}^{\infty} a^k t^k / k!$. Thus

$$\phi(t) = \sum_{k=1}^{\infty} \frac{2^k}{k\,!} t^k = e^{2t} - 1\,.$$

(d)



From the plot it appears that ϕ_4 is a good estimate for $|t| < 1/2$.


4.(a) The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t [-\phi_n(s)/2 + s]\, ds\,.$$

Setting $\phi_0(t) = 0$, $\phi_1(t) = t^2/2$. Continuing, $\phi_2(t) = t^2/2 - t^3/12$, $\phi_3(t) = t^2/2 - t^3/12 + t^4/96$, $\phi_4(t) = t^2/2 - t^3/12 + t^4/96 - t^5/960$, ... Based upon these we conjecture that $\phi_n(t) = \sum_{k=1}^{n} 4(-1/2)^{k+1} t^{k+1}/(k+1)!$ and use mathematical induction to verify this form for $\phi_n(t)$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual


(b)



(c) Recall from calculus that $e^{at} = 1 + \sum_{k=1}^{\infty} a^k t^k / k!$. Thus

$$\phi(t) = \sum_{k=1}^{\infty} 4\frac{(-1/2)^{k+1}}{k+1\,!}t^{k+1} = 4e^{-t/2} + 2t - 4\,.$$

(d)



From the plot it appears that ϕ_4 is a good estimate for $|t| < 2$.

6.(a) The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t \left[s^2\phi_n(s) - s\right] ds\,.$$


Set $\phi_0(t) = 0$. The iterates are given by $\phi_1(t) = -t^2/2$, $\phi_2(t) = -t^2/2 - t^5/10$, $\phi_3(t) = -t^2/2 - t^5/10 - t^8/80$, $\phi_4(t) = -t^2/2 - t^5/10 - t^8/80 - t^{11}/880, \ldots$. Upon inspection, it becomes apparent that

$$\phi_n(t) = -t^2\left[\frac{1}{2} + \frac{t^3}{2\cdot 5} + \frac{t^6}{2\cdot 5\cdot 8} + \ldots + \frac{(t^3)^{n-1}}{2\cdot 5\cdot 8\ldots[2+3(n-1)]}\right] =$$

$$= -t^2\sum_{k=1}^{n}\frac{(t^3)^{k-1}}{2\cdot 5\cdot 8\ldots[2+3(k-1)]}.$$

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

(b)



(c) Using the identity $\phi_n(t) = \phi_1(t) + [\phi_2(t) - \phi_1(t)] + [\phi_3(t) - \phi_2(t)] + \ldots + [\phi_n(t) - \phi_{n-1}(t)]$, consider the series $\phi_1(t) + \sum_{k=1}^{\infty}[\phi_{k+1}(t) - \phi_k(t)]$. Fix any t value now. We use the Ratio Test to prove the convergence of this series:

$$\left|\frac{\phi_{k+1}(t) - \phi_k(t)}{\phi_k(t) - \phi_{k-1}(t)}\right| = \left|\frac{\frac{(-t^2)(t^3)^k}{2\cdot 5\cdots(2+3k)}}{\frac{(-t^2)(t^3)^{k-1}}{2\cdot 5\cdots(2+3(k-1))}}\right| = \frac{|t|^3}{2+3k}.$$


The limit of this quantity is 0 for any fixed t as $k \to \infty$, and we obtain that $\phi_n(t)$ is convergent for any t.

7.(a) The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t \left[s^2 + \phi_n^2(s)\right] ds\,.$$

Set $\phi_0(t) = 0$. The first three iterates are given by $\phi_1(t) = t^3/3$, $\phi_2(t) = t^3/3 + t^7/63$, $\phi_3(t) = t^3/3 + t^7/63 + 2t^{11}/2079 + t^{15}/59535$.

(b)



The iterates appear to be converging.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

10.(a) The approximating functions are defined recursively by


$$\phi_{n+1}(t) = \int_0^t \left[\frac{3s^2+4s+2}{2(\phi_n(s)-1)}\right] ds\,.$$

Note that $1/(2y-2) = -(1/2)\sum_{k=0}^{6} y^k + O(y^7)$. For computational purposes, use the geometric series sum to replace the above iteration formula by

$$\phi_{n+1}(t) = -\frac{1}{2}\int_0^t \left[(3s^2+4s+2)\sum_{k=0}^{6}\phi_n^k(s)\right] ds\,.$$

Set $\phi_0(t) = 0$. The first four approximations are given by $\phi_1(t) = -t - t^2 - t^3/2$, $\phi_2(t) = -t - t^2/2 + t^3/6 + t^4/4 - t^5/5 - t^6/24 + \ldots$, $\phi_3(t) = -t - t^2/2 + t^4/12 - 3t^5/20 + 4t^6/45 + \ldots$, $\phi_4(t) = -t - t^2/2 + t^4/8 - 7t^5/60 + t^6/15 + \ldots$

(b)



The approximations appear to be converging to the exact solution, which can be found by separating the variables: $\phi(t) = 1 - \sqrt{1+2t+2t^2+t^3}$.

12.(a) $\phi_n(0) = 0$, for every $n \geq 1$. Let $a \in (0\,,1]$. Then $\phi_n(a) = 2na\,e^{-na^2} = 2na/e^{na^2}$. Using l'Hospital's rule, $\lim_{z\to\infty} 2az/e^{az^2} = \lim_{z\to\infty} 1/ze^{az^2} = 0$. Hence $\lim_{n\to\infty}\phi_n(a) = 0\,.$

(b) $\int_0^1 2nx\,e^{-nx^2}dx = -e^{-nx^2}\big|_0^1 = 1 - e^{-n}$. Therefore,

$$\lim_{n\to\infty}\int_0^1 \phi_n(x)dx \neq \int_0^1 \lim_{n\to\infty}\phi_n(x)dx\,.$$

13.(a) Recall that Eq.(9) states that $\phi(t) = \int_0^t 2s[1+\phi(s)]\,ds$. Since $\phi(t) = \sum_{k=1}^{\infty} t^{2k}/k!$,

$$2s[1+\phi(s)] = 2s\sum_{k=0}^{\infty}\frac{s^{2k}}{k!} = 2\sum_{k=0}^{\infty}\frac{s^{2k+1}}{k!}$$

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

Integrating term-by-term,

$$\int_0^t 2s[1+\phi(s)]\,ds = \int_0^t 2\sum_{k=0}^{\infty}\frac{s^{2k+1}}{k!}\,ds = 2\sum_{k=0}^{\infty}\frac{1}{k!}\int_0^t s^{2k+1}\,ds$$
$$= 2\sum_{k=0}^{\infty}\frac{1}{k!}\frac{t^{2k+2}}{2k+2} = \sum_{k=0}^{\infty}\frac{t^{2(k+1)}}{(k+1)!} = \sum_{k=1}^{\infty}\frac{t^{2k}}{k!} = \phi(t)$$

and $\phi(t)$ is a solution of Eq.(9).

(b) Recall that the initial value problem in Eq.(8) is $y' = 2t(1+y), y(0) = 0$. Letting $y = \phi(t) = \sum_{k=1}^{\infty} t^{2k}/k!$,

$$y' = \phi'(t) = \sum_{k=1}^{\infty}\frac{2kt^{2k-1}}{k!} = 2\sum_{k=1}^{\infty}\frac{t^{2k-1}}{(k-1)!} = 2t\sum_{k=1}^{\infty}\frac{t^{2k-2}}{(k-1)!}$$
$$= 2t\sum_{k=0}^{\infty}\frac{t^{2k}}{k!} = 2t\left[1+\sum_{k=1}^{\infty}\frac{t^{2k}}{k!}\right] = 2t[1+\phi(t)]$$

and $y(0) = \phi(0) = \sum_{k=1}^{\infty} 0 = 0$, so $y = \phi(t)$ satisfies the initial value problem $y' = 2t(1+y), y(0) = 0$.

(c) Since $e^t = \sum_{k=0}^{\infty}\frac{t^k}{k!}$,

$$\phi(t) = \sum_{k=1}^{\infty}\frac{t^{2k}}{k!} = -1 + \sum_{k=0}^{\infty}\frac{t^{2k}}{k!} = -1 + \sum_{k=0}^{\infty}\frac{(t^2)^k}{k!} = -1 + e^{t^2}$$

(d) Separating the variables gives $(1/(1+y))\,dy = 2t\,dt$, and integration yields $\ln|1+y| = t^2 + c$. Applying the initial condition, $\ln 1 = 0 + c$, so $c = 0$ and $\ln|1+y| = t^2$. Solving for y first gives $|1+y| = e^{t^2}$, so $1+y = \pm e^{t^2}$. In order for $y(0) = 0$ to be true, we choose $1+y = e^{t^2}$, and thus $y = -1 + e^{t^2}$.

(e) Consider the first-order linear equation $y' - 2ty = 0$. The integrating factor will be $\mu(t) = e^{\int -2t\,dt} = e^{-t^2}$. Since multiplying the differential equation by $\mu(t)$ yields

$$\frac{d}{dt}\left(e^{-t^2}y\right) = 2te^{-t^2}$$

we have $e^{-t^2}y = \int 2te^{-t^2}\,dt = -e^{-t^2} + c$. The initial condition $y(0) = 0$ may now be applied to show that $c = 1$, and $y = e^{t^2}(-e^{-t^2}+1) = -1 + e^{t^2}$.

14. Let t be fixed, such that $(t, y_1), (t, y_2) \in D$. Without loss of generality, assume that $y_1 < y_2$. Since f is differentiable with respect to y, the mean value theorem asserts that there exists $\xi \in (y_1, y_2)$ such that $f(t, y_1) - f(t, y_2) = f_y(t, \xi)(y_1 - y_2)$. This means that $|f(t, y_1) - f(t, y_2)| = |f_y(t, \xi)|\,|y_1 - y_2|$. Since, by assumption, $\partial f/\partial y$ is continuous in D, f_y attains a maximum K on any closed and bounded subset of D. Hence $|f(t, y_1) - f(t, y_2)| \le K\,|y_1 - y_2|$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

15. For a sufficiently small interval of t, $\phi_{n-1}(t)$, $\phi_n(t) \in D$. Since f satisfies a Lipschitz condition, $|f(t,\phi_n(t)) - f(t,\phi_{n-1}(t))| \le K\,|\phi_n(t) - \phi_{n-1}(t)|$. Here $K = \max|f_y|$.

16.(a) $\phi_1(t) = \int_0^t f(s\,,0)ds$. Hence $|\phi_1(t)| \le \int_0^{|t|} |f(s\,,0)|\,ds \le \int_0^{|t|} M ds = M\,|t|$, in which M is the maximum value of $|f(t\,,y)|$ on D.

(b) By definition, $\phi_2(t) - \phi_1(t) = \int_0^t [f(s\,,\phi_1(s)) - f(s\,,0)]\,ds$. Taking the absolute value of both sides, $|\phi_2(t) - \phi_1(t)| \le \int_0^{|t|} |[f(s\,,\phi_1(s)) - f(s\,,0)]|\,ds$. Based on the results in Problems 14 and 15,

$$|\phi_2(t) - \phi_1(t)| \le \int_0^{|t|} K\,|\phi_1(s) - 0|\,ds \le KM \int_0^{|t|} |s|\,ds\,.$$

Evaluating the last integral, we obtain that $|\phi_2(t) - \phi_1(t)| \le MK\,|t|^2\,/2$.

(c) Suppose that

$$|\phi_i(t) - \phi_{i-1}(t)| \le \frac{MK^{i-1}\,|t|^i}{i!}$$

for some $i \ge 1$. By definition,

$$\phi_{i+1}(t) - \phi_i(t) = \int_0^t [f(s,\phi_i(s)) - f(s,\phi_{i-1}(s))]\,ds\,.$$

It follows that

$$|\phi_{i+1}(t) - \phi_i(t)| \le \int_0^{|t|} |f(s,\phi_i(s)) - f(s,\phi_{i-1}(s))|\,ds$$

$$\le \int_0^{|t|} K\,|\phi_i(s) - \phi_{i-1}(s)|\,ds \le \int_0^{|t|} K\frac{MK^{\,i-1}\,|s|^i}{i!}ds =$$

$$= \frac{MK^{\,i}\,|t|^{i+1}}{(i+1)!} \le \frac{MK^{\,i}h^{i+1}}{(i+1)!}\,.$$

Hence, by mathematical induction, the assertion is true.

17.(a) Use the triangle inequality, $|a+b| \le |a| + |b|$.

(b) For $|t| \le h$, $|\phi_1(t)| \le Mh$, and $|\phi_n(t) - \phi_{n-1}(t)| \le MK^{\,n-1}h^n/(n\,!)$. Hence

$$|\phi_n(t)| \le M\sum_{i=1}^{n} \frac{K^{\,i-1}h^i}{i\,!} = \frac{M}{K}\sum_{i=1}^{n} \frac{(Kh)^i}{i\,!}\,.$$

(c) The sequence of partial sums in (b) converges to $M(e^{Kh} - 1)/K$. By the comparison test, the sums in (a) also converge. Since individual terms of a convergent series must tend to zero, $|\phi_n(t) - \phi_{n-1}(t)| \to 0$, and it follows that the sequence $|\phi_n(t)|$ is convergent.

18.(a) Let $\phi(t) = \int_0^t f(s\,,\phi(s))ds$ and $\psi(t) = \int_0^t f(s\,,\psi(s))ds$. Then by linearity of the integral, $\phi(t) - \psi(t) = \int_0^t [f(s\,,\phi(s)) - f(s\,,\psi(s))]\,ds$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

(b) It follows that $|\phi(t) - \psi(t)| \le \int_0^t |f(s\,,\phi(s)) - f(s\,,\psi(s))|\,ds\,.$

(c) We know that f satisfies a Lipschitz condition, $|f(t\,,y_1) - f(t\,,y_2)| \le K\ |y_1 - y_2|$, based on $|\partial f/\partial y| \le K$ in D. Therefore,

$$|\phi(t) - \psi(t)| \le \int_0^t |f(s\,,\phi(s)) - f(s\,,\psi(s))|\,ds \le \int_0^t K\,|\phi(s) - \psi(s)|\,ds.$$

2.9

1. Writing the equation for each $n \ge 0\,,\ y_1 = -0.9\,y_0\,,\ y_2 = -0.9\,y_1 = (-0.9^2)y_0$, $y_3 = -0.9\,y_2 = (-0.9)^3 y_0$ and so on, it is apparent that $y_n = (-0.9)^n\,y_0\,.$ The terms constitute an alternating series, which converge to zero, regardless of $y_0\,.$

2. Write the equation for each $n \ge 0$, $y_1 = \sqrt{3}\,y_0$, $y_2 = \sqrt{4/2}\,y_1$, $y_3 = \sqrt{5/3}\,y_2$, ... Upon substitution, we find that $y_2 = \sqrt{(4\cdot 3)/2}\,y_1$, $y_3 = \sqrt{(5\cdot 4\cdot 3)/(3\cdot 2)}\,y_0$, ... It can be proved by mathematical induction, that

$$y_n = \frac{1}{\sqrt{2}}\sqrt{\frac{(n+2)!}{n!}}\ y_0 \ = \frac{1}{\sqrt{2}}\sqrt{(n+1)(n+2)}\ y_0\,.$$

This sequence is divergent, except for $y_0 = 0\,.$

3. Writing the equation for each $n \ge 0\,,\ y_1 = -y_0\,,\ y_2 = y_1\,,\ y_3 = -y_2\,,\ y_4 = y_3\,,$ and so on. It can be shown that

$$y_n = \begin{cases} y_0, & \text{for } n = 4k \text{ or } n = 4k-1 \\ -y_0, & \text{for } n = 4k-2 \text{ or } n = 4k-3 \end{cases}$$

The sequence is convergent only for $y_0 = 0\,.$

5. Let y_n be the balance at the end of the nth month. Then $y_{n+1} = (1 + r/12)y_n + 25$. We have $y_n = \rho^n[y_0 - 25/(1-\rho)] + 25/(1-\rho)$, in which $\rho = (1 + r/12)$. Here r is the annual interest rate, given as 8%. Thus $y_{36} = (1.0066)^{36}\,[1000 + 12\cdot 25/r] - 12\cdot 25/r = \$2,283.63.$

6. Let y_n be the balance due at the end of the nth month. The appropriate difference equation is $y_{n+1} = (1 + r/12)\,y_n - P\,.$ Here r is the annual interest rate and P is the monthly payment. The solution, in terms of the amount borrowed, is given by $y_n = \rho^n[y_0 + P/(1-\rho)] - P/(1-\rho)$, in which $\rho = (1 + r/12)$ and $y_0 = 8,000\,.$ To figure out the monthly payment P, we require that $y_{36} = 0$. That is, $\rho^{36}[y_0 + P/(1-\rho)] = P/(1-\rho)$. After the specified amounts are substituted, we find that $P = \$258.14.$

7. Let y_n be the balance due at the end of the nth month. The appropriate difference equation is $y_{n+1} = (1 + r/12)\,y_n - P$, in which $r = .09$ and P is the monthly payment. The initial value of the mortgage is $y_0 = \$100,000.$ Then the balance

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

due at the end of the n-th month is $y_n = \rho^n[y_0 + P/(1-\rho)] - P/(1-\rho)$, where $\rho = (1 + r/12)$. In terms of the specified values, $y_n = (1.0075)^n[10^5 - 12P/r] + 12P/r$. Setting $n = 30 \cdot 12 = 360$, and $y_{360} = 0$, we find that $P = \$804.62$. For the monthly payment corresponding to a 20 year mortgage, set $n = 240$ and $y_{240} = 0$ to find that $P = \$899.73$. The total amount paid during the term of the loan is $360 \times 804.62 = \$289,663.20$ for the 30-year loan and is $240 \times 899.73 = \$215,935.20$ for the 20-year loan.


8. Let y_n be the balance due at the end of the nth month, with y_0 the initial value of the mortgage. The appropriate difference equation is $y_{n+1} = (1 + r/12)\, y_n - P$, in which $r = 0.1$ and $P = \$1000$ is the maximum monthly payment. Given that the life of the mortgage is 20 years, we require that $y_{240} = 0$. The balance due at the end of the n-th month is $y_n = \rho^n[y_0 + P/(1-\rho)] - P/(1-\rho)$. In terms of the specified values for the parameters, the solution of $(1.00833)^{240}[y_0 - 12 \cdot 1000/0.1] = -12 \cdot 1000/0.1$ is $y_0 = \$103,624.62$.

15.(a) $\delta_2 = (\rho_2 - \rho_1)/(\rho_3 - \rho_2) = (3.449 - 3)/(3.544 - 3.449) = 4.7263$.

(b) diff$= (|\delta - \delta_2|/\delta) \cdot 100 = (|4.6692 - 4.7363|/4.6692) \cdot 100 \approx 1.22\%$.

(c) Assuming $(\rho_3 - \rho_2)/(\rho_4 - \rho_3) = \delta$, $\rho_4 \approx 3.5643$

(d) A period 16 solution appears near $\rho \approx 3.565$.



(e) Note that $(\rho_{n+1} - \rho_n) = \delta_n^{-1}(\rho_n - \rho_{n-1})$. With the assumption that $\delta_n = \delta$, we have $(\rho_{n+1} - \rho_n) = \delta^{-1}(\rho_n - \rho_{n-1})$, which is of the form $y_{n+1} = \alpha\, y_n$, $n \geq 3$. It follows that $(\rho_k - \rho_{k-1}) = \delta^{3-k}(\rho_3 - \rho_2)$ for $k \geq 4$. Then

$$\begin{aligned}\rho_n &= \rho_1 + (\rho_2 - \rho_1) + (\rho_3 - \rho_2) + (\rho_4 - \rho_3) + \ldots + (\rho_n - \rho_{n-1}) \\ &= \rho_1 + (\rho_2 - \rho_1) + (\rho_3 - \rho_2)\left[1 + \delta^{-1} + \delta^{-2} + \ldots + \delta^{3-n}\right] \\ &= \rho_1 + (\rho_2 - \rho_1) + (\rho_3 - \rho_2)\left[\frac{1 - \delta^{4-n}}{1 - \delta^{-1}}\right].\end{aligned}$$

Hence $\lim_{n\to\infty} \rho_n = \rho_2 + (\rho_3 - \rho_2)\left[\frac{\delta}{\delta - 1}\right]$. Substitution of the appropriate values yields

$$\lim_{n\to\infty} \rho_n = 3.5699$$

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

PROBLEMS

1. The equation is *linear*. It can be written in the form $y' + 2y/x = x^2$, and the integrating factor is $\mu(x) = e^{\int (2/x)\,dx} = e^{2\ln x} = x^2$. Multiplication by $\mu(x)$ yields $x^2y' + 2yx = (yx^2)' = x^4$. Integration with respect to x and division by x^2 gives that $y = x^3/5 + c/x^2$.

5. The equation is *exact*. Algebraic manipulations give the symmetric form of the equation, $(2xy + y^2 + 1)dx + (x^2 + 2xy)dy = 0$. We can check that $M_y = 2x + 2y = N_x$, so the equation is really exact. Integrating M with respect to x gives that $\psi(x, y) = x^2y + xy^2 + x + g(y)$, then $\psi_y = x^2 + 2xy + g'(y) = x^2 + 2xy$, so we get that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the solution is defined implicitly as $x^2y + xy^2 + x = c$.

6. The equation is *linear*. It can be written in the form $y' + (1 + (1/x))y = 1/x$ and the integrating factor is $\mu(x) = e^{\int 1+(1/x)\,dx} = e^{x+\ln x} = xe^x$. Multiplication by $\mu(x)$ yields $xe^xy' + (xe^x + e^x)y = (xe^xy)' = e^x$. Integration with respect to x and division by xe^x shows that the general solution of the equation is $y = 1/x + c/(xe^x)$. The initial condition implies that $0 = 1 + c/e$, which means that $c = -e$ and the solution is $y = 1/x - e/(xe^x) = x^{-1}(1 - e^{1-x})$.

7. The equation is *linear*. It can be written in the form $y' + 2y/x = \sin x/x^2$ and the integrating factor is $\mu(x) = e^{\int (2/x)\,dx} = e^{2\ln x} = x^2$. Multiplication by $\mu(x)$ gives $x^2y' + 2xy = (x^2y)' = \sin x$, and after integration with respect to x and division by x^2 we obtain the general solution $y = (c - \cos x)/x^2$. The initial condition implies that $c = 4 + \cos 2$ and the solution becomes $y = (4 + \cos 2 - \cos x)/x^2$.

10. The equation is *exact*. It is easy to check that $M_y = 1 = N_x$. Integrating M with respect to x gives that $\psi(x, y) = x^3/3 + xy + g(y)$, then $\psi_y = x + g'(y) = x + e^y$, which means that $g'(y) = e^y$, so we obtain that $g(y) = e^y$. Therefore the solution is defined implicitly as $x^3/3 + xy + e^y = c$.

11. The equation is *exact*. We can check that $M_y = 1 = N_x$. Integrating M with respect to x gives that $\psi(x, y) = x^2/2 + xy + g(y)$, then $\psi_y = x + g'(y) = x + 2y$, which means that $g'(y) = 2y$, so we obtain that $g(y) = y^2$. Therefore the general solution is defined implicitly as $x^2/2 + xy + y^2 = c$. The initial condition gives us $c = 17$, so the solution is $x^2 + 2xy + 2y^2 = 34$.

12. The equation is *separable*. Separation of variables leads us to the equation

$$\frac{dy}{y} = \frac{1 - e^x}{1 + e^x}dx.$$

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

Note that $1 + e^x - 2e^x = 1 - e^x$. We obtain that

$$\ln|y| = \int \frac{1-e^x}{1+e^x}dx = \int 1 - \frac{2e^x}{1+e^x}dx = x - 2\ln(1+e^x) + \tilde{c}.$$

This means that $y = ce^x(1+e^x)^{-2}$, which also can be written as $y = c/\cosh^2(x/2)$ after some algebraic manipulations.

13. The equation is *exact*. The symmetric form is $(-e^{-x}\cos y + e^{2y}\cos x)dx + (-e^{-x}\sin y + 2e^{2y}\sin x)dy = 0$. We can check that $M_y = e^{-x}\sin y + 2e^{2y}\cos x = N_x$. Integrating M with respect to x gives that $\psi(x,y) = e^{-x}\cos y + e^{2y}\sin x + g(y)$, then $\psi_y = -e^{-x}\sin y + 2e^{2y}\sin x + g'(y) = -e^{-x}\sin y + 2e^{2y}\sin x$, so we get that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the solution is defined implicitly as $e^{-x}\cos y + e^{2y}\sin x = c$.

14. The equation is *linear*. The integrating factor is $\mu(x) = e^{-\int 3\,dx} = e^{-3x}$, which turns the equation into $e^{-3x}y' - 3e^{-3x}y = (e^{-3x}y)' = e^{-x}$. We integrate with respect to x to obtain $e^{-3x}y = -e^{-x} + c$, and the solution is $y = ce^{3x} - e^{2x}$ after multiplication by e^{3x}.

15. The equation is *linear*. The integrating factor is $\mu(x) = e^{\int 2\,dx} = e^{2x}$, which gives us $e^{2x}y' + 2e^{2x}y = (e^{2x}y)' = e^{-x^2}$. The antiderivative of the function on the right hand side can not be expressed in a closed form using elementary functions, so we have to express the solution using integrals. Let us integrate both sides of this equation from 0 to x. We obtain that the left hand side turns into

$$\int_0^x (e^{2s}y(s))'ds = e^{2x}y(x) - e^0 y(0) = e^{2x}y - 3.$$

The right hand side gives us $\int_0^x e^{-s^2}\,ds$. So we found that

$$y = e^{-2x}\int_0^x e^{-s^2}\,ds + 3e^{-2x}.$$

16. The equation is *exact*. Algebraic manipulations give us the symmetric form $(y^3 + 2y - 3x^2)dx + (2x + 3xy^2)dy = 0$. We can check that $M_y = 3y^2 + 2 = N_x$. Integrating M with respect to x gives that $\psi(x,y) = xy^3 + 2xy - x^3 + g(y)$, then $\psi_y = 3xy^2 + 2x + g'(y) = 2x + 3xy^2$, which means that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the solution is $xy^3 + 2xy - x^3 = c$.

17. The equation is *separable*, because $y' = e^{x+y} = e^x e^y$. Separation of variables yields the equation $e^{-y}dy = e^x dx$, which turns into $-e^{-y} = e^x + c$ after integration and we obtain the implicitly defined solution $e^x + e^{-y} = c$.

19. The equation is *linear*. Division by t gives $y' + (1 + (1/t))y = e^{2t}/t$, so the integrating factor is $\mu(t) = e^{\int(1+(1/t))dt} = e^{t+\ln t} = te^t$. The equation turns into $te^t y' + (te^t + e^t)y = (te^t y)' = e^{3t}$. Integration therefore leads to $te^t y = e^{3t}/3 + c$ and the solution is $y = e^{2t}/(3t) + ce^{-t}/t$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

22. The equation is *homogeneous*. (See Section 2.2, Problem 25) We can see that

$$y' = \frac{x+y}{x-y} = \frac{1+(y/x)}{1-(y/x)}.$$

We substitute $u = y/x$, which means also that $y = ux$ and then $y' = u'x + u = (1+u)/(1-u)$, which implies that

$$u'x = \frac{1+u}{1-u} - u = \frac{1+u^2}{1-u},$$

a separable equation. Separating the variables yields

$$\frac{1-u}{1+u^2}du = \frac{dx}{x},$$

and then integration gives $\arctan u - \ln(1+u^2)/2 = \ln|x| + c$. Substituting $u = y/x$ back into this expression and using that

$$-\ln(1+(y/x)^2)/2 - \ln|x| = -\ln(|x|\sqrt{1+(y/x)^2}) = -\ln(\sqrt{x^2+y^2})$$

we obtain that the solution is $\arctan(y/x) - \ln(\sqrt{x^2+y^2}) = c$.

23. The equation is *homogeneous*. (See Section 2.2, Problem 25) Algebraic manipulations show that it can be written in the form

$$y' = \frac{3y^2+2xy}{2xy+x^2} = \frac{3(y/x)^2+2(y/x)}{2(y/x)+1}.$$

Substituting $u = y/x$ gives that $y = ux$ and then

$$y' = u'x + u = \frac{3u^2+2u}{2u+1},$$

which implies that

$$u'x = \frac{3u^2+2u}{2u+1} - u = \frac{u^2+u}{2u+1},$$

a separable equation. We obtain that $(2u+1)du/(u^2+u) = dx/x$, which in turn means that $\ln(u^2+u) = \ln|x| + \tilde{c}$. Therefore, $u^2+u = cx$ and then substituting $u = y/x$ gives us the solution $(y^2/x^3) + (y/x^2) = c$.

25. Let y_1 be a solution, i.e. $y_1' = q_1 + q_2y_1 + q_3y_1^2$. Now let $y = y_1 + (1/v)$ also be a solution. Differentiating this expression with respect to t and using that y is also a solution we obtain $y' = y_1' - (1/v^2)v' = q_1 + q_2y + q_3y^2 = q_1 + q_2(y_1 + (1/v)) + q_3(y_1 + (1/v))^2$. Now using that y_1 was also a solution we get that $-(1/v^2)v' = q_2(1/v) + 2q_3(y_1/v) + q_3(1/v^2)$, which, after some simple algebraic manipulations turns into $v' = -(q_2 + 2q_3y_1)v - q_3$.

27.(a) The equation is $y' = (1-y)(x+by) = x + (b-x)y - by^2$. We set $y = 1 + (1/v)$ and differentiate: $y' = -v^{-2}v' = x + (b-x)(1+(1/v)) - b(1+(1/v))^2$, which, after simplification, turns into $v' = (b+x)v + b$.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

(b) When $x = at$, the equation is $v' - (b + at)v = b$, so the integrating factor is $\mu(t) = e^{-bt-at^2/2}$. This turns the equation into $(v\mu(t))' = b\mu(t)$, so $v\mu(t) = \int b\mu(t)dt$, and then $v = (b\int \mu(t)dt)/\mu(t)$.

28. Substitute $v = y'$, then $v' = y''$. The equation turns into $t^2v' + 2tv = (t^2v)' = 1$, which yields $t^2v = t + c_1$, so $y' = v = (1/t) + (c_1/t^2)$. Integrating this expression gives us the solution $y = \ln t - (c_1/t) + c_2$.

29. Set $v = y'$, then $v' = y''$. The equation with this substitution is $tv' + v = (tv)' = 1$, which gives $tv = t + c_1$, so $y' = v = 1 + (c_1/t)$. Integrating this expression yields the solution $y = t + c_1 \ln t + c_2$.

30. Set $v = y'$, so $v' = y''$. The equation is $v' + tv^2 = 0$, which is a separable equation. Separating the variables we obtain $dv/v^2 = -tdt$, so $-1/v = -t^2/2 + c$, and then $y' = v = 2/(t^2 + c_1)$. Now depending on the value of c_1, we have the following possibilities: when $c_1 = 0$, then $y = -2/t + c_2$, when $0 < c_1 = k^2$, then $y = (2/k)\arctan(t/k) + c_2$, and when $0 > c_1 = -k^2$ then

$$y = (1/k)\ln|(t - k)/(t + k)| + c_2.$$

We also divided by $v = y'$ when we separated the variables, and $v = 0$ (which is $y = c$) is also a solution.

31. Substitute $v = y'$ and $v' = y''$. The equation is $2t^2v' + v^3 = 2tv$. This is a *Bernoulli* equation (See Section 2.4, Problem 19), so the substitution $z = v^{-2}$ yields $z' = -2v^{-3}v'$, and the equation turns into $2t^2v'v^3 + 1 = 2t/v^2$, i.e. into $-2t^2z'/2 + 1 = 2tz$, which in turn simplifies to $t^2z' + 2tz = (t^2z)' = 1$. Integration yields $t^2z = t + c$, which means that $z = (1/t) + (c/t^2)$. Now $y' = v = \pm\sqrt{1/z} = \pm t/\sqrt{t + c_1}$ and another integration gives

$$y = \pm\frac{2}{3}(t - 2c_1)\sqrt{t + c_1} + c_2.$$

The substitution also loses the solution $v = 0$, i.e. $y = c$.

33. Set $y' = v(y)$. Then $y'' = v'(y)(dy/dt) = v'(y)v(y)$. We obtain the equation $v'v + y = 0$, where the differentiation is with respect to y. This is a separable equation which simplifies to $vdv = -ydy$. We obtain that $v^2/2 = -y^2/2 + c$, so $y' = v(y) = \pm\sqrt{c - y^2}$. We separate the variables again to get $dy/\sqrt{c - y^2} = \pm dt$, so $\arcsin(y/\sqrt{c}) = t + d$, which means that $y = \sqrt{c}\sin(\pm t + d) = c_1\sin(t + c_2)$.

34. Set $y' = v(y)$. Then $y'' = v'(y)(dy/dt) = v'(y)v(y)$. We obtain the equation $yv'v - v^3 = 0$, where the differentiation is with respect to y. This separable equation gives us $dv/v^2 = dy/y$, which means that $-1/v = \ln|y| + c$, and then $y' = v = 1/(c - \ln|y|)$. We separate variables again to obtain $(c - \ln|y|)dy = dt$, and then integration yields the implicitly defined solution $cy - (y\ln|y| - y) = t + d$. Also, $y = c$ is a solution which we lost when we divided by $v = 0$.

37. Set $v = y'$, then $v' = y''$. The equation with this substitution turns into the equation $(1 + t^2)v' + 2tv = ((1 + t^2)v)' = -3t^{-2}$. Integrating this we get that

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

$(1+t^2)v = 3t^{-1} + c$, and $c = -5$ from the initial conditions. This means that $y' = v = 3/(t(1+t^2)) - 5/(1+t^2)$. The partial fraction decomposition of the first expression shows that $y' = 3/t - 3t/(1+t^2) - 5/(1+t^2)$ and then another integration here gives us that $y = 3\ln t - (3/2)\ln(1+t^2) - 5\arctan t + d$. The initial conditions identify $d = 2 + (3/2)\ln 2 + 5\pi/4$, and we obtained the solution.

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual

Elementary Differential Equations and Boundary Value Problems 11th edition Boyce Solutions Manual