Foundations of College Chemistry 16th edition Hein Solutions Manual

SKU: 9781119768159-SOLUTIONS

CHAPTER 2

STANDARDS FOR MEASUREMENT

SOLUTIONS TO REVIEW QUESTIONS

1. The exponent will be positive for a large number and negative for a small number.
2. The exponent will decrease.
3. The last digit in a measurement is uncertain because if the quantity were to be measured multiple times, the last digit would vary.
4. It must be written in scientific notation as 6.420×10^5 g.
5. Zeroes are significant when they are between nonzero digits or at the end of a number that includes a decimal point.
6. Rule 1. When the first digit after those you want to retain is 4 or less, that digit and all others to its right are dropped. The last digit retained is not changed.

   Rule 2. When the first digit after those you want to retain is 5 or greater, that digit and all others to the right of it are dropped and the last digit retained is increased by one.
7. No, the number of significant digits in the calculated value may not be more than the number of significant figures in any of the measurements.
8. Yes, the number of significant digits depends on the precision of each of the individual measurements and the calculated value may have more or fewer significant figures than the original measurements as long as the precision is no greater than the measurement with the lowest precision. An example of a calculation with an increase in significant figures is in Example 2.9.
9. $100\text{ cm} = 1\text{ m}$

   $1{,}000{,}000{,}000\text{ nm} = 1\text{ m} \quad (1\text{ cm})\left(\frac{1\text{ m}}{100\text{ cm}}\right)\left(\frac{1{,}000{,}000{,}000\text{ nm}}{\text{m}}\right) = 10{,}000{,}000\text{ nm}$
10. 1000 mg = 1 g

    $1000\text{ g} = 1\text{ kg} \quad (1\text{ kg})\left(\frac{1000\text{ g}}{\text{kg}}\right)\left(\frac{1000\text{ mg}}{\text{g}}\right) = 1{,}000{,}000\text{ mg}$
11. Weight is a measure of how much attraction the earth's gravity has for an object (or person). In this case, the farther the astronaut is from the Earth the less gravitational force is pulling on him or her. Less gravitational attraction means the astronaut will weigh less. The mass of the astronaut is the amount of matter that makes up him or her. This does not change as the astronaut moves away from the Earth.
12. They are equivalent units.
13. $(3.5\text{ in.})\left(\frac{2.54\text{ cm}}{1\text{ in.}}\right) = 8.9\text{ cm}$
14. Heat is a form of energy, while temperature is a measure of the intensity of heat (how hot the system is).

Foundations of College Chemistry 16th edition Hein Solutions Manual

15. The number of degrees between the freezing and boiling point of water are

Fahrenheit	180°F
Celsius	100°C
Kelvin	100 K

16. The three materials would sort out according to their densities with the most dense (mercury) at the bottom and the least dense (glycerin) at the top. In the cylinder, the solid magnesium would sink in the glycerin and float on the liquid mercury.

17. Order of increasing density: ethyl alcohol, vegetable oil, salt, lead

18. The density of ice must be less than 0.91 g/mL and greater than 0.789 g/mL.

19. The density of water is 1.0 g/mL at approximately 4°C. However, when water changes from a liquid to a solid at 0°C there is actually an increase in volume. The density of ice at 0°C is 0.917 g/mL. Therefore, ice floats in water because solid water is less dense than liquid water.

20. If you collect a container of oxygen gas, you should store it with the mouth up. Oxygen gas is denser than air.

21. density of gold = 19.3 g/mL density of silver = 10.5 g/mL

$$(25 \text{ g gold})\left(\frac{1 \text{ mL}}{19.3 \text{ g}}\right) = 1.3 \text{ mL}$$

$$(25 \text{ g silver})\left(\frac{1 \text{ mL}}{10.5 \text{ g}}\right) = 2.4 \text{ mL}$$

25 g of silver has the greater volume.

Foundations of College Chemistry 16th edition Hein Solutions Manual

SOLUTIONS TO EXERCISES

1. (a) kilogram = 1000 grams
   (b) centimetre = 1/100 of a meter (0.01 m)
   (c) microliter = 1/1,000,000 of a liter (0.000001 L)
   (d) millimetre = 1/1000 of a meter (0.001 m)
   (e) decilitre = 1/10 of a liter (0.1 L)

2. (a) 1000 meters = 1 kilometer
   (b) 0.1 gram = 1 decigram
   (c) 0.000001 liter = 1 microliter
   (d) 0.01 meter = 1 centimeter
   (e) 0.001 liter = 1 milliliter

3. (a) gram = g
   (b) microgram = μg
   (c) centimetre = cm
   (d) micrometer = μm
   (e) millilitre = mL
   (f) decilitre = dL

4. (a) milligram = mg
   (b) kilogram = kg
   (c) meter = m
   (d) nanometer = nm
   (e) angstrom= Å
   (f) microlitre = μL

5. | | | |
   |---|---|---|
   | (a) | 2050 | the first zero is significant; the last zero is not significant |
   | (b) | 9.00×10^2 | zeros are significant |
   | (c) | 0.0530 | the first two zeros are not significant; the last zero is significant |
   | (d) | 0.075 | zeros are not significant |
   | (e) | 300. | zeros are significant |
   | (f) | 285.00 | zeros are significant |

6. | | | |
   |---|---|---|
   | (a) | 0.005 | zeros are not significant |
   | (b) | 1500 | zeros are not significant |
   | (c) | 250. | zero is significant |
   | (d) | 10.000 | zeros are significant |
   | (e) | 6.070×10^4 | zeros are significant |
   | (f) | 0.2300 | the first zero is not significant; the last two zeros are significant |

Foundations of College Chemistry 16th edition Hein Solutions Manual

7. Significant figures

(a) 0.025	(2 sig. fig.)	(c) 0.0404	(3 sig. fig.)
(b) 22.4	(3 sig. fig.)	(d) 5.50×10^3	(3 sig. fig.)

8. Significant figures

(a) 40.0	(3 sig. fig.)	(c) 129,042	(6 sig. fig.)
(b) 0.081	(2 sig. fig.)	(d) 4.090×10^{-3}	(4 sig. fig.)

9. Round each of the following numbers to four significant figures:

(a) 47.70 (c) 100.2

(b) 1.026 (d) 16.50

10. Round each of the following numbers to four significant figures:

(a) 0.07794 (c) 5.005

(b) 21.63 (d) 74.29

11. Exponential notation

(a) 6.7×10^4 (c) 7.8×10^{-3}

(b) 6.54×10^{-2} (d) 4.11×10^5

12. Exponential notation

(a) 4.56×10^{-2} (c) 4.030×10^1

(b) 4.0822×10^3 (d) 1.2×10^7

13. (a)
$$\begin{array}{r} 12.62 \\ 1.5 \\ 0.25 \\ \hline 14.37 = 14.4 \end{array}$$

(b) $(2.25 \times 10^3)(4.80 \times 10^4) = 10.8 \times 10^7 = 1.08 \times 10^8$

(c) $\dfrac{(452)(6.2)}{14.3} = 195.97 = 2.0 \times 10^2$

(d) $(0.0394)\,(12.8) = 0.504$

(e) $\dfrac{0.4278}{59.6} = 0.00718 = 7.18 \times 10^{-3}$

(f) $10.4 + (3.75)(1.5 \times 10^4) = 5.6 \times 10^4$

Foundations of College Chemistry 16th edition Hein Solutions Manual

14. (a)
$$\begin{array}{r} 15.2 \\ -2.75 \\ \underline{15.67} \\ 28.1 \end{array}$$

(b) $(4.68)(12.5) = 58.5$

(c) $\dfrac{182.6}{4.6} = 40.$

(d)
$$\begin{array}{r} 1986 \\ 23.84 \\ \underline{\quad 0.012} \\ 2009.852 \end{array} = 2010. = 2.010 \times 10^3$$

(e) $\dfrac{29.3}{(284)(415)} = 2.49 \times 10^{-4}$

(f) $(2.92 \times 10^{-3})(6.14 \times 10^{5}) = 1.79 \times 10^{3}$

15. Fractions to decimals (3 significant figures)

(a) $\dfrac{5}{6} = 0.833$

(b) $\dfrac{3}{7} = 0.429$

(c) $\dfrac{12}{16} = 0.750$

(d) $\dfrac{9}{18} = 0.500$

16. Decimals to fractions

(a) $0.25 = \dfrac{1}{4}$

(b) $0.625 = \dfrac{5}{8}$

(c) $1.67 = 1\dfrac{2}{3}$ or $\dfrac{5}{3}$

(d) $0.888 = \dfrac{8}{9}$

17. (a) $3.42x = 6.5$

$x = \dfrac{6.5}{3.42} = 1.9$

(b) $\dfrac{x}{12.3} = 7.05$

$x = (7.05)(12.3) = 86.7$

(c) $\dfrac{0.525}{x} = 0.25$

$0.525 = 0.25x$

$x = \dfrac{0.525}{0.25} = 2.1$

Foundations of College Chemistry 16th edition Hein Solutions Manual

18. (a) $x = \dfrac{212 - 32}{1.8}$

$x = 1.0 \times 10^{2}$

(c) $72 = 1.8x + 32$

$72 - 32 = 1.8x$

$40. = 1.8x$

$\dfrac{40.}{1.8} = x$

$22 = x$

(b) $8.9\dfrac{\text{g}}{\text{mL}} = \dfrac{40.90\ \text{g}}{x}$

$\left(8.9\dfrac{\text{g}}{\text{mL}}\right)x = 40.90\ \text{g}$

$x = \dfrac{40.90\ \text{g}}{8.9\dfrac{\text{g}}{\text{mL}}} = 4.6\ \text{mL}$

19. (a) $(28.0\ \text{cm})\left(\dfrac{1\ \text{m}}{100\ \text{cm}}\right) = 0.280\ \text{m}$

(b) $(1000.\ \text{m})\left(\dfrac{1\ \text{km}}{1000\ \text{m}}\right) = 1.000\ \text{km}$

(c) $(9.28\ \text{cm})\left(\dfrac{10\ \text{mm}}{1\ \text{cm}}\right) = 92.8\ \text{mm}$

(d) $(10.68\ \text{g})\left(\dfrac{1000\ \text{mg}}{1\ \text{g}}\right) = 1.068 \times 10^{4}\ \text{mg}$

(e) $(6.8 \times 10^{4}\ \text{mg})\left(\dfrac{1\ \text{g}}{1000\ \text{mg}}\right)\left(\dfrac{1\ \text{kg}}{1000\ \text{g}}\right) = 6.8 \times 10^{-2}\ \text{kg}$

(f) $(8.54\ \text{g})\left(\dfrac{1\ \text{kg}}{1000\ \text{g}}\right) = 0.0854\ \text{kg}$

(g) $(25.0\ \text{mL})\left(\dfrac{1\ \text{L}}{1000\ \text{mL}}\right) = 2.50 \times 10^{-2}\ \text{L}$

(h) $(22.4\ \text{L})\left(\dfrac{10^{6}\ \mu\text{L}}{1\ \text{L}}\right) = 2.24 \times 10^{7}\ \mu\text{L}$

20. (a) $(4.5\ \text{cm})\left(\dfrac{1\ \text{m}}{100\ \text{cm}}\right)\left(\dfrac{1\ \text{Å}}{10^{-10}\ \text{m}}\right) = 4.5 \times 10^{8}\ \text{Å}$

(b) $(12\ \text{nm})\left(\dfrac{10^{-9}\ \text{m}}{1\ \text{nm}}\right)\left(\dfrac{100\ \text{cm}}{1\ \text{m}}\right) = 1.2 \times 10^{-6}\ \text{cm}$

Foundations of College Chemistry 16th edition Hein Solutions Manual

(c) $(8.0\text{ km})\left(\frac{1000\text{ m}}{1\text{ km}}\right)\left(\frac{1000\text{ mm}}{1\text{ m}}\right)=8.0\times10^{6}\text{ mm}$

(d) $(164\text{ mg})\left(\frac{1\text{ g}}{1000\text{ mg}}\right)=0.164\text{ g}$

(e) $(0.65\text{ kg})\left(\frac{1000\text{ g}}{1\text{ kg}}\right)\left(\frac{1000\text{ mg}}{1\text{ g}}\right)=6.5\times10^{5}\text{ mg}$

(f) $(5.5\text{ kg})\left(\frac{1000\text{ g}}{1\text{ kg}}\right)=5.5\times10^{3}\text{ g}$

(g) $(0.468\text{ L})\left(\frac{1000\text{ mL}}{1\text{ L}}\right)=468\text{ mL}$

(h) $(9.0\ \mu\text{L})\left(\frac{1\text{ L}}{10^{6}\ \mu\text{L}}\right)\left(\frac{1000\text{ mL}}{1\text{ L}}\right)=9.0\times10^{-3}\text{ mL}$

21. (a) $(42.2\text{ in.})\left(\frac{2.54\text{ cm}}{1\text{ in.}}\right)=107\text{ cm}$

(b) $(0.64\text{ mi})\left(\frac{5280\text{ ft}}{1\text{ mi}}\right)\left(\frac{12\text{ in.}}{1\text{ ft}}\right)=4.1\times10^{4}\text{ in.}$

(c) $(2.00\text{ in.}^{2})\left(\frac{2.54\text{ cm}}{1\text{ in.}}\right)^{2}=12.9\text{ cm}^{2}$

(d) $(42.8\text{ kg})\left(\frac{2.205\text{ lb}}{\text{kg}}\right)=94.4\text{ lb}$

(e) $(3.5\text{ qt})\left(\frac{946\text{ mL}}{1\text{ qt}}\right)=3.3\times10^{3}\text{ mL}$

(f) $(20.0\text{ L})\left(\frac{1\text{ qt}}{0.946\text{ L}}\right)\left(\frac{1\text{ gal}}{4\text{ qt}}\right)=5.29\text{ gal}$

22. (a) The conversation is: m → cm → in. → ft

$(35.6\text{ m})\left(\frac{100\text{ cm}}{1\text{ m}}\right)\left(\frac{1\text{ in.}}{2.54\text{ cm}}\right)\left(\frac{1\text{ ft}}{12\text{ in.}}\right)=117\text{ ft}$

(b) $(16.5\text{ km})\left(\frac{1\text{ mi}}{1.609\text{ km}}\right)=10.3\text{ mi}$

(c) $(4.5\text{ in.}^{3})\left(\frac{2.54\text{ cm}}{1\text{ in.}}\right)^{3}\left(\frac{10\text{ mm}}{1\text{ cm}}\right)^{3}=7.4\times10^{4}\text{ mm}^{3}$

(d) $(95\text{ lb})\left(\frac{453.6\text{ g}}{1\text{ lb}}\right)=4.3\times10^{4}\text{ g}$

Foundations of College Chemistry 16th edition Hein Solutions Manual

(e) $(20.0\text{ gal})\left(\frac{4\text{ qt}}{1\text{ gal}}\right)\left(\frac{0.946\text{ L}}{1\text{ qt}}\right)=75.7\text{ L}$

(f) The conversation is: $\text{ft}^3 \rightarrow \text{in.}^3 \rightarrow \text{cm}^3 \rightarrow \text{m}^3$

$$\left(4.5\times10^4\text{ ft}^3\right)\left(\frac{12\text{ in.}}{1\text{ ft}}\right)^3\left(\frac{2.54\text{ cm}}{1\text{ in.}}\right)^3\left(\frac{1\text{ m}}{100\text{ cm}}\right)^3=1.3\times10^3\text{ m}^3$$

23. First convert 6 ft 6 in. into inches and then convert to centimeters and meters.

$$(6\text{ ft})\left(\frac{12\text{ in.}}{\text{ft}}\right)+6\text{ in.}=78\text{ in.}$$

$$(78\text{ in.})\left(\frac{2.54\text{ cm}}{\text{in.}}\right)=\left(2.0\times10^2\text{ cm}\right)$$

$$\left(2.0\times10^2\text{ cm}\right)\left(\frac{1\text{ m}}{100\text{ cm}}\right)=2.0\text{ m}\left(2\text{ significant figures}\right)$$

24. We must convert ounces to mL.

$$(46.0\text{ oz})\left(\frac{29.6\text{ mL}}{1\text{ oz}}\right)=1.36\times10^3\text{ mL}\left(3\text{ significant figures}\right)$$

25. The conversion is: $\text{L} \rightarrow \text{mL} \rightarrow \text{mg} \rightarrow \text{g}$

$$(1\text{ L})\left(\frac{1000\text{ mL}}{1\text{ L}}\right)\left(\frac{500.\text{ mg}}{100.\text{ mL}}\right)\left(\frac{1\text{ g}}{1000\text{ mg}}\right)=5.00\text{ g}$$

26. The conversion is: $\text{tablet} \rightarrow \text{g} \rightarrow \text{mg} \rightarrow \text{grains}$

$$(1\text{ tablet})\left(\frac{0.500\text{ g}}{1\text{ tablet}}\right)\left(\frac{1000\text{ mg}}{1\text{ g}}\right)\left(\frac{1\text{ grain}}{60\text{ mg}}\right)=8.33\text{ grains}$$

27. The conversion is: $\text{hr} \rightarrow \text{min} \rightarrow \text{s} \rightarrow \text{m} \rightarrow \text{km} \rightarrow \text{mi}$

$$(5\text{ hours})\left(\frac{60\text{ min}}{1\text{ hr}}\right)\left(\frac{60\text{ s}}{1\text{ min}}\right)\left(\frac{0.11\text{ m}}{1\text{ s}}\right)\left(\frac{1\text{ km}}{1000\text{ m}}\right)\left(\frac{1\text{ mi}}{1.61\text{ km}}\right)=1.2\text{ miles}$$

28. The conversion is: $\text{cm} \rightarrow \text{in.} \rightarrow \text{ft} \rightarrow \text{yr} \rightarrow \text{day}$

$$(1\text{ cm})\left(\frac{1\text{ in.}}{2.54\text{ cm}}\right)\left(\frac{1\text{ ft}}{12\text{ in.}}\right)\left(\frac{1\text{ yr}}{3.38\text{ ft}}\right)\left(\frac{365\text{ day}}{1\text{ yr}}\right)=3.54\text{ days}$$

29. The conversion is: $\text{lb} \rightarrow \text{g} \rightarrow \text{mg} \rightarrow \text{ant}$

$$(6.7\text{ lb ant})\left(\frac{454\text{ g ant}}{1\text{ lb ant}}\right)\left(\frac{1000\text{ mg ant}}{1\text{ g ant}}\right)\left(\frac{1\text{ ant}}{85\text{ mg ant}}\right)=3.6\times10^4\text{ ant}$$

Foundations of College Chemistry 16th edition Hein Solutions Manual

30. The conversion is: $\text{lb} \rightarrow \text{g} \rightarrow \text{egg}$

$$(0.75 \text{ lb egg})\left(\frac{454 \text{ g egg}}{1 \text{ lb egg}}\right)\left(\frac{1 \text{ lg egg}}{58.5 \text{ g egg}}\right) = 5.8 \text{ lg eggs}$$

31. The conversion is: $\text{lb body} \rightarrow \text{lb fat} \rightarrow \text{kg fat}$

$$(225 \text{ lb body})\left(\frac{11.2 \text{ lb fat}}{100 \text{ lb body}}\right)\left(\frac{1 \text{ kg}}{2.205 \text{ lb}}\right) = 11.4 \text{ kg fat}$$

32. The conversion is: $\text{carats} \rightarrow \text{mg} \rightarrow \text{g} \rightarrow \text{lb}$

$$(5.75 \text{ carats})\left(\frac{200 \text{ mg}}{1 \text{ carat}}\right)\left(\frac{1 \text{ g}}{1000 \text{ mg}}\right)\left(\frac{1 \text{ lb}}{453.6 \text{ g}}\right) = 2.54\times10^{-3} \text{ lb}$$

33. The conversion is: $\frac{\text{yd}}{\text{s}} \rightarrow \frac{\text{mi}}{\text{s}} \rightarrow \frac{\text{km}}{\text{s}} \rightarrow \frac{\text{m}}{\text{s}} \rightarrow \frac{\text{m}}{\text{min}}$

$$\left(\frac{100. \text{ yd}}{52 \text{ s}}\right)\left(\frac{1 \text{ mi}}{1760 \text{ yd}}\right)\left(\frac{1.609 \text{ km}}{1 \text{ mi}}\right)\left(\frac{1000 \text{ m}}{1 \text{ km}}\right)\left(\frac{60 \text{ s}}{1 \text{ min}}\right) = 1.1\times10^{2} \frac{\text{m}}{\text{min}}$$

34. The conversion is: $\frac{\text{mi}}{\text{hr}} \rightarrow \frac{\text{km}}{\text{hr}} \rightarrow \frac{\text{m}}{\text{hr}} \rightarrow \frac{\text{cm}}{\text{hr}} \rightarrow \frac{\text{cm}}{\text{min}} \rightarrow \frac{\text{cm}}{\text{s}}$

$$\left(\frac{133 \text{ mi}}{1 \text{ hr}}\right)\left(\frac{1.609 \text{ km}}{1 \text{ mi}}\right)\left(\frac{1000 \text{ m}}{1 \text{ km}}\right)\left(\frac{100 \text{ cm}}{1 \text{ m}}\right)\left(\frac{1 \text{ hr}}{60 \text{ min}}\right)\left(\frac{1 \text{ min}}{60 \text{ s}}\right) = 5.94\times10^{3} \frac{\text{cm}}{\text{s}}$$

35. (a) $29{,}035 \text{ ft} - 21{,}002 \text{ ft} = 8033 (\text{total feet climbed})$

$$(16 \text{ hr})\left(\frac{60 \text{ min}}{\text{hr}}\right) = 960 \text{ min} \quad 960 \text{ min} + 42 \text{ min} = 1.0\times10^{3} \text{ min}$$

$$\left(\frac{8003 \text{ ft}}{1.0\times10^{3} \text{ min}}\right)\left(\frac{1 \text{ mi}}{5280 \text{ ft}}\right) = 1.5\times10^{-3} \frac{\text{mi}}{\text{min}}$$

(b) $$\left(\frac{8003 \text{ ft}}{1.0\times10^{3} \text{ min}}\right)\left(\frac{1 \text{ mi}}{5280 \text{ ft}}\right)\left(\frac{1.609 \text{ km}}{\text{mi}}\right)\left(\frac{1000 \text{ m}}{\text{km}}\right)\left(\frac{1 \text{ min}}{60 \text{ s}}\right) = 4.1\times10^{-2} \frac{\text{m}}{\text{s}}$$

36. (a) The conversion is: $\frac{\text{m}}{\text{hr}} \rightarrow \frac{\text{cm}}{\text{hr}} \rightarrow \frac{\text{in.}}{\text{hr}} \rightarrow \frac{\text{ft}}{\text{hr}} \rightarrow \frac{\text{ft}}{\text{min}}$

$$\left(\frac{4500 \text{ m}}{5 \text{ hr}}\right)\left(\frac{100 \text{ cm}}{\text{m}}\right)\left(\frac{1 \text{ in.}}{2.54 \text{ cm}}\right)\left(\frac{1 \text{ ft}}{12 \text{ in.}}\right)\left(\frac{1 \text{ hr}}{60 \text{ min}}\right) = 50 \frac{\text{ft}}{\text{min}}$$

(b) The conversion is: $\frac{\text{m}}{\text{hr}} \rightarrow \frac{\text{km}}{\text{hr}} \rightarrow \frac{\text{km}}{\text{min}} \rightarrow \frac{\text{km}}{\text{s}}$

$$\left(\frac{4500 \text{ m}}{5 \text{ hr}}\right)\left(\frac{1 \text{ km}}{1000 \text{ m}}\right)\left(\frac{1 \text{ hr}}{60 \text{ min}}\right)\left(\frac{1 \text{ min}}{60 \text{ s}}\right) = 3\times10^{-4} \frac{\text{km}}{\text{s}}$$

Foundations of College Chemistry 16th edition Hein Solutions Manual

37. The conversion is: $\frac{\text{mi}}{\text{hr}} \rightarrow \frac{\text{ft}}{\text{hr}} \rightarrow \frac{\text{in.}}{\text{hr}} \rightarrow \frac{\text{cm}}{\text{hr}} \rightarrow \frac{\text{cm}}{\text{min}} \rightarrow \frac{\text{cm}}{\text{s}}$

$$\left(\frac{155 \text{ mi}}{\text{hr}}\right)\left(\frac{5280 \text{ ft}}{1 \text{ mi}}\right)\left(\frac{12 \text{ in}}{1 \text{ ft}}\right)\left(\frac{2.54 \text{ cm}}{1 \text{ in}}\right)\left(\frac{1 \text{ hr}}{60 \text{ min}}\right)\left(\frac{1 \text{ min}}{60 \text{ sec}}\right) = \frac{6930 \text{ cm}}{\text{sec}}$$

38. The conversion is: $\frac{\text{mi}}{\text{hr}} \rightarrow \frac{\text{ft}}{\text{hr}} \rightarrow \frac{\text{in}}{\text{hr}} \rightarrow \frac{\text{cm}}{\text{hr}} \rightarrow \frac{\text{cm}}{\text{min}} \rightarrow \frac{\text{cm}}{\text{s}}$

$$\left(\frac{125 \text{ mi}}{\text{hr}}\right)\left(\frac{5280 \text{ ft}}{1 \text{ mi}}\right)\left(\frac{12 \text{ in}}{1 \text{ ft}}\right)\left(\frac{2.54 \text{ cm}}{1 \text{ in}}\right)\left(\frac{1 \text{ hr}}{60 \text{ min}}\right)\left(\frac{1 \text{ min}}{60 \text{ sec}}\right) = \frac{5590 \text{ cm}}{\text{sec}}$$

39. The conversion is: $\text{gal} \rightarrow \text{qt} \rightarrow \text{L} \rightarrow \text{mL} \rightarrow \text{cup}$

$$(2010 \text{ gal})\left(\frac{4 \text{ qt}}{1 \text{ gal}}\right)\left(\frac{1 \text{ L}}{1.06 \text{ qt}}\right)\left(\frac{1000 \text{ mL}}{1 \text{ L}}\right)\left(\frac{1 \text{ cup}}{473 \text{ mL}}\right) = 1.60 \times 10^4 \text{ cups coffee}$$

40. The conversion is: $\text{lb} \rightarrow \text{g} \rightarrow \text{number of tilapia}$

$$(475 \times 10^6 \text{ lb})\left(\frac{454 \text{ g}}{1 \text{ lb}}\right)\left(\frac{1 \text{ tilapia}}{535 \text{ g}}\right) = 4.03 \times 10^8 \text{ tilapia}$$

41. The conversion is: $\text{gal} \rightarrow \text{qt} \rightarrow \text{mL} \rightarrow \text{drops}$

$$(1.0 \text{ gal})\left(\frac{4 \text{ qt}}{\text{gal}}\right)\left(\frac{946 \text{ mL}}{\text{qt}}\right)\left(\frac{20. \text{ drops}}{\text{mL}}\right) = 7.6 \times 10^4 \text{ drops}$$

42. The conversion is: $\text{gal} \rightarrow \text{qt} \rightarrow \text{L}$

$$(42 \text{ gal})\left(\frac{4 \text{ qt}}{\text{gal}}\right)\left(\frac{0.946 \text{ L}}{\text{qt}}\right) = 160 \text{ L}$$

43. The conversion is: $\text{ft}^3 \rightarrow \text{in.}^3 \rightarrow \text{cm}^3 \rightarrow \text{mL}$

$$(1.00 \text{ ft}^3)\left(\frac{12 \text{ in.}}{1 \text{ ft}}\right)^3\left(\frac{2.54 \text{ cm}}{1 \text{ in.}}\right)^3\left(\frac{1 \text{ mL}}{1 \text{ cm}^3}\right) = 2.83 \times 10^4 \text{ mL}$$

44. $\text{V} = \text{A} \times \text{h}$ $\text{A} = \text{area}$ $\text{h} = \text{height}$ $\text{V} = \text{volume}$

The conversion is: $\frac{\text{cm}^3}{\text{nm}} \rightarrow \frac{\text{cm}^3}{\text{m}} \rightarrow \text{m}^2$

$$\text{A} = \frac{\text{V}}{\text{h}} = \left(\frac{200 \text{ cm}^3}{0.5 \text{ nm}}\right)\left(\frac{1 \text{ nm}}{10^{-9} \text{ m}}\right)\left(\frac{1 \text{ m}}{100 \text{ cm}}\right)^3 = 4 \times 10^5 \text{ m}^2$$

45. (a) $(27 \text{ cm})(21 \text{ cm})(4.4 \text{ cm}) = 2.5 \times 10^3 \text{ cm}^3$

(b) $2.5 \times 10^3 \text{ cm}^3$ is $2.5 \times 10^3 \text{ mL}\left(\frac{1 \text{ L}}{1000 \text{ mL}}\right) = 2.5 \text{ L}$

(c) $(2.5 \times 10^3 \text{ cm}^3)\left(\frac{1 \text{ in.}}{2.54 \text{ cm}}\right)^3 = 1.5 \times 10^2 \text{ in.}^3$

Foundations of College Chemistry 16th edition Hein Solutions Manual

46. $(16\text{ in.})(8\text{ in.})(10\text{ in.})\left(\frac{2.54\text{ cm}}{1\text{ in.}}\right)^3\left(\frac{1\text{ mL}}{1\text{ cm}^3}\right)\left(\frac{1\text{ L}}{1000\text{ mL}}\right)\left(\frac{1\text{ qt}}{0.946\text{ L}}\right)\left(\frac{1\text{ gal}}{4\text{ qt}}\right)=6\text{ gal}$

47. $(250.\text{ g ketone})\left(\frac{73.14\text{ gC}}{100\text{ g ketone}}\right)=183\text{ g C}$

48. $(475.\text{ g nectaryl})\left(\frac{10.98\text{ g H}}{100\text{ g nectaryl}}\right)=52.2\text{ g H}$

49. (a) $\left(\frac{\text{mass part}}{\text{mass whole}}\right)\times 100=\left(\frac{283.4\text{ g potassium dichromate}}{283.4\text{ g potassium dichromate}+650.0\text{ g water}}\right)\times 100$

$=\left(\frac{283.4\text{ g potassium dichromate}}{933.4\text{ g soln}}\right)\times 100=30.36\%\text{ potassium dichromate}$

(b) $(275.\text{ g solution})\left(\frac{30.36\text{ g potassium dichromate}}{100\text{ g solution}}\right)=83.5\text{ g potassium dichromate}$

(c) $(15.0\text{ g potassium dichromate})\left(\frac{100\text{ g soln}}{30.36\text{ g potassium dichromate}}\right)=49.4\text{ g soln}$

50. (a) $\left(\frac{\text{mass part}}{\text{mass whole}}\right)\times 100=\left(\frac{95.4\text{ g sodium nitrate}}{95.4\text{ g sodium nitrate}+753\text{ g water}}\right)\times 100$

$=\left(\frac{95.4\text{ g sodium nitrate}}{848\text{ g soln}}\right)\times 100=11.3\%\text{ sodium nitrate}$

(b) $(350.\text{ g solution})\left(\frac{11.3\text{ g sodium nitrate}}{100\text{ g solution}}\right)=39.6\text{ g sodium nitrate}$

(c) $(50.0\text{ g sodium nitrate})\left(\frac{100\text{ g soln}}{11.3\text{ g sodium nitrate}}\right)=442\text{ g soln}$

51. (a) °F=1.8°C + 32 = (1.8) (38.8) + 32 = 101.8°F

(b) Yes, the child has a fever since 101.8°F > 98.6°F

52. $°\text{F}=1.8°\text{C}+32\quad(1.8)(45)+32=113°\text{F}$ Summer!

53. (a) $\frac{162-32}{1.8}=72.2°\text{C}$ Remember to express the answer to the same precision as the original measurement.

(b) $°\text{C}+273=\text{K}\quad\frac{0.0-32}{1.8}+273=255.2\text{ K}$

(c) 8(−18) + 32 = − 0.40°F

(d) 212 – 273 = − 61°C

54. (a) 1.8(32) + 32 = 90.°F

(b) $\frac{-8.6-32}{1.8}=-22.6°\text{C}$

Foundations of College Chemistry 16th edition Hein Solutions Manual

(c) $273 + 273 = 546\ \text{K}$

(d) $°\text{C} = 100 - 273 = -173°\text{C}$

$(-173)(1.8) + 32 = -279°\text{F} = -300°\text{F}$ (1 significant figure in 100 K)

55. $°\text{C} = \dfrac{(°\text{F} - 32)}{1.8} = \dfrac{(425°\text{F} - 32°\text{F})}{1.8°\text{F}/°\text{C}} = 218°\text{C}$

$K = °\text{C} + 273 = 218 + 273 = 491\ K$

56. $°\text{C} = \dfrac{(°\text{F} - 32)}{1.8} = \dfrac{(247°\text{F} - 32°\text{F})}{1.8°\text{F}/°\text{C}} = 119°\text{C}$

$K = °\text{C} + 273 = 119 + 273 = 392\ K$

57. $°\text{F} = °\text{C}$

$°\text{F} = 1.8(°\text{C}) + 32$

$°\text{F} = 1.8(°\text{F}) + 32$ substitute °F for °C

$-32 = 0.8(°\text{F})$

$\dfrac{-32}{0.8} = °\text{F}$

$-40 = °\text{F}$

$-40°\text{F} = -40°\text{C}$

58. $°\text{F} = -°\text{C}$

$°\text{F} = 1.8(°\text{C}) + 32$ substitute $-°\text{C}$ for °F

$-°\text{C} = 1.8(°\text{C}) + 32$

$2.8(°\text{C}) = -32$

$°\text{C} = \dfrac{-32}{2.8}$

$°\text{C} = -11.4$

$-11.4°\text{C} = 11.4°\text{F}$

59. $°\text{C} = \dfrac{-60° - 32}{1.8} = \dfrac{-92°}{1.8} = -51°\text{C}$

60. $°\text{F} = (1.8 \times 801°) + 32 = 1470°\text{F}$

61. $\text{d} = \dfrac{\text{m}}{\text{v}} = \dfrac{59.82\ \text{g}}{65.0\ \text{mL}} = 0.920\ \dfrac{\text{g}}{\text{mL}}$

62. $\text{d} = \dfrac{\text{m}}{\text{v}} = \dfrac{20.41\ \text{g}}{25.2\ \text{mL}} = 0.810\ \dfrac{\text{g}}{\text{mL}}$

63. $50.92\ \text{g} - 25.23\ \text{g} = 25.69\ \text{g}(\text{mass of liquid})$

$\text{d} = \dfrac{\text{m}}{\text{v}} = \dfrac{25.69\ \text{g}}{25.0\ \text{mL}} = 1.03\ \dfrac{\text{g}}{\text{mL}}$

Foundations of College Chemistry 16th edition Hein Solutions Manual

64. $54.6\text{ mL} - 50.0\text{ mL} = 4.6\text{ mL}(\text{volume of zinc})$

$$d = \frac{m}{v} = \frac{32.95\text{ g}}{4.6\text{ mL}} = 7.2\frac{\text{g}}{\text{mL}}$$

65. The conversion is $\text{g} \rightarrow \text{mL}$

$$(15\text{ g})\left(\frac{1\text{ mL}}{0.929\text{ g}}\right) = 16\text{ mL}$$

66. The conversion is $\text{g} \rightarrow \text{mL}$

$$(75\text{ g})\left(\frac{1\text{ mL}}{1.20\text{ g}}\right) = 63\text{ mL}$$

67. The conversion is kg cayenne $\rightarrow$ kg salsa $\rightarrow$ lb salsa

$$(3.26\text{ kg cayenne})\left(\frac{100\text{ kg salsa}}{8.41\text{ kg cayenne}}\right)\left(\frac{2.20\text{ lb salsa}}{1\text{ kg salsa}}\right) = 85.3\text{ lb}$$

68. The conversion is kg pecan $\rightarrow$ NCB(nutty candy bars) $\rightarrow$ lb NCB

$$(7.0\text{ kg pecan})\left(\frac{100\text{ kg NCB}}{22.0\text{ kg pecan}}\right)\left(\frac{2.20\text{ lb NCB}}{1\text{ kg NCB}}\right) = 70.\text{lb NCB}$$

69. (a) $\frac{91.67\text{ g Cu}}{100\text{ g quarter}}$ and $\frac{8.33\text{ g Ni}}{100\text{ g quarter}}$

(b) The conversion is ton Cu $\rightarrow$ g Cu $\rightarrow$ g quarters $\rightarrow$ quarters

$$(3.0\text{ ton Cu})\left(\frac{2000\text{ lb Cu}}{1\text{ ton Cu}}\right)\left(\frac{454\text{ g Cu}}{1\text{ lb Cu}}\right)\left(\frac{100\text{ g quarters}}{91.67\text{ g Cu}}\right)\left(\frac{1\text{ quarters}}{5.670\text{ g quarters}}\right)$$

$$= 5.2\times10^{5}\text{ quarters}$$

70. (a) $\frac{92.5\text{ g Ag}}{100\text{ g sterling}}$ and $\frac{7.50\text{ g Cu}}{100\text{ g sterling}}$

(b) The conversion is lb Ag $\rightarrow$ g Ag $\rightarrow$ g sterling $\rightarrow$ silver hearts

$$(50.0\text{ lb Ag})\left(\frac{454\text{ g Ag}}{1\text{ lb Ag}}\right)\left(\frac{100\text{ g sterling}}{9.5\text{ g Ag}}\right)\left(\frac{1\text{ silver heart}}{35.0\text{ g sterling}}\right) = 701\text{ silver hearts}$$

71. a) 1055 kg

1) 1055 is the numerical value
2) kg is the unit
3) 4 significant figures
4) 105 is known, 5 is estimated
5) 1.1×10^3 kg

Foundations of College Chemistry 16th edition Hein Solutions Manual

b) 1,650,763 wavelengths
   1) 1,650,763 is the numerical value
   2) wavelengths is the unit
   3) 7 significant figures
   4) 165076 is known, 3 is estimated
   5) 1.7×10^6 wavelengths

c) 1.077 g/cm^3
   1) 1.077 is the numerical value
   2) g/cm^3 is the unit
   3) 4 significant figures
   4) 1.07 is known, 7 is estimated
   5) 1.1×10^0 g/cm^3

d) 0.00845 L
   1) 0.00845 is the numerical value
   2) L is the unit
   3) 3 significant figures
   4) 84 is known, 5 is estimated
   5) 8.5×10^{-3}L

e) 776,000 g
   1) 776,000 is the numerical value
   2) g is the unit
   3) 3 significant figures
   4) 77 is known, 6 is estimated
   5) 7.8×10^5 g

72. (a) report 10.01 grams
    (b) report 10.012 grams
    (c) report 10.0124 grams

73. (a) $(175 \text{ Skittles})\left(\frac{1.134 \text{ g Skittle}}{1 \text{ Skittle}}\right) = 198 \text{ g Skittles}$

    (b) $(175 \text{ Skittles})\left(\frac{5.3 \text{ mL Skittle}}{6 \text{ Skittle}}\right)\left(\frac{1 \text{ L Skittles}}{1000 \text{ mL Skittles}}\right) = 0.15 \text{ L Skittles}$

    (c) $(325.0 \text{ Skittles})\left(\frac{1 \text{ Skittles}}{1.134 \text{ g Skittles}}\right) = 286.6 \text{ Skittles}$

    (d) $(0.550 \text{ L Skittles})\left(\frac{1000 \text{ mL Skittle}}{1 \text{ L Skittle}}\right)\left(\frac{6 \text{ Skittles}}{5.3 \text{ mL Skittles}}\right) = 620 \text{ Skittles}$

Foundations of College Chemistry 16th edition Hein Solutions Manual

(e) The mass measurement is more precise. It is also more accurate. Using calculations similar to those above there should be 309 Skittles in 350 grams and the average value is 310 Skittles. There should be 367 Skittles in 0.325 L of Skittles and the average value is 384 Skittles.

74. A graduated cylinder would be the best choice for adding 100 mL of solvent to a reaction. While the volumetric flask is also labeled 100 mL, volumetric flasks are typically used for doing dilutions. The other three pieces of glassware could also be used, but they hold smaller volumes so it would take a longer time to measure out 100 mL. Also, because you would have to repeat the measurement many times using the other glassware, there is a greater chance for error.

75. The conversion is: $\text{g} \rightarrow \text{mL}$

$$(21.5\text{ g})\left(\frac{1\text{ mL}}{1.484\text{ g}}\right) = 14.5\text{ mL}$$

76. The conversion is: $\text{g} \rightarrow \text{mL}$

$$(25.27\text{ g})\left(\frac{1\text{ mL}}{0.97\text{ g}}\right) = 26\text{ mL}$$

77. The conversion is: $\text{day} \rightarrow \text{cups} \rightarrow \text{mg} \rightarrow \text{g} \rightarrow \text{lb}$

$$(1\text{ day})\left(\frac{4.00\times10^{8}\text{ cups}}{\text{day}}\right)\left(\frac{160\text{ mg}}{\text{cup}}\right)\left(\frac{1\text{ g}}{1000\text{ mg}}\right)\left(\frac{1\text{ lb}}{453.6\text{ g}}\right) = 1\times10^{5}\text{ lb}$$

78. Mass of gear carried by paladin after drinking strength potion = 115 lb + 50.0 lb = 165 lb

Amount of mass potion paladin can carry = 165 lb – 92 lb = 73 lb

The conversion is: $\text{lb} \rightarrow \text{g} \rightarrow \text{mL} \rightarrow \text{vials potion}$

$$(73\text{ lb})\left(\frac{454\text{ g}}{1\text{ lb}}\right)\left(\frac{1\text{ ml}}{93\text{ g}}\right)\left(\frac{1\text{ vial potion}}{50.0\text{ mL}}\right) = 3.43\text{ vials potion}$$

You would only be able to collect 3 vials, because 4 would put you over your mass limit.

79. The conversion is : $\text{sequins} \rightarrow \text{cm}^3 \rightarrow \text{g} \rightarrow \text{kg}$

$$(4560\text{ sequins})\left(\frac{0.0241\text{ cm}^3}{1\text{ sequins}}\right)\left(\frac{41.6\text{ g sequins}}{1\text{ cm}^3}\right)\left(\frac{1\text{ kg}}{1000\text{ g}}\right) = 4.57\text{ kg sequins}$$

The conversion is: $\text{kg} \rightarrow \text{lb}$

$$(4.57\text{ kg sequins})\left(\frac{2.20\text{ lb}}{1\text{ kg}}\right) = 10.1\text{ lb sequins}$$

Foundations of College Chemistry 16th edition Hein Solutions Manual

80. $V = \text{side}^3 = (0.50\ \text{m})^3 = 0.13\ \text{m}^3$ (volume of the cube)

$$(0.13\ \text{m}^3)\left(\frac{100.\ \text{cm}}{\text{m}}\right)^3\left(\frac{1\ \text{L}}{1000\ \text{cm}^3}\right) = 130\ \text{L}$$ (volume of cube)

Yes, the cube will hold the solution. 130 L – 8.5 L = 120 L additional solution is necessary to fill the container.

81. The conversion is: $\frac{\mu\text{g}}{\text{m}^3} \rightarrow \frac{\mu\text{g}}{\text{L}} \rightarrow \frac{\mu\text{g}}{\text{day}}$

$$\left(\frac{180\ \mu\text{g}}{1\ \text{m}^3}\right)\left(\frac{1\ \text{m}^3}{1000\ \text{L}}\right)\left(2\times10^4\ \frac{\text{L}}{\text{day}}\right) = 4000\ \mu\text{g ingested/day} \quad \text{(1 significant figure)}$$

Yes, the nurse is at risk. This is well over the toxic limit.

82. (a) The conversion is ac ft → L → kL

$$(1380\ \text{ac ft})\left(\frac{1.233\times10^6\ \text{L}}{1\ \text{ac ft}}\right)\left(\frac{1\ \text{kL}}{1000\ \text{L}}\right) = 1.70\times10^6\ \text{kL}$$

(b) The conversion is day → kL → L → qt → gal

$$(31\ \text{day})\left(\frac{1.70\times10^6\ \text{kL}}{1\ \text{day}}\right)\left(\frac{1000\ \text{L}}{1\ \text{kL}}\right)\left(\frac{1.06\ \text{qt}}{1\ \text{L}}\right)\left(\frac{1\ \text{gal}}{4\ \text{qt}}\right) = 1.40\times10^{10}\ \text{gal}$$

83. (a) Convert 20.27 K to °C

$\text{K} - 273.15 = °\text{C}$

$20.27\ \text{K} - 273.15 = -252.88°\text{C}$

(b) Convert 20.27 K to °F

$°\text{F} = (1.8\times °\text{C}) + 32$

$°\text{F} = (1.8\times -252.88) + 32 = -455.18 + 32$

$°\text{F} = -423.18°\text{F}$

84. $°\text{F} = 1.8°\text{C} + 32$

Convert 36°C to °F $(1.8)(36) + 32 = 97°\text{F}$

Convert 38°C to °F $(1.8)(38) + 32 = 100°\text{F}$

Sauropods have a body temperature close to the body temperature of other warm-blooded mammals such as dogs, humans, and cows.

85. The conversion is : L → dL → mg → g

$$(4.7\ \text{L})\left(\frac{10\ \text{dL}}{1\ \text{L}}\right)\left(\frac{130\ \text{mg}}{1\ \text{dL}}\right)\left(\frac{1\ \text{g}}{1000\ \text{mg}}\right) = 6.1\ \text{g}$$

86. A sample of gold will sink to the bottom of the mercury and a sample of iron pyrite will float.

Foundations of College Chemistry 16th edition Hein Solutions Manual

87. The conversion is : El → ton → lb

$$(5.3\text{ El})\left(\frac{6.00\text{ ton}}{1\text{ El}}\right)\left(\frac{2000\text{ lb}}{1\text{ ton}}\right)=6.4\times10^{4}\text{ lb}$$

88. The conversion is: hands → in. → cm → m

$$(14.2\text{ hands})\left(\frac{4\text{ in.}}{1\text{ hand}}\right)\left(\frac{2.54\text{ cm}}{1\text{ in.}}\right)\left(\frac{1\text{ m}}{100\text{ cm}}\right)=1.44\text{ m}$$

89. The conversion is: days → hr → gal → qt → L

$$(30\text{ days})\left(\frac{24\text{ hr}}{1\text{ day}}\right)\left(\frac{22.5\text{ gal}}{12\text{ hr}}\right)\left(\frac{4\text{ qt}}{1\text{ gal}}\right)\left(\frac{0.946\text{ L}}{1\text{ qt}}\right)=5.1\times10^{3}\text{ L}$$

90. $d=\frac{\text{m}}{\text{V}}$ The cube with the largest volume has the lowest density. Use Table 2.6.

Cube A - lowest density 1.74 g/mL - magnesium
Cube B 2.7 g/mL - aluminium
Cube C - highest density 10.5 g/mL - silver

91. The conversion is: ft → in. → cm → m → nm → nanotubes

$$(40.0\text{ ft})\left(\frac{12\text{ in.}}{1\text{ ft}}\right)\left(\frac{2.54\text{ cm}}{1\text{ in.}}\right)\left(\frac{1\text{ m}}{100\text{ cm}}\right)\left(\frac{10^{9}\text{ nm}}{1\text{ m}}\right)\left(\frac{1\text{ nanotube}}{1.3\text{ nm}}\right)=9.4\times10^{9}\text{ nanotubes}$$

92. $$(380\text{ mg Mg})\left(\frac{0.25\text{ c peanuts}}{67\text{ mg Mg}}\right)=1.4\text{ c peanuts}$$

93. The conversion is: acres → ft^2 → mi^2 → km^2

$$(125\text{ acres})\left(\frac{43560\text{ ft}^2}{1\text{ acre}}\right)\left(\frac{1\text{ mi}}{5280\text{ ft}}\right)^2\left(\frac{1.609\text{ km}}{1\text{ mi}}\right)^2=0.506\text{ km}^2$$

94. The volume of the aluminum cube is

$$\text{V}=\frac{\text{m}}{d}=\frac{500.\text{ g}}{2.70\frac{\text{g}}{\text{mL}}}=185\text{ mL}$$

Density of Al is 2.70 g/mL.

This is the same volume as the gold cube thus

$\text{m}=d\text{V}=(185\text{ mL})(19.3\text{ g/mL})=3.57\times10^{3}$ g of gold

Density of Au is 19.3 g/mL.

95. $d=\frac{\text{m}}{\text{V}}=\frac{24.12\text{ g}}{25.0\text{ mL}}=\frac{0.965\text{ g}}{\text{mL}}$ (density of water at 90°C)

96. $150.50\text{ g}-88.25\text{ g}=62.25\text{ g}$ (mass of liquid)

$d=\frac{\text{m}}{\text{V}}$ thus $\text{V}=\frac{\text{m}}{d}=\frac{62.25\text{ g}}{1.25\frac{\text{g}}{\text{mL}}}=49.8\text{ mL}$ (volume of liquid)

The container must hold at least 50 mL.

Foundations of College Chemistry 16th edition Hein Solutions Manual

97. H_2O $\dfrac{50\text{ g}}{1.0\dfrac{\text{g}}{\text{mL}}} = 50\text{ mL}$

alcohol $\dfrac{50\text{ g}}{0.789\dfrac{\text{g}}{\text{mL}}} = 60\text{ mL}$

Ethyl alcohol has the greater volume due to its lower density.


98. Volume of sulfuric acid

$$\left(\frac{1\text{ mL}}{1.84\text{ g}}\right)(100.\text{ g}) = 54.3\text{ mL}$$

99. The conversion is: $\text{cup} \rightarrow \text{oz} \rightarrow \text{qt} \rightarrow \text{L} \rightarrow \text{mg}$

$$(2.00\text{ cup coffee})\left(\frac{10\text{ oz coffee}}{1\text{ cup coffee}}\right)\left(\frac{1\text{ qt coffee}}{32\text{ oz coffee}}\right)\left(\frac{1\text{ L coffee}}{1.057\text{ qt coffee}}\right)\left(\frac{31.4\text{ mg NMP}}{1\text{ L coffee}}\right)$$
$$= 18.6\text{ mg NMP}$$

100.



Since $d = \dfrac{\text{m}}{\text{V}}$, as the volume increases, the density decreases. As solids are heated the density decreases due to an increase in the volume of the solid.

101. $\text{V} = (2.00\text{ cm})(15.0\text{ cm})(6.00\text{ cm})\left(\dfrac{1\text{ mL}}{1\text{ cm}^3}\right) = 180.\text{mL}\ \ (\text{volume of bar})$

$$d = \frac{\text{m}}{\text{V}} = 3300\text{ g}/180.\text{ mL} = 18.3\text{ g/mL}$$

The density of pure gold is 19.3 g/mL (from Table 2.5), therefore, the gold bar is not pure gold, since its density is only 18.3 g/mL, or it is hollow inside.

102. $\text{m} = d\text{V} = (0.789\text{ g/mL})(35.0\text{ mL}) = 27.6\text{ g ethyl alcohol}$

$27.6\text{ g} + 49.28\text{ g} = 76.9\text{ g}\ \ (\text{mass of cylinder and alcohol})$

103. The conversion is $\text{g} \rightarrow \text{oz} \rightarrow \text{gr} \rightarrow \text{scruples}$

$$(695\text{ g})\left(\frac{12\text{ oz}}{373\text{ g}}\right)\left(\frac{480\text{ gr}}{\text{oz}}\right)\left(\frac{1\text{ scruple}}{20\text{ gr}}\right) = 537\text{ scruples}$$

Foundations of College Chemistry 16th edition Hein Solutions Manual

104. The conversion is: days → teaspoons → mL

$$(10 \text{ days})\left(\frac{2 \text{ teaspoons} \times 4}{\text{day}}\right)\left(\frac{5 \text{ mL}}{\text{teaspoon}}\right) = 400 \text{ mL}$$

Since you will need a total of 400 mL for the 10 days and the bottle contains 500 mL, you have purchased enough.

105. (a) The conversion is kg → g → mL → L

$$(8.14 \text{ kg brine})\left(\frac{1000 \text{ g brine}}{1 \text{ kg brine}}\right)\left(\frac{1 \text{ mL brine}}{1.28 \text{ g brine}}\right)\left(\frac{1 \text{ L brine}}{1000 \text{ mL brine}}\right) = 6.36 \text{ L brine}$$

(b) The conversion is g NaCl → g brine

$$(150.0 \text{ g NaCl})\left(\frac{100 \text{ g brine}}{7.55 \text{ g NaCl}}\right) = 1990 \text{ g brine}$$

106. The density of lead is 11.34 g/mL. The density of aluminum is 2.70 g/mL. The density of silver is 10.5 g/mL. The density of the unknown piece of metal can be calculated from the mass (20.25 g) and the volume (57.5 mL − 50 mL = 7.5 mL) of the metal. Density of the unknown metal = 20.25 g/7.5 mL = 2.7 g/mL. The metal must be aluminum.

107.

Volume of slug	$30.7 \text{ mL} - 25.0 \text{ mL} = 5.7 \text{ mL}$
Density of slug	$d = \frac{\text{m}}{\text{V}} = \frac{15.454 \text{ g}}{5.7 \text{ mL}} = 2.7 \text{ g/mL}$
Mass of liquid, cylinder, and slug	125.934 g
Mass of slug (subtract)	−15.454 g
Mass of cylinder (subtract)	−89.450 g
Mass of the liquid	21.030 g

Density of liquid $d = \frac{\text{m}}{\text{V}} = \frac{21.030 \text{ g}}{25.0 \text{ mL}} = 0.841 \text{ g/mL}$

108. $$(225 \text{ mL bronze needed})\left(\frac{7.8 \text{ g bronze needed}}{1 \text{ mL bronze needed}}\right)\left(\frac{100 \text{ g bronze melted}}{90.0 \text{ g bronze needed}}\right)$$
$$= 2.0 \times 10^3 \text{ g bronze melted}$$

109. The conversion is: km → m → cm → in. → ft → mi → hr → min → s → ns

$$(730.0 \text{ km})\left(\frac{1000 \text{ m}}{1 \text{ km}}\right)\left(\frac{100 \text{ cm}}{1 \text{ m}}\right)\left(\frac{1 \text{ in.}}{2.54 \text{ cm}}\right)\left(\frac{1 \text{ ft}}{12 \text{ in.}}\right)\left(\frac{1 \text{ mi}}{5280 \text{ ft}}\right)\left(\frac{1 \text{ hr}}{1.86 \times 10^8 \text{ mi}}\right)\left(\frac{60 \text{ min}}{1 \text{ hr}}\right)\left(\frac{60 \text{ s}}{1 \text{ min}}\right)$$
$$\left(\frac{10^9 \text{ ns}}{1 \text{ s}}\right) = 8.78 \times 10^6 \text{ ns}$$

8,780,000 ns is good to only 3 significant figures. You would need at least 6 significant digits to detect the difference between 8,780,000 ns and (8,780,000 + 60) ns. Sometimes experimenters do not see differences due to the precision of their measuring techniques.

Foundations of College Chemistry 16th edition Hein Solutions Manual

110. $(1.5\text{ m})\left(\frac{100\text{ cm}}{1\text{ m}}\right)\left(\frac{1\text{ in.}}{2.54\text{ cm}}\right)\left(\frac{1\text{ ft}}{12\text{ in.}}\right)=4.9\text{ ft}$

$(4\text{ m})\left(\frac{100\text{ cm}}{1\text{ m}}\right)\left(\frac{1\text{ in.}}{2.54\text{ cm}}\right)\left(\frac{1\text{ ft}}{12\text{ in.}}\right)=13\text{ ft}$

$(27°\text{C}\times 1.8)+32=81°\text{F}$

Foundations of College Chemistry 16th edition Hein Solutions Manual