Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

SKU: 9781138044494-SOLUTIONS

Chapter 2

Solving linear systems

2.1 Vectors and linear equations 11
2.2 Matrix operations 16
2.3 Inverse matrices 19
2.4 LU decomposition 24
2.5 Transpose and permutation 25

2.1 Vectors and linear equations

Exercise 2.1.1.

1. Redo Example 2.1.1 with the first elementary row operation $R_2 - R_1$.

Solution:

System	Matrix representation	Elementary matrix
$\begin{cases} x - y = 1 \\ x + y = 2 \end{cases}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$	
$\Downarrow R_2 - R_1$		$E_1 = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$
$\begin{cases} x - y = 1 \\ 2y = 1 \end{cases}$	$\begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$	
$\Downarrow \frac{1}{2} R_2$	$\begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$	$E_3 = \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}$
$\begin{cases} x - y = 1 \\ y = \frac{1}{2} \end{cases}$	$\begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$	
$\Downarrow R_1 + R_2$	$\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}$	$E_4 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$
$\begin{cases} x = \frac{3}{2} \\ y = \frac{1}{2} \end{cases}$	$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}$	

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{3}{2} \\ \frac{1}{2} \end{bmatrix}.$$

2. Determine whether the following matrices are in reduced row echelon form and row echelon form, respectively:

$$a) \begin{bmatrix} 1 & 0 & 0 & 9 & -2 \\ 0 & 1 & 0 & -2 & \frac{1}{2} \\ 0 & 0 & 1 & -5 & \frac{1}{2} \end{bmatrix}, \quad b) \begin{bmatrix} 1 & 0 & 0 & 9 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -5 \end{bmatrix}, \quad c) \begin{bmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & \frac{1}{2} \\ 0 & 0 & 1 & \frac{1}{2} \end{bmatrix}.$$

Solution: a) and b) are in reduced row echelon form (hence also in row echelon form). c) is in row echelon form. □

3. Solve the following systems using Gauss–Jordan eliminations:

$$a) \begin{cases} x + 3z = 1 \\ 2x + 3y = 3 \\ 4y + 5z = 5 \end{cases} \quad b) \begin{cases} x + 2y + 3z = 1 \\ 2x + 3y + 4z = 3 \\ 3x + 4y + 5z = 5 \end{cases} \quad c) \begin{cases} x + 2y + 3z = 1 \\ 2x + 3y + 4z = 3 \\ 5x + 9y + 13z = 7 \end{cases}$$

Solution: a) We re-write the system of linear equations in the matrix form $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 1 & 0 & 3 \\ 2 & 3 & 0 \\ 0 & 4 & 5 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix}.$$

Then the corresponding augmented matrix is

$$[A : \mathbf{b}] = \begin{bmatrix} 1 & 0 & 3 & 1 \\ 2 & 3 & 0 & 3 \\ 0 & 4 & 5 & 5 \end{bmatrix}.$$

By the elementary row operations on $[A : \mathbf{b}]$ we have

$$\begin{bmatrix} 1 & 0 & 3 & 1 \\ 2 & 3 & 0 & 3 \\ 0 & 4 & 5 & 5 \end{bmatrix} \xrightarrow{R_2 - 2R_1} \begin{bmatrix} 1 & 0 & 3 & 1 \\ 0 & 3 & -6 & 1 \\ 0 & 4 & 5 & 5 \end{bmatrix}$$

$$\xrightarrow{R_2/3} \begin{bmatrix} 1 & 0 & 3 & 1 \\ 0 & 1 & -2 & \frac{1}{3} \\ 0 & 4 & 5 & 5 \end{bmatrix}$$

$$\xrightarrow{R_3 - 4R_2} \begin{bmatrix} 1 & 0 & 3 & 1 \\ 0 & 1 & -2 & \frac{1}{3} \\ 0 & 0 & 13 & \frac{11}{3} \end{bmatrix}$$

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

$$\xrightarrow{R_3/13} \begin{bmatrix} 1 & 0 & 3 & 1 \\ 0 & 1 & -2 & \frac{1}{3} \\ 0 & 0 & 1 & \frac{11}{39} \end{bmatrix}$$

$$\xrightarrow[R_1-3R_3]{R_2+2R_3} \begin{bmatrix} 1 & 0 & 0 & \frac{6}{39} \\ 0 & 1 & 0 & \frac{35}{39} \\ 0 & 0 & 1 & \frac{11}{39} \end{bmatrix}.$$

Then we have an equivalent system with augmented matrix

$$\begin{bmatrix} 1 & 0 & 0 & \frac{6}{39} \\ 0 & 1 & 0 & \frac{35}{39} \\ 0 & 0 & 1 & \frac{11}{39} \end{bmatrix}.$$

The solution is

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{6}{39} \\ \frac{35}{39} \\ \frac{11}{39} \end{bmatrix}.$$

b) We re-write the system of linear equations in the matrix form $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix}.$$

Then the corresponding augmented matrix is

$$[A : \mathbf{b}] = \begin{bmatrix} 1 & 2 & 3 & 1 \\ 2 & 3 & 4 & 3 \\ 3 & 4 & 5 & 5 \end{bmatrix}.$$

By the elementary row operations on $[A : \mathbf{b}]$ we have

$$\begin{bmatrix} 1 & 2 & 3 & 1 \\ 2 & 3 & 4 & 3 \\ 3 & 4 & 5 & 5 \end{bmatrix} \xrightarrow[R_3-3R_1]{R_2-2R_1} \begin{bmatrix} 1 & 2 & 3 & 1 \\ 0 & -1 & -2 & 1 \\ 0 & -2 & -4 & 2 \end{bmatrix}$$

$$\xrightarrow{(-1)\cdot R_2} \begin{bmatrix} 1 & 2 & 3 & 1 \\ 0 & 1 & 2 & -1 \\ 0 & -2 & -4 & 2 \end{bmatrix}$$

$$\xrightarrow[R_1-2R_2]{R_3+2R_2} \begin{bmatrix} 1 & 0 & -1 & 3 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then we have an equivalent system with augmented matrix

$$\begin{bmatrix} 1 & 0 & -1 & 3 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

The solution is

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \\ 0 \end{bmatrix} + t \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}, \, t \in \mathbb{R}.$$

c) We re-write the system of linear equations in the matrix form $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 5 & 9 & 13 \end{bmatrix}, \, \mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} 1 \\ 3 \\ 7 \end{bmatrix}.$$

Then the corresponding augmented matrix is

$$[A : \mathbf{b}] = \begin{bmatrix} 1 & 2 & 3 & 1 \\ 2 & 3 & 4 & 3 \\ 5 & 9 & 13 & 7 \end{bmatrix}.$$

By the elementary row operations on $[A : \mathbf{b}]$ we have

$$\begin{bmatrix} 1 & 2 & 3 & 1 \\ 2 & 3 & 4 & 3 \\ 5 & 9 & 13 & 7 \end{bmatrix} \xrightarrow[R_3 - 5R_1]{R_2 - 2R_1} \begin{bmatrix} 1 & 2 & 3 & 1 \\ 0 & -1 & -2 & 1 \\ 0 & -1 & -2 & 2 \end{bmatrix}$$

$$\xrightarrow{(-1)\cdot R_2} \begin{bmatrix} 1 & 2 & 3 & 1 \\ 0 & 1 & 2 & -1 \\ 0 & -1 & -2 & 2 \end{bmatrix}$$

$$\xrightarrow[R_1 - 2R_2]{R_3 + R_2} \begin{bmatrix} 1 & 0 & -1 & 3 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Then we have an equivalent system with augmented matrix

$$\begin{bmatrix} 1 & 0 & -1 & 3 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

where the last equation is contradictory. The system has no solution. □

□

4. Consider a linear system $Ax = b$ with A an $m \times n$ matrix and b an $m \times 1$ matrix. Is it true if there is more than one solution for x in $\mathbb{R}^n$, there must be infinitely many? You may use the fact that

$$A(x + y) = Ax + Ay$$
$$A(tx) = tAx,$$

for every x, $y \in \mathbb{R}^n$ and $t \in \mathbb{R}$,

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

which is called the **linearity** of matrix multiplication.

Solution: Let x_1 and x_2 be distinct solutions. Then $A(x_1-x_2) = Ax_1-Ax_2 = 0$ and for every $t \in \mathbb{R}$, we have

$$A(x_1 + t(x_1 - x_2)) = Ax_1 + tA(x_1 - x_2) = b.$$

Hence $x = x_1 + t(x_1 - x_2)$, $t \in \mathbb{R}$ are all solutions of $Ax = b$. That is, $Ax = b$ has infinitely many solutions. □

5. Let k be a real number. Consider the following linear system of equations:

$$\begin{cases} x_2 + 2x_3 + x_4 = 1 \\ 2x_1 + x_2 + 3x_3 = 2 \\ x_1 + 4x_3 + 2x_4 = 3 \\ kx_2 + x_4 = 1. \end{cases} \tag{2.1}$$

Find all possible values of k such that system (2.1) i) has a unique solution; ii) has no solutions and iii) has infinitely many solutions.

Solution: We re-write the system of linear equations in the matrix form $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 0 & 1 & 2 & 1 \\ 2 & 1 & 3 & 0 \\ 1 & 0 & 4 & 2 \\ 0 & k & 0 & 1 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 1 \end{bmatrix}.$$

Then the corresponding augmented matrix is

$$[A : \mathbf{b}] = \begin{bmatrix} 0 & 1 & 2 & 1 & 1 \\ 2 & 1 & 3 & 0 & 2 \\ 1 & 0 & 4 & 2 & 3 \\ 0 & k & 0 & 1 & 1 \end{bmatrix}.$$

By the elementary row operations on $[A : \mathbf{b}]$ we have

$$\begin{bmatrix} 0 & 1 & 2 & 1 & 1 \\ 2 & 1 & 3 & 0 & 2 \\ 1 & 0 & 4 & 2 & 3 \\ 0 & k & 0 & 1 & 1 \end{bmatrix} \xrightarrow{R_3 \leftrightarrow R_1} \begin{bmatrix} 1 & 0 & 4 & 2 & 3 \\ 2 & 1 & 3 & 0 & 2 \\ 0 & 1 & 2 & 1 & 1 \\ 0 & k & 0 & 1 & 1 \end{bmatrix}$$

$$\xrightarrow{R_2 - 2R_1} \begin{bmatrix} 1 & 0 & 4 & 2 & 3 \\ 0 & 1 & -5 & -4 & -4 \\ 0 & 1 & 2 & 1 & 1 \\ 0 & k & 0 & 1 & 1 \end{bmatrix}$$

$$\xrightarrow[R_3 - R_2]{R_4 - kR_2} \begin{bmatrix} 1 & 0 & 4 & 2 & 3 \\ 0 & 1 & -5 & -4 & -4 \\ 0 & 0 & 7 & 5 & 5 \\ 0 & 0 & 5k & 1+4k & 1+4k \end{bmatrix}$$

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

$$\xrightarrow{R_3/7} \begin{bmatrix} 1 & 0 & 4 & 2 & 3 \\ 0 & 1 & -5 & -4 & -4 \\ 0 & 0 & 1 & 5/7 & 5/7 \\ 0 & 0 & 5k & 1+4k & 1+4k \end{bmatrix}$$

$$\xrightarrow{R_4-(5k)\cdot R_3} \begin{bmatrix} 1 & 0 & 4 & 2 & 3 \\ 0 & 1 & -5 & -4 & -4 \\ 0 & 0 & 1 & 5/7 & 5/7 \\ 0 & 0 & 0 & 1+3k/7 & 1+3k/7 \end{bmatrix}.$$

Then we have an equivalent system with augmented matrix

$$\begin{bmatrix} 1 & 0 & 4 & 2 & 3 \\ 0 & 1 & -5 & -4 & -4 \\ 0 & 0 & 1 & 5/7 & 5/7 \\ 0 & 0 & 0 & 1+3k/7 & 1+3k/7 \end{bmatrix},$$

i) If $1+3k/7 \neq 0$, that is, $k \neq -7/3$ the original system has a unique solutiuon;

ii) If $1+3k/7 = 0$, that is, $k = -7/3$ the original system has infinitely many solutions.

Therefore, for every $k \in \mathbb{R}$, the system has at least one solution. □

2.2 Matrix operations

Exercise 2.2.1.

1. Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$. i) Compute AB. ii) Does BA exist?

Solution: i)

$$AB = \begin{bmatrix} 1 & 2 & 1 & 2 \\ 3 & 4 & 3 & 4 \end{bmatrix}.$$

ii) BA does not exist since their sizes do not match as B is 2×4 and A is 2×2. □

2. Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 1 & -1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix}$. i) Compute AB. ii) If B is block partitioned into $B = [B_1 : B_2]$, is it true $AB = [AB_1 : AB_2]$?

Solution: i)

$$AB = \begin{bmatrix} 3 & 1 & 2 & 1 \\ 7 & 1 & 4 & 3 \end{bmatrix}$$

ii) According to the definition of matrix multiplication, the statement is true. □

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

3. Show Lemma 2.2.6.

Solution: Follow the approach for the proof of Theorem 2.2.5. □

4. Let $A = [a_1 : a_2 \cdots : a_n]$, $B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$ be $m \times n$ and $n \times r$ matrices. Show that $AB = a_1b_1 + a_2b_2 + \cdots + a_nb_n$.

Solution: By the definition of matrix multiplication we have

$$\begin{aligned}(AB)_{i,j} &= a_{i,1}b_{1,j} + a_{i,2}b_{2,j} + \cdots + a_{i,n}b_{n,j} \\ &= (a_1b_1)_{i,j} + (a_2b_2)_{i,j} + \cdots + (a_nb_n)_{i,j} \\ &= (a_1b_1 + a_2b_2 + \cdots + a_nb_n)_{i,j}.\end{aligned}$$

Therefore, we have $AB = a_1b_1 + a_2b_2 + \cdots + a_nb_n$. □

5. Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 1 & -1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix}$. Use Question 4 to compute AB.

Solution: Let $A = [a_1 : a_2]$, $B = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$

$$\begin{aligned}AB &= a_1b_1 + a_2b_2 \\ &= \begin{bmatrix} 1 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & -1 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 2 \\ 4 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1 & -1 & 0 & 1 \\ 3 & -3 & 0 & 3 \end{bmatrix} + \begin{bmatrix} 2 & 2 & 2 & 0 \\ 4 & 4 & 4 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 3 & 1 & 2 & 1 \\ 7 & 1 & 4 & 3 \end{bmatrix}.\end{aligned}$$

□

6. Let A and B be $m \times n$ and $n \times r$ matrices. Show that i) every column of AB is a linear combination of the columns of A; ii) every row of AB is a linear combination of the rows of B.

Solution: Let $A = [a_1 : a_2 \cdots : a_n]$, $B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$ be $m \times n$ and $n \times r$ matrices.

$$(AB)_{i,j} = a_{i,1}b_{1,j} + a_{i,2}b_{2,j} + \cdots + a_{i,n}b_{n,j}.$$

The j-th column of AB is

$$(AB)_j = a_1b_{1,j} + a_2b_{2,j} + \cdots + a_nb_{n,j},$$

which is a linear combination of the columns of A. □

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

7. Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$. Find all matrices B such that $AB = BA$.

Solution: Let $B = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then $BA = AB$ leads to a linear system of $x = (a, b, c, d) \in \mathbb{R}^4$:

$$\begin{cases} 3b - 2c = 0 \\ 2a + 3b - 2d = 0 \\ 3a + 3c - 3d = 0 \\ 3b - 2c = 0. \end{cases}$$

Using Guass-Jordan elimination, we obtain the solution

$$x = s \begin{bmatrix} -1 \\ 2/3 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \ s, t \in \mathbb{R}.$$

That is,

$$B = \begin{bmatrix} t - s & 2s/3 \\ s & t \end{bmatrix}.$$

8. Let A and B be $n \times n$ matrices. Explain that in general we have $(A + B)(A - B) \neq A^2 - B^2$ and $(A + B)^2 \neq A^2 + 2AB + B^2$.

Solution: We note that in general, $AB \neq BA$. So $(A + B)(A - B) = A^2 - AB + BA - B^2 \neq A^2 - B^2$ and $(A + B)^2 \neq A^2 + 2AB + B^2$ if $AB \neq BA$.

9. Let A be an $n \times n$ matrix. Define $V = \{B : AB = BA\}$. Show that i) $V \neq \emptyset$; ii) if $B_1 \in V$ and $B_2 \in V$, then every linear combination of B_1 and B_2 is in V.

Solution: i) Since $AI = IB$ and $I \in V$, we have $V \neq \emptyset$.

ii) If $B_1 \in V$ and $B_2 \in V$, then for every $s \in \mathbb{R}$ and $t \in \mathbb{R}$, we have

$$\begin{aligned} (sB_1 + tB_2)A =& sB_1A + tB_2A \\ =& sAB_1 + tAB_2 \\ =& A(sB_1 + tB_2). \end{aligned}$$

Therefore, we have $sB_1 + tB_2 \in V$. We have shown that V is a vector space. □

10. Give an example that $A^2 = 0$ but $A \neq 0$.

Solution:

$$A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}.$$

We have $A^2 = 0$ but $A \neq 0$.

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

11. Give an example that $A^2 = I$ but $A \neq \pm I$.

Solution:

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

We have $A^2 = I$ but $A \neq \pm I$.

12. Let A be an $n \times n$ matrix. If we want to define a limit $\lim_{m\to\infty} A^m$, how would you define the closeness (distance) between matrices?

Solution: If we identify A as a vector in $\mathbb{R}^{n^2}$. Then we can borrow norms on $\mathbb{R}^{n^2}$ for defining the distance between matrices. For example, we can define

$$\text{distance}(A,\, B) = \left(\sum_{i=1}^{n} \sum_{j=1}^{n} (a_{ij} - b_{ij})^2 \right)^{\frac{1}{2}},$$

where a_{ij} and b_{ij} are entries at $(i,\, j)$-position of A and B, respectively. □

2.3 Inverse matrices

Exercise 2.3.1.

1. Determine whether or not the following matrices are invertible. Find the inverse of each matrix if it exists.

$$a) \quad \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \quad b) \quad \begin{bmatrix} -1 & 2 \\ 3 & 6 \end{bmatrix}, \quad c) \quad \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}.$$

Solution: a) $\begin{bmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix}$, b) $\begin{bmatrix} -\frac{1}{2} & \frac{1}{6} \\ \frac{1}{4} & \frac{1}{12} \end{bmatrix}$, c) not invertible.

□

2. Determine whether or not the following matrices are invertible. Find the inverse of each matrix if it exists.

$$a) \quad \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \quad b) \quad \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 3 & 1 \end{bmatrix}, \quad c) \quad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Solution: a) $\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$, b) $\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 6 & -3 & 1 \end{bmatrix}$, b) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{6} & 0 \\ 0 & 0 & 1 \end{bmatrix}$

□

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

3. Determine whether or not the following matrices are invertible. Find the inverse of each matrix if it exists.

$$a)\ \begin{bmatrix}1&2&0\\3&4&0\\0&0&1\end{bmatrix},\quad b)\ \begin{bmatrix}1&0&0\\0&-1&2\\0&3&6\end{bmatrix},\quad c)\ \begin{bmatrix}1&2&0\\3&6&0\\0&0&1\end{bmatrix}.$$

Solution: a) $\begin{bmatrix}-2&1&0\\\frac{3}{2}&-\frac{1}{2}&0\\0&0&1\end{bmatrix}$, b) $\begin{bmatrix}1&0&0\\0&-\frac{1}{2}&\frac{1}{6}\\0&\frac{1}{4}&\frac{1}{12}\end{bmatrix}$, c) not invertible.

□

4. For the given matrix A, use elimination to find A^{-1} and record each elementary row operation and the corresponding elementary matrix at the same time.

$$A=\begin{bmatrix}3&0&1\\2&4&2\\1&-1&5\end{bmatrix}.$$

Solution:

$$\begin{bmatrix}3&0&1&1&0&0\\2&4&2&0&1&0\\1&-1&5&0&0&1\end{bmatrix}$$

Row operation: $R_1 \leftrightarrow R_3$ — Elementary Matrix: $E_1=\begin{bmatrix}0&0&1\\0&1&0\\1&0&0\end{bmatrix}$

$$\Downarrow$$

$$\begin{bmatrix}1&-1&5&0&0&1\\2&4&2&0&1&0\\3&0&1&1&0&0\end{bmatrix}$$

Row operation: R_2-2R_1 — Elementary Matrix: $E_2=\begin{bmatrix}1&0&0\\-2&1&0\\0&0&1\end{bmatrix}$

$$\Downarrow$$

$$\begin{bmatrix}1&-1&5&0&0&1\\0&6&-8&0&1&-2\\3&0&1&1&0&0\end{bmatrix}$$

Row operation: R_3-3R_1 — Elementary Matrix: $E_3=\begin{bmatrix}1&0&0\\0&1&0\\-3&0&1\end{bmatrix}$

$$\Downarrow$$

$$\begin{bmatrix}1&-1&5&0&0&1\\0&6&-8&0&1&-2\\0&3&-14&1&0&-3\end{bmatrix}$$

Row operation: $R_3-R_2/2$ — Elementary Matrix: $E_4=\begin{bmatrix}1&0&0\\0&1&0\\0&-\frac{1}{2}&1\end{bmatrix}$

$$\Downarrow$$

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

$$\begin{bmatrix} 1 & -1 & 5 & 0 & 0 & 1 \\ 0 & 6 & -8 & 0 & 1 & -2 \\ 0 & 0 & -10 & 1 & -\frac{1}{2} & -2 \end{bmatrix}$$

Row operation	Elementary Matrix
$R_3/(-10)$	$E_5 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{10} \end{bmatrix}$

$$\Downarrow$$

$$\begin{bmatrix} 1 & -1 & 5 & 0 & 0 & 1 \\ 0 & 6 & -8 & 0 & 1 & -2 \\ 0 & 0 & 1 & -\frac{1}{10} & \frac{1}{20} & \frac{1}{5} \end{bmatrix}$$

Row operations	Elementary Matrices
$R_1 - 5R_3$	$E_6 = \begin{bmatrix} 1 & 0 & -5 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$,
$R_2 + 8R_3$	$E_7 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 8 \\ 0 & 0 & 1 \end{bmatrix}$

$$\Downarrow$$

$$\begin{bmatrix} 1 & -1 & 0 & \frac{1}{2} & -\frac{1}{4} & 0 \\ 0 & 6 & 0 & -\frac{4}{5} & \frac{14}{10} & -\frac{2}{5} \\ 0 & 0 & 1 & -\frac{1}{10} & \frac{1}{20} & \frac{1}{5} \end{bmatrix}$$

Row operation	Elementary Matrix
$R_2/6$	$E_8 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{6} & 0 \\ 0 & 0 & 1 \end{bmatrix}$

$$\Downarrow$$

$$\begin{bmatrix} 1 & -1 & 0 & \frac{1}{2} & -\frac{1}{4} & 0 \\ 0 & 1 & 0 & -\frac{4}{30} & \frac{14}{60} & -\frac{2}{30} \\ 0 & 0 & 1 & -\frac{1}{10} & \frac{1}{20} & \frac{1}{5} \end{bmatrix}$$

Row operation	Elementary Matrix
$R_1 + R_2$	$E_9 = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

$$\Downarrow$$

$$\begin{bmatrix} 1 & 0 & 0 & \frac{11}{30} & \frac{1}{60} & -\frac{2}{30} \\ 0 & 1 & 0 & -\frac{4}{30} & \frac{14}{60} & -\frac{2}{30} \\ 0 & 0 & 1 & -\frac{1}{10} & \frac{1}{20} & \frac{1}{5} \end{bmatrix}$$

Then we have

$$A^{-1} = \frac{1}{60} \begin{bmatrix} 22 & 1 & -4 \\ -8 & 14 & -4 \\ -6 & 3 & 12 \end{bmatrix},$$

which can be written as the product of elementary matrices $E_9E_8E_7E_6E_5E_4E_3E_2E_1$. □

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

5. For what values of $\lambda \in \mathbb{R}$ is the following matrix

$$A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & 4 & 2 \\ 1 & -1 & \lambda \end{bmatrix}$$

invertible?

Solution: We use elimination to determine when the reduced row echelon form is an identity matrix. We have

$$\begin{aligned} A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & 4 & 2 \\ 1 & -1 & \lambda \end{bmatrix} &\xrightarrow{R_2 \leftrightarrow R_1} \begin{bmatrix} 2 & 4 & 2 \\ 3 & 0 & 1 \\ 1 & -1 & \lambda \end{bmatrix} \\ &\xrightarrow{R_1/2} \begin{bmatrix} 1 & 2 & 1 \\ 3 & 0 & 1 \\ 1 & -1 & \lambda \end{bmatrix} \\ &\xrightarrow[R_3 - R_1]{R_2 - 2R_1} \begin{bmatrix} 1 & 2 & 1 \\ 0 & -6 & -2 \\ 0 & -3 & \lambda - 1 \end{bmatrix} \\ &\xrightarrow{R_2/(-6)} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & \frac{1}{3} \\ 0 & -3 & \lambda - 1 \end{bmatrix} \\ &\xrightarrow{R_3 + 3R_2} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & \frac{1}{3} \\ 0 & 0 & \lambda \end{bmatrix}. \end{aligned}$$

The reduced row echelon form of A is I if and only if $\lambda \neq 0$. That is, A is invertible if and only if $\lambda \neq 0$.

□

6. Let A be an $n \times n$ matrix. If $A = \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_n \end{bmatrix}$ satisfies that $r_2 = r_3 + r_1$, is A invertible?

Solution: A is not invertible since using elementary eliminations to substract r_1 and r_3 from r_2 will result in a matrix with a zero row which is not invertible.

□

7. Let A be an $n \times n$ matrix. If $A = \begin{bmatrix} c_1 : c_2 : \cdots : c_n \end{bmatrix}$ satisfies that $c_2 = c_3 + c_1$, is A invertible?

Solution: By the previous question, A^T is not invertible and hence A is not invertible. □

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

8. Let v, $w \in \mathbb{R}^n$ be vectors. Is the matrix $A = \begin{bmatrix} \|v\| & 1 \\ |v \cdot w| & \|w\| \end{bmatrix}$ invertible?

Solution: A is not invertible if and only if $\|v\| \cdot \|w\| = |v \cdot w|$. □

9. Give an example of a 3×3 dominant matrix and find its inverse.

10. Find a sufficient condition on a, b, c and $d \in \mathbb{R}$ such that the matrix

$$A = \begin{bmatrix} a^2 + b^2 & 2ab \\ 2cd & c^2 + d^2 \end{bmatrix}$$

is invertible.

11. Let $A = \begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$. i) Compute A^2; ii) Show that for every $k \geq 3$, $k \in \mathbb{N}$, $A^k = 0$.

Solution: $A^2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ and $A^3 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$. Therefore for every $k \geq 3$, $k \in \mathbb{N}$, $A^k = A^3 A^{k-3} = 0$. . □

12. Let A be an $n \times n$ matrix. Show that if $A^k = 0$, then $I - A$ is invertible and

$$(I - A)^{-1} = I + A + A^2 + \cdots + A^{k-1}.$$

Solution: Hint: We have $(I - A)(I + A + A^2 + \cdots + A^{k-1}) = I$. □

13. Let A be an $n \times n$ matrix and $A = tI + N$, $t \in \mathbb{R}$ with $N^4 = 0$ for some $k \in \mathbb{N}$. Compute A^4 in terms of t and N.

Solution: Note that $tIN = N(tI)$. We have $A^4 = N^4 + (tI)N^3 + (tI)^2N^2 + (tI)^3N + (tI)^4 = tN^3 + t^2N^2 + t^3N + t^4I$. □

14. Let $D = \text{diag}\{\lambda_1, \lambda_2, \cdots, \lambda_n\}$ be a diagonal matrix with the main diagonal entries $\lambda_1, \lambda_2, \cdots, \lambda_n$. Show that D is invertible if and only if $\lambda_i \neq 0$, for every $i = 1, 2, \cdots, n$.

Solution: Hint: if $\lambda_i \neq 0$, for every $i = 1, 2, \cdots, n$, the reduced row echelon form of D is the identity matrix. □

15. Let A be an $n \times n$ matrix. i) If $A^3 = I$, find A^{-1}; ii) If $A^k = I$ for some $k \in \mathbb{N}$, find A^{-1}; iii) If $A^k = 0$ for some $k \in \mathbb{N}$, is it possible that A is invertible?

Solution: i) $A^{-1} = A^2$, ii) $A^{-1} = A^{k-1}$, iii). No. Notice that AB is invertible if and only if both A and B are invertible, where A and B are square matrices of the same size. □

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

16. Show that A is invertible if and only if A^k is invertible for every $k \in \mathbb{N}, k \geq 1$.

Solution: Notice that AB is invertible if and only if both A and B are invertible. The conclusion follows. □

17. Let A and B be $n \times n$ invertible matrices. i) Give an example to show that $A+B$ may not be invertible; ii) Show that $A+B$ is invertible if and only if $A^{-1} + B^{-1}$ is invertible.

Solution: i) Let $A = I$ and $B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ be 2×2 matrixes. Then A and B are both invertible while $A + B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ is not invertible.

ii). Note that $A + B = A(A^{-1} + B^{-1})B$. $A + B$ is invertible if and only if $A^{-1} + B^{-1}$ is invertible. □

2.4 *LU* decomposition

Exercise 2.4.1.

1. Let $-l_{ij}$ be the entry of the 4×4 E_{ij}^{-1} matrix below the main diagonal. Which one of the following products can be obtained by directly writing $-l_{ij}$ into the (i, j) position of the products? i) $E_{31}^{-1}E_{32}^{-1}E_{41}^{-1}E_{42}^{-1}E_{43}^{-1}$; ii) $E_{32}^{-1}E_{21}^{-1}E_{31}^{-1}E_{42}^{-1}E_{43}^{-1}$.

Solution: The point is that the row of the identity matrix to be added to another row should not the changed. i) Row 1, 2, 3 were not changed when they are used to change row 4 and row Row 1 and 2 were not changed when they are used to change row 3. So the matrix $E_{31}^{-1}E_{32}^{-1}E_{41}^{-1}E_{42}^{-1}E_{43}^{-1}$ can be obtained by directly writing $-l_{ij}$ into the (i, j) position of the products.

ii) Since row 2 were changed with E_{21}^{-1} before applying to E_{32}^{-1}, $E_{32}^{-1}E_{21}^{-1}E_{31}^{-1}E_{42}^{-1}E_{43}^{-1}$ can not be obtained by directly writing $-l_{ij}$ into the (i, j) position of the products.

□

2. Find the *LU* decomposition of

$$A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & 4 & 2 \\ 1 & -1 & 5 \end{bmatrix}.$$

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

Solution: Using elementary eliminations we have

$$E_{32}E_{31}E_{21}A = \begin{bmatrix} 3 & 0 & 1 \\ 0 & 4 & \frac{4}{3} \\ 0 & 0 & 5 \end{bmatrix} := U,$$

where

$$E_{21}\begin{bmatrix} 1 & 0 & 0 \\ -\frac{2}{3} & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_{31} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -\frac{1}{3} & 0 & 1 \end{bmatrix}, E_{32} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & \frac{1}{4} & 1 \end{bmatrix}.$$

Then we have $A = LU$, where

$$L = E_{21}^{-1}E_{31}^{-1}E_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ \frac{2}{3} & 1 & 0 \\ \frac{1}{3} & -\frac{1}{4} & 1 \end{bmatrix}.$$

□

3. Let $b = (1, 2, 3)$ and $A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & 4 & 2 \\ 1 & -1 & 5 \end{bmatrix}$. Use the LU decomposition of A to solve system $Ax = b$.

Solution: Since $A = LU$, we have $LUx = b$. Let $Ux = y$. Then $Ly = b$ with solution $y = (1, \frac{4}{3}, 3)$. Solving $Ux = y$ we have

$$x = \left(\frac{2}{15}, \frac{2}{15}, \frac{3}{5}\right).$$

□

4. Is it true that a matrix A does not have an LU decomposition? Justify your answer.

Solution: Yes, for example

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

□

2.5 Transpose and permutation

Exercise 2.5.1.

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

1. Let $A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix}$. Find A^{-1} and A^T.

Solution: A is a permutation matrix and is orthogonal.

$$A^{-1} = A^T = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

□

2. Let $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}$. i) Find AA^T and A^TA. ii) Determine which one of AA^T and A^TA is invertible. iii) If one of AA^T and A^TA is invertible, does it contradict Theorem 2.3.6?

Solution: i)

$$AA^T = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, A^TA = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

ii) A^TA is invertible.

iii) No. Theorem 2.3.6 is about square matrices. □

3. Let

$$A = \begin{bmatrix} 0 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 5 \\ 3 & 4 & 5 & 6 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 2 & 3 \\ 3 & 4 & 5 & 6 \\ 2 & 3 & 4 & 5 \end{bmatrix}.$$

i) Find a permutation matrix P_1 such that $B = P_1A$; ii) Find a permutation matrix P_2 such that $A = P_2B$. iii) Compute P_1P_2 and P_2P_1.

Solution: Examining how the rows of A were rearranged to obtain B, we have

$$P_1 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}, P_2 = P_1^T = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}, P_1P_2 = P_2P_1 = I.$$

□

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

4. Let

$$A = \begin{bmatrix} 7 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 5 \\ 3 & 4 & 5 & 7 \end{bmatrix}.$$

Find a permutation matrix P, a lower triangular matrix L and a diagonal matrix D such that $PA = LU$.

Solution:

$$L = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1/7 & 13/25 & 1 & 0 \\ 2/7 & 19/25 & 1/2 & 1 \\ 3/7 & 1 & 0 & 0 \end{bmatrix}, U = \begin{bmatrix} 7 & 1 & 2 & 3 \\ 0 & 25/7 & 29/7 & 40/7 \\ 0 & 0 & 14/25 & 3/5 \\ 0 & 0 & 0 & -1/2 \end{bmatrix},$$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

□

5. Let $R_\theta = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$. Show that R_θ is an orthogonal matrix.

Solution: R_θ is an orthogonal matrix since we have

$$R_\theta {R_\theta}^T = I.$$

□

6. Let $x \in \mathbb{R}^n$ with $x^T x = 1$. Define the **Householder matrix** by

$$H = I - 2xx^T.$$

i) Show that H is an orthogonal matrix; ii) Show that H is symmetric.

Solution: i) Verify by the definition that $H^T H = I$. Indeed, we have

$$\begin{aligned} H^T H &= (I - 2xx^T)^T (I - 2xx^T) \\ &= (I - 2xx^T)(I - 2xx^T) \\ &= (I - 2xx^T) - 2xx^T(I - 2xx^T) \\ &= (I - 2xx^T) - 2xx^T + 4xx^T xx^T \\ &= I - 4xx^T + 4xx^T \\ &= I. \end{aligned}$$

ii) $H^T = (I - 2xx^T)^T = H$. H is symmetric. □

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

7. Let $S = \begin{bmatrix} I & A \\ A^T & O \end{bmatrix}$, where I is $m \times m$ and A is $m \times n$, O the zero matrix. Find a block diagonal matrix D and block lower triangular matrix L such that

$$S = LDL^T.$$

Solution: Let $L = \begin{bmatrix} I & O \\ -A^T & I \end{bmatrix}$. We have

$$LS = \begin{bmatrix} I & O \\ -A^T & I \end{bmatrix} \begin{bmatrix} I & A \\ A^T & O \end{bmatrix} = \begin{bmatrix} I & A \\ O & -A^TA \end{bmatrix}.$$

Then

$$LSL^T = \begin{bmatrix} I & O \\ -A^T & I \end{bmatrix} \begin{bmatrix} I & A \\ A^T & O \end{bmatrix} \begin{bmatrix} I & -A \\ O & I \end{bmatrix} = \begin{bmatrix} I & O \\ O & -A^TA \end{bmatrix}.$$

□

8. Show that AA^T is invertible if and only if the rows of A are linearly independent.

Solution: We know that the columns of A are linearly independent if and only if A^TA is invertible. If follows that $(A^T)^TA^T$ is invertible if and only if the columns of A^T are linearly independent. That is, AA^T is invertible if and only if the rows of A are linearly independent. □

9. We say A is **skew-symmetric** if $A^T = -A$. i) Show that if A is a skew-symmetric $n \times n$ matrix then $a_{ii} = 0$ for every $i = 1, 2, \cdots, n$. ii) If A is both symmetric and skew-symmetric, then $A = 0$.

Solution: i) We have $A^T = -A$. Then it follows that for every $i = 1, 2, \cdots, n$,

$$(A^T)_{ii} = (A)_{ii} = (-A)_{ii} \Rightarrow a_{ii} = -a_{ii}.$$

We have $a_{ii} = 0$ for every $i = 1, 2, \cdots, n$.

ii) If A is both symmetric and skew-symmetric, then $A^T = -A = A$. That is, $A = 0$. □

10. Let A be an $n \times n$ matrix. Show that i) $A + A^T$ is symmetric; ii) $A - A^T$ is skew-symmetric; iii) For every square matrix B, there exist a unique symmetric matrix B_1 and a unique skew-symmetric matrix B_2 such that $B = B_1 + B_2$.

Solution: i) Since $(A + A^T)^T = A^T + A = A + A^T$, $A + A^T$ is symmetric.

ii) Since $(A - A^T)^T = A^T - A = -(A - A^T)$, $A - A^T$ is skew-symmetric. □

11. A matrix is called **lower triangular** if every entry above the main diagonal is zero and is called **upper triangular** if every entry below the main diagonal is zero. Let A be an $n \times n$ invertible matrix. i) Show that if A is lower

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

triangular, A^{-1} is also lower triangular; ii) Show that if A is upper triangular, then A^{-1} is also upper triangular.

Solution: i) We use matrix partition. Suppose A and $B = A^{-1}$ are partitioned as following:

$$A = \begin{bmatrix} A_{11} & O \\ A_{21} & a_{nn} \end{bmatrix}, B = A^{-1} = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & b_{nn} \end{bmatrix}.$$

Then we have

$$BA = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & b_{nn} \end{bmatrix} \begin{bmatrix} A_{11} & O \\ A_{21} & a_{nn} \end{bmatrix} = \begin{bmatrix} B_{11}A_{11} + B_{12}A_{21} & B_{12}a_{nn} \\ B_{21}A_{11} + b_{22}A_{21} & b_{nn}a_{nn} \end{bmatrix} = I_n,$$

where I_n is the $n \times n$ identity matrix. Since $a_{nn} \neq 0$, we have $B_{12} = O$. It follows that $B_{11}A_{11} = I_{n-1}$. By the same token, we can show that right upper block of B_{11} is zero. Repeat the same argument on the sub-martrices of B_{11}, we obtain that B is lower triangular.

ii) Notice that if A is invertible we have $(A^{-1})^T = (A^T)^{-1}$. The statement follows from i). □

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual

Concise Introduction to Linear Algebra 1st edition Hu Solutions Manual