MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

SKU: 9781498755672-SOLUTIONS

Solutions Manual for A MATLAB Companion to Complex Variables.

Chapter 1

SECTION 1.1

```
SEC 1.1
Problem 1
format short
solution='PROBLEM 1'
a=(3-4i)/(1+2i)+2i
b=(2+i)/(3+5i)-1-i/2
c=(1+2*i/3)^3-i/(3-4i)^3
d=(1+2/(3i))^11
e=(2i/(3+4i)+conj(1/(5-4i)))^5
```

OUTPUT FROM THE ABOVE PROGRAM

solution =

PROBLEM 1

a =

-1

b =

-0.6765 - 0.7059i

c =

-0.3305 + 1.7112i

d =

7.4284 - 1.3889i

e =

0.0003 + 0.0216i

solution =

1

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

```
solution='PROBLEM 2'
%note that (1+i)^2=2i and (2i)^18=-2^18 since i raised to an even
%integer
%is plus or minus one;the minus sign holds if the integer not divisible
%by4

%so, without matlab, the  final answer is -2^18*(1+i)=-2^18-i2^18=
%-262144-i262144
%provided you get 2^18 off a pocket calculator or you can notice that
%2^18=8*32^3

format long
(1+i)^37

solution='PROBLEM 3a'
%part a
sa=0;
for n=0:10
    z=(i/2)^n;
    sa=sa+z;
end
sa

solution='PROBLEM 3b'
%part b
sb=(1-(i/2)^11)/(1-i/2)
%EXERCISES SEC 1.1
format short
solution='PROBLEM 1'
a=(3-4i)/(1+2i)+2i
b=(2+i)/(3+5i)-1-i/2
c=(1+2*i/3)^3-i/(3-4i)^3
d=(1+2/(3i))^11
e=(2i/(3+4i)+conj(1/(5-4i)))^5
```

PROBLEM 2

Without MATLAB

$$(1+i)^{37} = (1+i)^{36}(1+i) = \left((1+i)^2\right)^{18}(1+i) = 2^{18} i^{18}(1+i)$$

$$= -2^{18}(1+i) = -262144 - i262144$$

You can compute $2^{18} = 32*32*32*8 = 262144$

With MATLAB

>> format long

>> (1+i)^37

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

ans =

-2.621440000000000e+05 - 2.621440000000000e+05i

```
3.

solution='PROBLEM 3'
%part a
sa=0;
for n=0:10
    z=(i/2)^n;
    sa=sa+z;
end
sa

solution='PROBLEM 3b'
%part b
sb=(1-(i/2)^11)/(1-i/2)
```

3.

PROBLEM 3a

```
%problem 3 section1.1
format long
s=1;
for n=1:10
    s=s+(i/2)^n;
end
s
```

WHOSE OUTPUT IS

sa =

0.79980468750000 + 0.40039062500000i

solution =

PROBLEM 3b

>> format long
>> (1-(i/2)^11)/(1-i/2)

ans =

0.799804687500000 + 0.400390625000000i

3

SECTION 1.2

Problem 1.

```
%PROBLEMS 1 %solution='problem 1'
%parta
a=(1+i)*(sqrt(3)+i)
angle_product=angle(a)
angle_first=angle(1+i)
angle_second=angle(sqrt(3)+i)
angle_sum=angle_first+angle_second
%partbb=(1+i)*(-sqrt(3)+i)
angle_product=angle(b)
angle_first=angle(1+i)
angle_second=angle(-sqrt(3)+i)
angle_sum=angle_first+angle_second
```

OUTPUT FROM THE ABOVE PROGRAM

solution =

problem 1

a =

0.73205080756888 + 2.73205080756888i

angle_product =

1.30899693899575

angle_first =

0.78539816339745

angle_second =

0.52359877559830

angle_sum =

1.30899693899575

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

b =

-2.73205080756888 - 0.73205080756888i

angle_product =

-2.87979326579064

angle_first =

0.78539816339745

angle_second =

2.61799387799149

angle_sum =

3.40339204138894

```
% explanation of discrepancy;the principal angle of (1+i) is pi/4 while
%the
% principal angle of (-sqrt(3)+i)is 5pi/6. Their sum is 13pi/12 which
%is
% not a principal angle as it exceeds pi. MATLAB yields the principal
%angle of the product
% which is -11pi/12.
```

PROBLEM 2

```
%PROBLEM 2

%solution ='problem 2'
%parta
a=angle(cos(197)+i*sin(197))
b=angle(1/(4*cos(79)+4*i*sin(79)))
c=angle((1+2*i)^27)
theta=53.3*pi/180;
d= angle((cos(theta)+i*sin(theta))^8);
d=d/pi*180
```

OUTPUT OF THE ABOVE

5

a =

2.22125547743282

b =

2.68140899333462

c =

-1.52291115545749

d =

66.39999999999999

PROBLEM 3

part(a)

>> [x,y]=pol2cart(-207,3)

x =

2.82310952232897

y =

1.01491508262293

part(b)

[x,y]=pol2cart(50,4)

6

x =

3.85986411396845

y =

-1.04949941481572

```
4. function [x,y]=pold2cart(th,r);
%a function that will convert the polar form of a complex number to the
%rectangular form,the input angle is in degrees, x and y are real and
% imag parts
x=r*(cos(th*pi/180))
y=r*sin(th*pi/180)
clear

a)
>> pold2cart(5,60)

x =

   2.500000000000000


y =

   4.330127018922193

b)
>> pold2cart(2,-320)

x =

   1.532088886237956


y =

   1.285575219373079

>>
```

SECTION 1.3

PROBLEM 1

```
>> %part a
>> (-27*i)^(2/3)

ans =
```


MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

```
   4.5000 - 7.7942i

>> %partb
>> (1+i)^(1/15)

ans =

   1.0220 + 0.0536i

>> %partc
>> (-1-i)^(1/9)

ans =

   1.0038 - 0.2690i
```

PROBLEM 2

a) $(i^{1/2})^2 = i$ must be true since $z^{1/m}$ is defined in such a way that $\left(z^{1/m}\right)^m$ will always yield *z when m is a nonzero integer*. Verify this using De Moivre's Theorem.

b) MATLAB will compute $i^{3/2}$ as $\left(\sqrt{1}\right)^3 cis(\frac{3\pi}{4}) =$ .Raising this to the 2/3 power using the MATLAB convention(we have $\left(\left(\sqrt{1}\right)^3 cis(\frac{3\pi}{4})\right)^{2/3} = cis(\frac{\pi}{2}) = i,$ since $3\pi/4$ is a principle angle.

c) MATLAB will compute $-1^{4/3}$ as $\left(\sqrt[3]{1}\right)^4 cis(\frac{4\pi}{3}) = \frac{-1}{2} - i\frac{\sqrt{3}}{2}$ .Raising this to the 3/4 power using the MATLAB convention (the principle angle is $-2\pi/3$) we have $\left(cis(\frac{-2\pi}{3})\right)^{3/4} = cis(\frac{-\pi}{2}) = -i,$. We do not recover -1.

d) MATLAB will compute $-1^{3/4}$ as $\left(\sqrt[4]{1}\right)^3 cis(\frac{3\pi}{4}) = \frac{-1}{\sqrt{2}} + i\frac{\sqrt{1}}{2}$ .Raising this to the 4/3 power using the MATLAB convention (the principle angle is $3\pi/4$) we have $\left(cis(\frac{3\pi}{4})\right)^{4/3} = cis(\pi) = -1,$. We do recover -1.

e) MATLAB will compute $(-1+i)^{5/3}$ as $\left(\sqrt[3]{\sqrt{2}}\right)^5 cis(\frac{3\pi}{4}\frac{5}{3}) = \left(\sqrt[3]{\sqrt{2}}\right)^5 cis(\frac{5\pi}{4})$ .Raising this to the 3/5 power using the MATLAB convention (the principle angle is $-3\pi/4$) we have $\left(\left(\sqrt[3]{\sqrt{2}}\right)^5 cis(\frac{-3\pi}{4})\right)^{3/5} = \sqrt{2}cis(\frac{-9\pi}{20}) =$ 0.2212 - 1.3968i . We do not recover -1+i.

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

```
f) verification of part e) above
>> a=(-1+i)^(5/3)

a =

 -1.259921049894873 - 1.259921049894873i

>> a^(3/5)

ans =

  0.221231742082474 - 1.396802246667421i (which checks)
```

3.a)

```
>> coeffs=zeros(1,6);
>> coeffs(1)=1;
>> coeffs(6)=-1-i;
>> roots(coeffs)

ans =


  -0.9550 + 0.4866i
  -0.7579 - 0.7579i
   0.1677 + 1.0586i
   0.4866 - 0.9550i
   1.0586 + 0.1677i


>> ans.^24

ans =

   1.6310 + 5.0197i
   5.2780 - 0.0000i
  -4.2700 + 3.1023i
   1.6310 - 5.0197i
  -4.2700 - 3.1023i
```


MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual



The above is obtained if we use the line of code plot(ans,'*')

b)

```
>> coeffs=zeros(1,8);
>> coeffs(1)=1;
>> coeffs(8)=-3+4*i;
>> roots(coeffs)

ans =

  -0.4396 - 1.1792i
  -1.1961 - 0.3915i
  -1.0518 + 0.6910i
  -0.1155 + 1.2532i
   0.6478 - 1.0790i
   0.9077 + 0.8717i
   1.2475 - 0.1662i

>> ans.^5


ans =

  -3.0852 + 0.6691i
   0.0342 - 3.1567i
```


MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

3.0700 + 0.7358i
-1.4005 + 2.8293i
1.3388 + 2.8590i
-2.4467 - 1.9950i
2.4894 - 1.9414i

c) >> coeffs=zeros(1,100);
>> coeffs(1)=1;coeffs(100)=-13-7*i;
>> a=roots(coeffs);
>> b=a.^100;
>> plot(b,'*');axis equal;grid



The 99 values of $(13+7i)^{100/99}$.

PROBLEM 4

a) Refer to the code written for Example 2 and add an additional line
`sum_of_roots=sum(w)`
which will give you the sum of all the values of $z^{n/m}$.

b) sum_of_roots =

-1.143529715363911e-014 +6.661338147750939e-016i
The preceding is about 14 orders of magnitude less than any of the individual roots.

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

Had there been no rounding error we would have gotten zero.

b)

```
>> coeffs=zeros(1,8);
>> coeffs(1)=1;coeffs(8)=-1-i;
>> a=roots(coeffs);
>> b=a.^11;
>> sum(b)

ans =

   1.0991e-14 - 1.7319e-14i
```

Note that this result is small but not zero.

Problem 5

```
clf; clear
x=linspace (-3, 3, 301);
w=(x-1).^(2/5);
figure(1)
plot(x,w);grid;
figure(2)
plot(x, real(w),'-',x,imag(w),'*')
grid;figure(2)
```

The above results in these 2 figures:

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual




Figure 1

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual




Figure 2 (Note that real part is in solid line while imag part consists of *

Explanation:

Figure 1 above is a plot of the real part of $(z-1)^{2/5}$ where z = x satisfies -3 $\leq$ x $\leq$3. Notice that the function of z has a branch cut at z = 1 . The principle value of this function is $f(z)=\left(\sqrt[5]{|z-1|}\right)^2 \angle\left(\frac{2}{5}\theta\right)$ where $-\pi<\arg(z-1)\leq\pi$ and $\arg(z-1)=\theta$.

If $z=x>1,$ then $\arg(z\text{-}1)=0$ and for $x>1$ our plot is simply a graph of $f(z)=\left(\sqrt[5]{|x-1|}\right)^2$ for $x>1$. If $z=x<1,$ then $\arg(z\text{-}1)=\pi$ and $w=f(z)=\left(\sqrt[5]{|x-1|}\right)^2[\cos(2\pi/5)+i\sin(2\pi/5)]$. Now MATLAB *plots only the real part* of this because of the line of code: `plot(x,w);grid;`

We obtain a graph of $f(x)=\left(\sqrt[5]{|x-1|}\right)^2[\cos(2\pi/5)]\approx\left(\sqrt[5]{|x-1|}\right)^2(.309)$ for x<1. Study the graphs in Figure 1 and you'll see that for values of x equally displaced on either side of x = 1 that the plot to the left of x=1 achieves a height that is about .31 times the height on the right.

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

Turn now to Figure 2. If $z = x \geq 1$, the function f(z) is real, and a plot of the real part reveals the behavior of the function. If $z = x \leq 1$ then as we have seen above the real part of *f(z)* when *z=x* is $f(x) \approx \left(\sqrt[5]{|x-1|}\right)^2 (.309)$ and the imaginary part is $f(z) = \left(\sqrt[5]{|x-1|}\right)^2 [i \sin(2\pi/5)] \approx \left(\sqrt[5]{|x-1|}\right)^2 (.9511)$. The imaginary part of *w*, for $-3 < x < 3$ appears in the heavy* line in Figure 2.

PROBLEM 6

```
z=input('the complex value of z to be considered')
m=input('input the positive integer m')
n=input('the integer n')
v=zeros(1,m+1);%creates a row vector length m+1
%elements are zero.
v(1,1)=1;% the first coeff is 1, the coeff of w^m
v(1,m+1)=-z;% the constant term (the last coeff) is -z
%the following solves the equation w^m-z=0 for w=z^(1/m)
the_roots =roots(v);
final_answer=the_roots.^n
```

SECTION 1.4

PROBLEM 1

```
%sec1.4_prob1
syms x y real
w=1/x+1/(i*y)+i/((x-1)+i*y)^2;
u=real(w)
v=imag(w)
'pretty answers'
'u'
pretty(u)
'v'
pretty (v)
 u=subs(u,[x y],[1 1])
 v=subs(v,[x y],[1 1])
```

OUTPUT IS
>> sec1_4prob1

u =

1/x + (2*y*(x - 1))/(((x - 1)^2 - y^2)^2 + 4*y^2*(x - 1)^2)

v =

((x - 1)^2 - y^2)/(((x - 1)^2 - y^2)^2 + 4*y^2*(x - 1)^2) - 1/y

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

ans =

pretty answers

ans =

u

$$\frac{1}{x}+\frac{2y(x-1)}{((x-1)^2-y^2)^2+4y^2(x-1)^2}$$

ans =

v

$$\frac{(x-1)-y}{((x-1)^2-y^2)^2+4y^2(x-1)^2}-\frac{1}{y}$$

u =

1

v =

-2

>>

PROBLEM 2

```
2 %sec1_4problem 2
syms r1 r2 x1 x2 real
Z=1/(1/(r1+i*x1)+1/(r2+i*x2))
r=real(Z)
x=imag(Z)
 R=subs(r,[r1 r2 x1 x2],[2 2 4 4])
 X=subs(x,[r1 r2 x1 x2],[2 2 4 4])
 R=subs(r, [r1 r2 x1 x2],[2 2 4 -4])
 X=subs(x, [r1 r2 x1 x2],[2 2 4 -4])

WHOSE OUTPUT IS
```


MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

```
>> sec1_4prob2

Z =

1/(1/(r1 + x1*i) + 1/(r2 + x2*i))


r =

(r1/(r1^2 + x1^2) + r2/(r2^2 + x2^2))/((r1/(r1^2 + x1^2) +
r2/(r2^2 + x2^2))^2 + (x1/(r1^2 + x1^2) + x2/(r2^2 +
x2^2))^2)


x =

(x1/(r1^2 + x1^2) + x2/(r2^2 + x2^2))/((r1/(r1^2 + x1^2) +
r2/(r2^2 + x2^2))^2 + (x1/(r1^2 + x1^2) + x2/(r2^2 +
x2^2))^2)


R =

    1.0000


X =

    2.0000


R =

    5.0000


X =

     0

>>
```

PROBLEM 3

3.
```
%sec1.4_prob3
syms x y real
w=1/(x+i*y)^3;
u=real(w)
v=imag(w)
```


MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

```
'pretty answers'
'u'
pretty(u)
'v'
pretty (v)
```

OUTPUT IS

>> sec1_4prob3

u =

(x*(x^2 - y^2) - 2*x*y^2)/((x*(x^2 - y^2) - 2*x*y^2)^2 + (y*(x^2 - y^2) + 2*x^2*y)^2)

v =

-(y*(x^2 - y^2) + 2*x^2*y)/((x*(x^2 - y^2) - 2*x*y^2)^2 + (y*(x^2 - y^2) + 2*x^2*y)^2)

ans =

pretty answers

ans =

u

$$\frac{x(x^2-y^2)-2xy^2}{(x(x^2-y^2)-2xy^2)^2+(y(x^2-y^2)+2x^2y)^2}$$

ans=
v

$$-\frac{y(x^2-y^2)+2x^2y}{(x(x^2-y^2)-2xy^2)^2+(y(x^2-y^2)+2x^2y)^2}$$

PROBLEM 4

```
%sec1.4_prob4
syms x y real
w=((x-i*y)/(x+i*y))^6;
u=real(w);
u=simplify(u)
v=imag(w);
v=simplify(v)
mag2=u^2+v^2;
'simplified u^2+v^2'
```


MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

```
simplify(mag2)
```

OUTPUT IS
>> sec1_4prob4

u =

1 - (72*x^10*y^2 - 480*x^8*y^4 + 944*x^6*y^6 - 480*x^4*y^8 + 72*x^2*y^10)/(x^2 + y^2)^6

v =

-(4*x*y*(x^2 - 3*y^2)*(3*x^2 - y^2)*(x^6 - 15*x^4*y^2 + 15*x^2*y^4 - y^6))/(x^2 + y^2)^6

ans =

simplified u^2+v^2

ans =

1

The reason we must get one is this
$\left|\frac{\overline{z}}{z}\right| = 1,$ thus $\left|\left(\frac{\overline{z}}{z}\right)^6\right| = 1$ which means that $\sqrt{u^2+v^2} = \left|\left(\frac{\overline{z}}{z}\right)^6\right| = 1$

5.

```
%sec1.4_prob5
syms x y real
w=(x+i*y+1/(x+i*y))^8
u=real(w);
u=simplify(u)
v=imag(w);
v=simplify(v)
u1=subs(u,[x y],[0,1])
v1=subs(v,[x y],[0,1])
```

OUTPUT IS
u =

120*x^2*y^4 - 168*x^2*y^2 - 120*x^4*y^2 - 28*x^2*y^6 + 70*x^4*y^4 - 28*x^6*y^2 - (112*y^2 - 28)/(x^2 + y^2)^2 - (224*y^2 - 8)/(x^2 + y^2)^3 + 56/(x^2 + y^2) + (224*y^4 - 144*y^2 + 1)/(x^2 +

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

y^2)^4 - (256*y^6)/(x^2 + y^2)^7 + (128*y^8)/(x^2 + y^2)^8 + 56*x^2 + 28*x^4 + 8*x^6 + x^8 - 56*y^2 + 28*y^4 - 8*y^6 + y^8 + (32*y^2*(12*y^2 - 1))/(x^2 + y^2)^5 - (32*y^4*(8*y^2 - 5))/(x^2 + y^2)^6 + 70

v =

(8*x*y*(x^2 + y^2 - 1)*(x^2 + y^2 + 1)*(x^3 + x^2*y + x*y^2 + x + y^3 - y)*(x^3 - x^2*y + x*y^2 + x - y^3 + y)*(x^6 - 2*x^5*y + x^4*y^2 + 2*x^4 - 4*x^3*y^3 - x^2*y^4 + 4*x^2*y^2 + x^2 - 2*x*y^5 + 2*x*y - y^6 + 2*y^4 - y^2)*(x^6 + 2*x^5*y + x^4*y^2 + 2*x^4 + 4*x^3*y^3 - x^2*y^4 + 4*x^2*y^2 + x^2 + 2*x*y^5 - 2*x*y - y^6 + 2*y^4 - y^2))/(x^2 + y^2)^8

u1 =

0

v1 =

0

Notice that at z=0 y=1 that $z=i$. .Thus (z+1/z) =0 at z=i and the real and imaginary parts of the given expression must vanish.

6.
```
a=sym(sqrt(2));
b=sym(sqrt(3));
c=(a+i*b)^17
c1=expand(c)
cr=real(c1)
ci=imag(c1)
 % this is for part b below
d=1/(a-i*b)^13
d1=expand(d)
dr=real(d1)
di=imag(d1)
```

The numerical answers
c =

(2^(1/2) + 3^(1/2)*1i)^17

c1 =

- 493679*2^(1/2) + 3^(1/2)*303041i

cr =

-493679*2^(1/2)

ci =

303041*3^(1/2)

MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual

d =

1/(2^(1/2) - 3^(1/2)*1i)^13

d1 =

1/(12349*2^(1/2) + 3^(1/2)*17471i)

dr =

(12349*2^(1/2))/1220703125

di =

-(17471*3^(1/2))/1220703125

Thus the answers to part a) are cr and ci above
And the answers to part b) are dr and di above.

CHAPTER 2

SECTION 2.1

Note: in the following codes, you can insert the command `colormap(gray)` after using **mesh, meshz** or **surf** in order to obtain plots in shades of gray, with no coloring. For the most part we provide color solutions here.

PROBLEM 1

```
clf
%probl 2.1 number 1
x=linspace(.5,2,50);
y=x;
[x,y]=meshgrid(x,y);
z=x+i*y;
Z=z+1./z;
figure(1)
mesh(x,y,Z);xlabel('x');ylabel('y');title('using mesh');
figure(2)
 meshz(x,y,Z);xlabel('x');ylabel('y');title('using meshz');
 figure(3)
surf(x,y,Z);xlabel('x');ylabel('y');title('using surf');
% view(-50,30); %try this view for part d
```


MatLab Companion to Complex Variables 1st edition Wunsch Solutions Manual